

## WELL-POSEDNESS AND STATIONARY SOLUTIONS

MARTIN BURNS<sup>1</sup> AND MICHAEL GRINFELD<sup>2</sup>

<sup>1</sup>Department of Mathematics and Statistics, University of Strathclyde  
Glasgow G1 1XH, UK  
*E-mail:* martin.f.burns@strath.ac.uk

<sup>2</sup>Department of Mathematics and Statistics, University of Strathclyde  
Glasgow G1 1XH, UK  
*E-mail:* m.grinfeld@strath.ac.uk

*Dedicated to Professor Jeff Webb on the occasion of his retirement.*

**ABSTRACT.** In this paper we prove existence and uniqueness of variational inequality solutions for a bistable quasilinear parabolic equation arising in the theory of solid-solid phase transitions and discuss its stationary solutions, which can be discontinuous.

**AMS (MOS) Subject Classification.** 35B32, 35D30, 35J62, 35K59

### 1. INTRODUCTION

To generalise the Ginzburg Landau phase transition theory to high gradients in the order parameter ( $u$ ), Rosenau [7, 8] proposed the following free energy functional:

$$E[u](t) = \int_{\Omega} [W(u) + \epsilon\Psi(|\nabla u|)] dx, \quad (1.1)$$

where the diffusion coefficient  $\epsilon > 0$ , the interface energy  $\Psi(s)$  is a convex function of its variable that grows linearly in  $s$ ; for example, below we take

$$\Psi(s) = \sqrt{1 + s^2} - 1,$$

$W(u)$  is the bulk energy, which we take to be a double well one, and fix

$$W(u) = \frac{u^4}{4} - \frac{u^2}{2}.$$

The formal  $L^2$  gradient flow of (1.1) is

$$u_t = \epsilon \nabla \cdot (\psi(\nabla u)) + f(u), \quad (1.2)$$

where  $f(u) = -W'(u) := u - u^3$ ,

$$\psi(\nabla u) = \frac{\nabla u}{\sqrt{1 + |\nabla u|^2}},$$

and  $(x, t) \in \Omega \times (0, T) \equiv Q_T$  for some bounded domain  $\Omega \subset \mathbb{R}^n$ ,  $T > 0$ . (1.2) has to be supplemented with suitable initial and boundary conditions; here we consider the physically relevant Neumann boundary conditions,  $\psi(\nabla u) \cdot \underline{n} = 0$  on  $\partial\Omega$  which, since  $\psi(0) = 0$ , implies that  $\nabla u \cdot n = 0$  on  $\partial\Omega$ .

In this paper, using the methods of [5] we prove a well-posedness result for (1.2) Although this result holds for any dimension  $n$ , here we restrict ourselves to the one-dimensional case  $\Omega \equiv (0, L)$ ,  $L > 0$ . As shown in [2], the bifurcation structure for the stationary problem associated with (1.2) depends on the parameter  $\epsilon$  as well as the length  $L$  of the interval; these issues will be discussed in more detail in Section 4.

## 2. PRELIMINARIES

In this section we briefly recall some properties of the function space  $BV(\Omega)$ . A function of bounded variation is a  $u \in L^1(\Omega)$  whose partial derivatives in the sense of distributions are measures with finite total variation

$$\int_{\Omega} |u_x| dx = \sup \left\{ \int_{\Omega} u v_x dx : v \in C_0^{\infty}(\Omega), |v(x)| \leq 1 \text{ for } x \in \Omega \right\}$$

The space  $BV(\Omega)$  endowed with the norm

$$\|u\|_{BV} = \|u\|_{L^1(\Omega)} + \int_{\Omega} |u_x| dx$$

is a Banach space. The topology on  $BV$  which we will require is the  $BV$ -weak\* topology defined by

$$u_j \xrightarrow{BV-w^*} u \Leftrightarrow u_j \rightarrow u \text{ in } L^1(\Omega) \text{ and } u_{jx} \rightharpoonup u_x \text{ in } M(\Omega)$$

where  $M(\Omega)$  is the space of bounded measures on  $\Omega$  and  $u_{jx} \rightharpoonup u_x$  in  $M(\Omega)$  means that

$$\int_{\Omega} u_{jx} \varphi dx \rightarrow \int_{\Omega} u_x \varphi dx$$

for all  $\varphi \in C_0(\Omega)$ .

We also have the following compactness property: for every bounded sequence  $\{u_j\}$ , there exists a subsequence  $\{u_{j_k}\}$  and a function  $u$  in  $BV(\Omega)$  such that  $u_{j_k} \xrightarrow{BV-w^*} u$ .

Following [6], we define  $\int_{\Omega} \Psi(u_x)$  and if  $\Psi(s) = \sqrt{1 + s^2}$  we arrive at the following definition

$$\int_{\Omega} \sqrt{1 + |u_x|^2} dx = \sup_{v \in C_0^{\infty}} \left\{ - \int_{\Omega} u v_x dx + \int_{\Omega} \sqrt{1 - v^2} dx : |v(x)| \leq 1 \forall x \in \Omega \right\}.$$

Hence we obtain the following useful estimate:

$$\int_{\Omega} |u_x| dx - |\Omega| \leq \int_{\Omega} \sqrt{1 + |u_x|^2} - 1 dx \leq \int_{\Omega} |u_x| dx + |\Omega| \quad (2.1)$$

for all  $u \in BV(\Omega)$ .

### 3. Existence and uniqueness of weak solutions of the parabolic problem

The problem we are considering is

$$\begin{aligned} u_t &= (\psi(u_x))_x + f(u), & (x, t) \in Q_T \equiv \Omega \times (0, T), \\ u_x(0, t) &= u_x(L, t) = 0, & (x, t) \in \partial\Omega \times (0, T), \\ u(x, 0) &= u_0(x), & x \in \Omega, \end{aligned} \quad (3.1)$$

where  $(0, T)$  is any finite time interval over which we will prove existence of solutions. Note that without loss of generality we have put  $\epsilon = 1$ .

First, we need to define our notion of a weak solution. To begin with, let us suppose that  $u$  is smooth enough so that we can justify the calculations which follow. For smooth test functions  $v \in C^\infty(Q_T)$ , we multiply our equation by  $v - u$  and integrate by parts using Neumann boundary conditions to obtain

$$\int_{Q_T} (u_t - f(u))(v - u) dx dt + \int_{Q_T} \psi(u_x)(v_x - u_x) dx dt = 0. \quad (3.2)$$

Since  $\Psi(s)$  is convex, we have that  $\Psi(v_x) - \Psi(u_x) \geq \Psi'(u_x)(v_x - u_x)$  and hence

$$\int_{Q_T} (u_t - f(u))(v - u) dx dt + \int_{Q_T} (\Psi(v_x) - \Psi(u_x)) dx dt \geq 0,$$

for smooth functions  $v \in C^\infty(Q_T)$ . This motivates the following definition of a weak solution to our problem.

**Definition 3.1.** *Let  $M(Q_T)$  denote the space of bounded measures on  $Q_T$ . A function  $u \in L^\infty(Q_T) \cap L^\infty((0, T), BV(\Omega)) \cap \{u : u_x \in M(Q_T)\}$  is called a weak solution of problem (3.1) if  $u_t \in L^2(Q_T)$  and  $u$  satisfies the variational inequality*

$$\int_{Q_T} (u_t - f(u))(v - u) dx dt + \int_{Q_T} (\Psi(v_x) - \Psi(u_x)) dx dt \geq 0 \quad (3.3)$$

for all  $v \in L^\infty(Q_T) \cap \{v : v_x \in M(Q_T)\}$ .

(Thus  $v_x$ , the distributional derivative of the function  $v$ , will be a measure with finite total variation.)

By the above discussion, classical solutions of (3.1) automatically satisfy variational inequality (3.3). To see that a smooth solution of (3.3) also satisfies (3.1), choose as a test function  $v = u + ch$  where  $h \in C^\infty$ ,  $c \in \mathbb{R}$ , so that (3.3) becomes

$$\int_{Q_T} (u_t - f(u))(ch) dx dt + \int_{Q_T} \Psi(u_x + ch_x) dx dt \geq \int_{Q_T} \Psi(u_x) dx dt.$$

Hence from the Taylor series of  $\Psi(u_x + ch_x)$  we have

$$c \int_{Q_T} (u_t - f(u))h dx dt + c \int_{Q_T} \Psi'(u_x)h_x dx dt + \frac{c^2}{2} \int_{Q_T} \Psi''(u_x)(h_x)^2 + \dots \geq 0.$$

Considering firstly,  $c > 0$ , then  $c < 0$  and letting  $c \rightarrow 0$  from above and below yields

$$\int_{Q_T} (u_t - f(u))h \, dx \, dt + \int_{Q_T} \psi(u_x)h_x \, dx \, dt = 0, \quad \forall h \in C_0^\infty(Q_T).$$

Integrating by parts and using the boundary conditions, we see that  $u$  classically satisfies (3.1).

**Theorem 3.2.** *The problem (3.1) admits a unique variational inequality solution for all  $T > 0$  for every  $u_0(x) \in BV(\Omega)$ .*

For  $\gamma > 0$ , consider the following regularised problem:

$$\begin{aligned} u_t &= (\psi(u_x))_x + f(u), & (x, t) &\in \Omega \times (0, T), \\ u_x(0, t) &= u_x(L, t) = 0, & (x, t) &\in \partial\Omega \times (0, T), \\ u(x, 0) &= u_0^\gamma(x), & x &\in \Omega, \end{aligned}$$

where  $u_0^\gamma(x)$  satisfy

$$\begin{aligned} u_0^\gamma &\in C^\infty(\bar{\Omega}), & u_{0x}^\gamma &= 0 \text{ on } \partial\Omega, \\ \|u_0^\gamma - u_0\|_{L^\infty(\Omega)} &\rightarrow 0 \text{ as } \gamma \rightarrow 0, & \|u_0^\gamma\|_{L^\infty(\Omega)} &\leq \|u_0\|_{L^\infty(\Omega)} + 1 = m_0, \\ \text{and } \int_{\Omega} |u_{0x}^\gamma| \, dx &\leq C(\Omega) \int_{\Omega} |u_{0x}| \, dx. \end{aligned}$$

The existence of such a sequence of regularising initial data  $u_0^\gamma \in C^\infty(\Omega)$  follows from the fact the initial data  $u_0 \in BV(\Omega)$  and because the space  $C^\infty(\Omega)$  is dense in the space of functions of bounded variation. Let  $u^\gamma(x, t)$  represent the unique classical solution to the regularised problem arising from the regularising initial data  $u_0^\gamma(x)$ ; these exist by standard parabolic theory. We want to show that there exists a limit  $u \in BV(\Omega)$  of  $u^\gamma$  in  $L^1(Q_T)$  as  $\gamma \rightarrow 0$ , which will be a weak solution to our problem and that it does not depend on the choice of the sequence  $u^\gamma$ . As in [5], we will need to establish a series of convergence properties for, and a priori bounds on, the approximating solutions  $u^\gamma$ . Namely we show

**Lemma 3.3.**

- A: *the sequence  $\{u^\gamma\}$  is uniformly bounded in  $L^\infty(Q_T)$  and the sequence  $\{u_t^\gamma\}$  is uniformly bounded in  $L^2(Q_T)$*
- B: *the sequence  $\{u^\gamma\}$  is uniformly bounded in  $BV(Q_T)$  and in  $L^\infty((0, T), BV(\Omega))$*
- C: *the sequence  $\{u^\gamma\}$  converges in the space  $L^\infty((0, T), L^2(\Omega))$  and the sequence  $\{u^\gamma(t, \cdot)\}$  converges in the space  $L^2(\Omega)$  for all  $t \in [0, T]$ .*

*Proof.* [A]: In what follows, let  $Q_\tau$  denote the space-time cylinder  $\Omega \times (0, \tau)$  where  $\tau$  is arbitrary in  $[0, T]$ . First of all, we have that

$$\|u^\gamma\|_{L^\infty(Q_T)} < m_0, \tag{3.4}$$

where  $m_0 > 1$ , by the parabolic maximum principle and properties of  $f(\cdot)$ .

We show next that the sequence  $\{u_t^\gamma\}$  is uniformly bounded in  $L^2(Q_T)$ . Multiply the regularised problem by  $u_t^\gamma$  and integrate over  $Q_\tau$ :

$$\begin{aligned} \int_{Q_\tau} (u_t^\gamma)^2 dx dt &= - \int_{Q_\tau} \psi(u_x^\gamma) u_{tx}^\gamma dx dt + \int_{Q_\tau} f(u^\gamma) u_t^\gamma dx dt \\ &= - \int_0^\tau \frac{d}{dt} \int_\Omega \Psi(u_x^\gamma) dx dt + \int_0^\tau \frac{d}{dt} \int_\Omega F(u^\gamma) dx dt \\ &= - \int_\Omega (\Psi(u_x^\gamma)|_{t=\tau} - \Psi(u_0^\gamma)) dx + \int_\Omega (F(u^\gamma)|_{t=\tau} - F(u_0^\gamma)) dx, \end{aligned}$$

where  $\tau$  is arbitrary in  $[0, T]$ . Hence

$$\begin{aligned} \|u_t^\gamma\|_{L^2(Q_\tau)}^2 + \int_\Omega \Psi(u_x^\gamma)|_{t=\tau} dx + \int_\Omega \left[ \frac{(u^\gamma)^4}{4} \Big|_{t=\tau} + \frac{(u_0^\gamma)^2}{2} \right] dx \\ \leq \int_\Omega \Psi(u_{0x}^\gamma) dx + \left( \frac{m_0^4}{4} + \frac{m_0^2}{2} \right) |\Omega|, \end{aligned} \quad (3.5)$$

from the bounds we have on  $u_0^\gamma$  and  $u^\gamma$ . Hence using the bound on  $\int_\Omega \Psi(u_x) dx$  and the bound on the total variation of the regularised initial data, it follows from (3.5) taking  $\tau = T$ , that

$$\begin{aligned} \|u_t^\gamma\|_{L^2(Q_T)}^2 &\leq \int_\Omega \Psi(u_{0x}^\gamma) dx + \left( \frac{m_0^4}{4} + \frac{m_0^2}{2} \right) |\Omega| \\ &\leq \int_\Omega |u_{0x}^\gamma| dx + \left( \frac{m_0^4}{4} + \frac{m_0^2}{2} + 1 \right) |\Omega| \\ &\leq C(\Omega) \int_\Omega |u_{0x}| dx + C_1 < \infty, \end{aligned} \quad (3.6)$$

since  $u_0 \in BV(\Omega)$ . Thus we have that the sequence  $\{u_t^\gamma\}$  is uniformly bounded in  $L^2(Q_T)$  and therefore also in  $L^1(Q_T)$ .

[B]: We will also need to show that the sequence  $\{u^\gamma\}$  is uniformly bounded in the space  $L^\infty((0, T), BV(\Omega))$  and also that  $\{u^\gamma\}$  is uniformly bounded in  $BV(Q_T)$ . To see the former, first note that (3.5) also implies that

$$\int_\Omega \Psi(u_x^\gamma)|_{t=\tau} dx \leq C(\Omega) \int_\Omega |u_{0x}| dx + \left( \frac{m_0^4}{4} + \frac{m_0^2}{2} \right) |\Omega|,$$

but since  $\tau$  was arbitrary in  $[0, T]$  we have, using the lower bound on  $\int_\Omega \Psi(u_x)$  once again, that for all  $t \in [0, T]$

$$\begin{aligned} C(\Omega) \int_\Omega |u_{0x}| dx + \left( \frac{m_0^4}{4} + \frac{m_0^2}{2} \right) |\Omega| &\geq \int_\Omega \Psi(u_x^\gamma) dx \geq \int_\Omega |u_x^\gamma| dx - |\Omega|, \\ \text{so that } \int_\Omega |u_x^\gamma| dx &\leq C(\Omega) \int_\Omega |u_{0x}| dx + C_1 \leq C_2 \quad \forall t \in [0, T]. \end{aligned} \quad (3.7)$$

This, together with the fact that  $u^\gamma(t, \cdot) \in L^1(\Omega)$  for all  $t \in [0, T]$  implies that

$$\|u^\gamma(t, \cdot)\|_{BV(\Omega)} < C_3 \quad \forall t \in [0, T], \quad \text{with } C_3 \text{ independent of } \gamma \text{ and of } t,$$

so that  $\sup_{0 < t \leq T} \|u^\gamma(t, \cdot)\|_{BV(\Omega)} < C_3$ . Hence we have that the sequence  $\{u^\gamma\}$  is indeed uniformly bounded in  $L^\infty((0, T), BV(\Omega))$ .

Since  $u^\gamma(t, \cdot) \in L^1(\Omega) \forall t \in [0, T]$ , we infer that  $u^\gamma \in L^1(Q_T)$  and (3.7) implies that

$$\int_0^T \int_\Omega |u_x^\gamma| dx dt \leq C_2 T,$$

we have that

$$\|u^\gamma\|_{BV(Q_T)} < C_4, \quad (3.8)$$

for  $C_4$  independent of  $\gamma$  and so  $u^\gamma$  is also uniformly bounded in  $BV(Q_T)$ .

[C]: We now establish that the sequence  $\{u^\gamma(t, \cdot)\}$  converges in the space  $L^2(\Omega)$  as  $\gamma \rightarrow 0$  for all  $t \in [0, T]$  and that the sequence  $\{u^\gamma\}$  converges in the space  $L^\infty((0, T), L^2(\Omega))$  as  $\gamma \rightarrow 0$ . To this end, consider  $u^{\gamma_m}$  and  $u^{\gamma_n}$  both satisfying the regularised problem, multiply the difference of the two equations by the difference  $u^{\gamma_m} - u^{\gamma_n}$ , then integrate over  $Q_\tau$  to obtain

$$\begin{aligned} \frac{1}{2} \int_{Q_\tau} \frac{\partial}{\partial t} (u^{\gamma_m} - u^{\gamma_n})^2 dx dt &= - \int_{Q_\tau} (\psi(u_x^{\gamma_m}) - \psi(u_x^{\gamma_n})) (u_x^{\gamma_m} - u_x^{\gamma_n}) dx dt \\ &\quad + \int_{Q_\tau} (f(u^{\gamma_m}) - f(u^{\gamma_n})) (u^{\gamma_m} - u^{\gamma_n}) dx dt \end{aligned} \quad (3.9)$$

But since the function  $\psi(s)$  is monotonic, the first term on the right-hand side of (3.9) is non-positive and so (3.9) becomes

$$\begin{aligned} \int_0^\tau \frac{d}{dt} \left( \int_\Omega (u^{\gamma_m} - u^{\gamma_n})^2 dx \right) dt &\leq 2 \int_{Q_\tau} (f(u^{\gamma_m}) - f(u^{\gamma_n})) (u^{\gamma_m} - u^{\gamma_n}) dx dt \\ &= 2 \int_{Q_\tau} [(u^{\gamma_m} - u^{\gamma_n}) - \{(u^{\gamma_m})^3 - (u^{\gamma_n})^3\}] (u^{\gamma_m} - u^{\gamma_n}) dx dt \\ &= 2 \int_{Q_\tau} [1 - \{(u^{\gamma_m})^2 + u^{\gamma_m} u^{\gamma_n} + (u^{\gamma_n})^2\}] (u^{\gamma_m} - u^{\gamma_n})^2 dx dt \\ &\leq 2 \int_{Q_\tau} |[\{(u^{\gamma_m})^2 + u^{\gamma_m} u^{\gamma_n} + (u^{\gamma_n})^2\} - 1]| |u^{\gamma_m} - u^{\gamma_n}|^2 dx dt \\ &\leq 2|3m_0^2 - 1| \int_{Q_\tau} |u^{\gamma_m} - u^{\gamma_n}|^2 dx dt \\ &= \int_0^\tau |6m_0^2 - 2| \left( \int_\Omega (u^{\gamma_m} - u^{\gamma_n})^2 dx \right) dt \end{aligned} \quad (3.10)$$

Thus if we define  $C(m_0) = |6m_0^2 - 2|$  then we have, since  $\tau$  is arbitrary in  $[0, T]$

$$\frac{d}{dt} \int_\Omega (u^{\gamma_m} - u^{\gamma_n})^2 dx \leq C(m_0) \int_\Omega (u^{\gamma_m} - u^{\gamma_n})^2 dx.$$

Hence Gronwall's inequality implies that

$$\int_\Omega (u^{\gamma_m} - u^{\gamma_n})^2 dx \leq e^{C(m_0)\tau} \int_\Omega (u_0^{\gamma_m} - u_0^{\gamma_n})^2 dx,$$

so that from (3.10)

$$\begin{aligned} \int_{\Omega} (u^{\gamma_m} - u^{\gamma_n})^2|_{t=\tau} dx &\leq \int_0^{\tau} C(m_0) \left( \int_{\Omega} (u^{\gamma_m} - u^{\gamma_n})^2 dx \right) dt + \int_{\Omega} (u_0^{\gamma_m} - u_0^{\gamma_n})^2 dx \\ &\leq (C(m_0)\tau e^{C(m_0)\tau} + 1) \int_{\Omega} (u_0^{\gamma_m} - u_0^{\gamma_n})^2 dx, \end{aligned}$$

but since  $\tau$  was arbitrary in  $[0, T]$  and  $u^{\gamma_m}$  and  $u^{\gamma_n}$  both satisfy the regularised problem, we have

$$\|u^{\gamma_m}(t, \cdot) - u^{\gamma_n}(t, \cdot)\|_{L^2(\Omega)} \rightarrow 0 \text{ as } \gamma_m, \gamma_n \rightarrow 0, \text{ for all } t \in [0, T].$$

So  $u^{\gamma_n}(t, \cdot)$  is Cauchy in  $L^2(\Omega)$  for all  $t \in [0, T]$  hence the sequence  $u^{\gamma_n}(t, \cdot)$  converges in  $L^2(\Omega)$  for all  $t \in [0, T]$  and from this it follows that  $u^{\gamma}$  converges in  $L^\infty((0, T), L^2(\Omega))$ .  $\square$

We now pass to the limit as  $\gamma \rightarrow 0$  making use of the above properties of the sequence  $u^{\gamma}$ . We have shown that there exists a unique  $u \in L^\infty((0, T), L^2(\Omega))$  such that

$$\|u^{\gamma}(t, \cdot) - u(t, \cdot)\|_{L^2(\Omega)} \rightarrow 0 \text{ as } \gamma \rightarrow 0 \forall t \in [0, T] \text{ and } \|u^{\gamma} - u\|_{L^\infty(Q_T)} \rightarrow 0 \text{ as } \gamma \rightarrow 0,$$

but then this implies convergence in  $L^1$  also so that we have

$$\|u^{\gamma}(t, \cdot) - u(t, \cdot)\|_{L^1(\Omega)} \rightarrow 0 \text{ as } \gamma \rightarrow 0 \forall t \in [0, T],$$

and

$$\|u^{\gamma} - u\|_{L^1(Q_T)} \rightarrow 0 \text{ as } \gamma \rightarrow 0 \tag{3.11}$$

using the Cauchy Schwarz inequality.

We have also shown uniform boundedness of  $u_t^{\gamma}$  in  $L^2(Q_T)$ , hence  $\|u_t^{\gamma}\|_{L^2(Q_T)} \leq C$  and so by weak compactness in  $L^2(Q_T)$ , we can extract a subsequence that we still denote as  $\{u_t^{\gamma}\}$  which is such that

$$u_t^{\gamma} \rightharpoonup u_t \text{ in } L^2(Q_T) \text{ with } u_t \in L^2(Q_t).$$

This implies that given  $\varphi \in L^2(\Omega)$  we have

$$\int_0^t \langle u_t^{\gamma}(x, s), \varphi \rangle_{L^2(\Omega)} ds = \langle u^{\gamma}(x, t), \varphi \rangle_{L^2(\Omega)} - \langle u_0^{\gamma}(x), \varphi \rangle_{L^2(\Omega)},$$

and letting  $\gamma \rightarrow 0$  gives

$$\int_0^t \langle u_t(x, s), \varphi \rangle_{L^2(\Omega)} ds = \langle u(x, t), \varphi \rangle_{L^2(\Omega)} - \langle u_0(x), \varphi \rangle_{L^2(\Omega)},$$

from which it follows that the limit function  $u(x, t)$  satisfies the initial condition,  $u(x, 0) = u_0(x)$ , and following the same reasoning as for (3.4), the limit function  $u$  is also uniformly bounded in  $L^\infty(Q_T)$ .

We now prove that the limit function  $u$  is in  $BV(Q_T)$ . We have shown that the sequence  $\{u^{\gamma}\}$  is uniformly bounded in  $BV(Q_T)$ . Hence we can extract a subsequence

denoted  $\{u^{\gamma_i}\}$  that converges weakly to a  $BV$  function  $\eta$ , say. That is,  $u^{\gamma_i}(x, t) \rightharpoonup \eta(x, t)$  in  $BV(Q_T)$ -weak-\* with  $\eta \in BV(Q_T)$ , but this means that  $u^{\gamma_i} \rightarrow \eta$  in  $L^1(Q_T)$ , so from (3.11) by the uniqueness of the limit we must have

$$\eta = u \in BV(Q_T) \quad (3.12)$$

Hence by definition of  $BV$  functions on  $Q_T$ , we conclude from (3.12) that weak first derivative in space of  $u$  is a bounded measure on  $Q_T$ .

We can now show that the limit function  $u$  is such that  $u(t, \cdot) \in BV(\Omega)$  for every  $t \in [0, T]$ . That the sequence  $\{u^\gamma\}$  is uniformly bounded in  $L^\infty((0, T), BV(\Omega))$  means that

$$\|u^{\gamma_i}(t, \cdot)\|_{BV(\Omega)} < C_5, \quad \text{for almost every } t \in [0, T].$$

Fix  $t_0$  arbitrary in  $[0, T]$ . We can extract a subsequence  $\{u^{\gamma_j}\}$  of  $\{u^{\gamma_i}\}$  such that  $u^{\gamma_j}(t_0, \cdot) \rightharpoonup U(t_0, \cdot)$  weak-\* in  $BV(\Omega)$  with  $U(t_0, \cdot) \in BV(\Omega)$ . But this means that  $u^{\gamma_j}(t_0, \cdot) \rightarrow U(t_0, \cdot)$  in  $L^1(\Omega)$  and so we have once again from (3.11) that  $u(t, \cdot) = U(t, \cdot) \in BV(\Omega)$  for all  $t \in [0, T]$  since  $t_0$  was arbitrary in  $[0, T]$ .

It has been shown in [5] that for  $u \in BV(\Omega)$  and  $\Psi$  convex, the functional  $\int_\Omega \Psi(u_x) dx$  is lower semi-continuous with respect to the  $L^1$  convergence. Hence, since we know that  $u(t, \cdot) \in BV(\Omega)$  for almost all  $t \in [0, T]$  and that  $\|u^\gamma(t, \cdot) - u(t, \cdot)\|_{L^1(\Omega)} \rightarrow 0$  as  $\gamma \rightarrow 0$  for all  $t \in [0, T]$  we must have that

$$\int_\Omega \Psi(u_x) dx \leq \liminf_{\gamma \rightarrow 0} \int_\Omega \Psi(u_x^\gamma) dx \quad \text{for all } t \in [0, T]. \quad (3.13)$$

We noted earlier that from (3.5), it follows that

$$\int_\Omega \Psi(u_x^\gamma) dx \leq C(\Omega) \int_\Omega |u_{0x}| + \left(\frac{m_0^4}{4} + \frac{m_0^2}{2}\right) |\Omega| \quad \forall t \in [0, T]. \quad (3.14)$$

Hence taking the limit inferior of (3.14) as  $\gamma \rightarrow 0$ , we see that

$$\begin{aligned} \int_\Omega (|u_x| - 1) dx &\leq \int_\Omega \Psi(u_x) dx \\ &\leq \liminf_{\gamma \rightarrow 0} \int_\Omega \Psi(u_x^\gamma) dx \leq C(\Omega) \int_\Omega |u_{0x}| dx + \left(\frac{m_0^4}{4} + \frac{m_0^2}{2}\right) |\Omega| \quad \forall t \in [0, T] \end{aligned}$$

Thus we are lead to conclude that  $\|u(t, \cdot)\|_{BV(\Omega)} < \infty$  for almost all  $t \in [0, T]$  and consequently

$$u \in L^\infty((0, T), BV(\Omega)).$$

For later, note that one may integrate (3.13) on  $[0, T]$  to obtain

$$\liminf_{\gamma \rightarrow 0} \int_{Q_T} \Psi(u_x^\gamma) dx dt \geq \int_{Q_T} \Psi(u_x) dx dt.$$

An additional result that we will need when passing to the limit as  $\gamma \rightarrow 0$  is that as  $\|u^\gamma - u\|_{L^1(Q_T)} \rightarrow 0$  as  $\gamma \rightarrow 0$ ,  $\|f(u) - f(u^\gamma)\|_{L^1(Q_T)} \rightarrow 0$ . This follows easily when



one considers

$$\begin{aligned} & \int_0^T \int_{\Omega} |f(u) - f(u^\gamma)| dx dt = \int_0^T \int_{\Omega} |u - u^3 - (u^\gamma - (u^\gamma)^3)| dx dt \\ & \leq \int_0^T \int_{\Omega} |u - u^\gamma| dx dt + \int_0^T \int_{\Omega} |u - u^\gamma| |u^2 + uu^\gamma + (u^\gamma)^2| dx dt \\ & \leq \int_0^T \int_{\Omega} |u - u^\gamma| dx dt + 3m_0^2 \int_0^T \int_{\Omega} |u - u^\gamma| dx dt \rightarrow 0 \text{ as } \gamma \rightarrow 0. \end{aligned}$$

So far we have shown that the limit function  $u$  is such that

$$u \in L^\infty(Q_T) \cap L^\infty((0, T), BV(\Omega)) \cap \{u : u_x \in M(Q_T)\},$$

so that all that remains is to be proven is that the limit function  $u$  satisfies the variational inequality (3.3). Note that the variational inequality holds for the solutions  $u^\gamma$  of the regularised problems with test functions taken from the smooth sequence  $\{v^n\}_{n \in \mathbb{N}} \subset C^\infty(Q_T)$  i.e.

$$\int_{Q_T} (u_t^\gamma - f(u^\gamma))(v^n - u^\gamma) dx dt + \int_{Q_T} \Psi(v_x^n) - \Psi(u_x^\gamma) dx dt \geq 0. \quad (3.15)$$

It is shown in [6] that the space  $C^\infty(Q_T)$  is dense in  $BV(Q_T)$  equipped with the topology defined by the distance

$$d(u, w) = \|u - w\|_{L^1(Q_T)} + \left| \int_{Q_T} |u_x| - \int_{Q_T} |w_x| \right| + \left| \int_{Q_T} \Psi(u_x) - \int_{Q_T} \Psi(w_x) \right|,$$

which means that one can approximate  $BV(Q_T)$  functions by a sequence of  $C^\infty(Q_T)$  functions, i.e. for  $v \in BV(Q_T)$ , there exists a sequence  $\{v^n\} \in C^\infty(Q_T)$  such that

$$\begin{aligned} & \int_{Q_T} \Psi(v_x^n) \rightarrow \int_{Q_T} \Psi(v_x) \text{ as } n \rightarrow \infty \\ \text{and } & \int |v^n - v| dx dt \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned} \quad (3.16)$$

This combined with all the properties that have been established for solutions  $u^\gamma$  to the regularised problem, means that one may pass to the limit as  $n \rightarrow \infty$  and subsequently as  $\gamma \rightarrow 0$  in (3.15) to obtain the result.

As usual, in order to prove uniqueness of a weak solution to our problem we suppose non-uniqueness and derive a contradiction. Hence suppose there are two weak solutions  $u_1$  and  $u_2$  satisfying problem (3.1) and therefore the variational inequality (3.3) with

$$u_1(x, 0) = u_2(x, 0) = u_0(x). \quad (3.17)$$

Take the variational inequality first with  $u = u_1, v = u_2$  and then with  $u = u_2, v = u_1$  so that

$$\int_{Q_\tau} \left( \frac{\partial u_1}{\partial t} - f(u_1) \right) (u_2 - u_1) dx dt + \int_{Q_\tau} \Psi((u_2)_x) - \Psi((u_1)_x) dx dt \geq 0,$$

and

$$\int_{Q_\tau} \left( \frac{\partial u_2}{\partial t} - f(u_2) \right) (u_1 - u_2) dx dt + \int_{Q_\tau} \Psi((u_1)_x) - \Psi((u_2)_x) dx dt \geq 0.$$

Adding these two inequalities gives

$$\int_{Q_\tau} \frac{\partial(u_1 - u_2)}{\partial t} (u_1 - u_2) dx dt \leq \int_{Q_\tau} (f(u_1) - f(u_2))(u_1 - u_2) dx dt.$$

As before

$$\begin{aligned} \int_{\Omega} (u_1 - u_2)^2 dx|_{t=\tau} &\leq \int_0^\tau C(m_0) \left( \int_{\Omega} (u_1 - u_2)^2 dx \right) dt + \int_{\Omega} (u_1(x, 0) - u_2(x, 0))^2 dx \\ &\leq (C(m_0)\tau e^{C(m_0)T} + 1) \int_{\Omega} (u_1(x, 0) - u_2(x, 0))^2 dx, \end{aligned}$$

using again the Gronwall inequality. Thus it follows from (3.17) that

$$\|u_1(\tau, \cdot) - u_2(\tau, \cdot)\|_{L^2(\Omega)} = 0,$$

and uniqueness follows from  $\tau$  being arbitrary in  $[0, T]$ .

#### 4. STATIONARY SOLUTIONS IN ONE DIMENSION

The one-dimensional Neumann stationary problem for (1.2)

$$\begin{aligned} \left( \frac{u'}{\sqrt{1 + (u')^2}} \right)' + \lambda f(u) &= 0, \\ u'(0) = u'(L) &= 0, \end{aligned} \tag{4.1}$$

$\lambda = 1/\epsilon$ , is studied in [2] and it is shown through an analysis of the time map associated with the equation that the bifurcation behaviour depends not only on  $\lambda$  but also on the length  $L$  of the interval. For fixed  $L$ , the following proposition is proven using Liapunov-Schmidt reduction

**Proposition 4.1.** *The  $k$ -th bifurcation from the trivial solution of (4.1) is a supercritical pitchfork if  $L > k\pi/\sqrt{2}$  and a subcritical pitchfork if the inequality is reversed.*

It is also shown that for any given value of  $L$ , there is a value  $\lambda^*(L)$  beyond which there cannot exist classical, i.e.  $C^2((0, L) \cap C^1([0, L])$ , solutions to (4.1) and solutions at  $\lambda = \lambda^*(L)$  develop infinite gradient.

Solutions to (4.1) are defined in the  $BV$  sense as functions of bounded variation which satisfy the variational inequality

$$-\lambda \int_{\Omega} f(u)(v - u) dx + \int_{\Omega} \Psi(v_x) - \Psi(u_x) dx \geq 0 \quad \forall v \in BV(\Omega), \tag{4.2}$$

which is obtained from (3.3) if one assumes that  $u_t = 0$ . If without loss of generality one considers monotone decreasing solutions to (4.1), a theorem proven in [2] is that

(discontinuous) solutions constructed by patching together different level curves of the Hamiltonian

$$H(u, u') = 1 - \frac{1}{\sqrt{1 + (u')^2}} - \lambda W(u),$$

which satisfy  $u_x = 0$  at  $x = 0$  and  $x = L$  are solutions to (4.1) in the  $BV$  sense. Hence there exists a continuum of discontinuous stationary solutions beyond  $\lambda^*(L)$ . One can easily generate initial conditions which, for  $L$  fixed and  $\lambda > \lambda^*(L)$ , converge to a discontinuous stationary solution of (1.2) by taking

$$u_0(x) = -\alpha \tanh\left(\beta\left(\frac{x}{L} - \gamma\right)\right), \quad (4.3)$$

which serves as an approximation to the discontinuous steady state with a discontinuity at some  $x_0 = \gamma L$  for  $\gamma \in (0, 1)$  and where  $u_0(0) = -u_0(L) = \alpha \in (0, 1)$  and  $\beta$  is large and such that  $u'_0(x_0) = -\frac{\alpha\beta}{L}$ . In Figure 1, we fix  $L = 2.5$  (supercritical) and  $\lambda = 5 > \lambda^*(L) \approx 4.019534$  and solve (1.2), (4.3) with  $\alpha = 0.98$ ,  $\beta = 500$  and  $\gamma = 0.24, 0.5$  and  $0.76$  respectively and the solutions indeed converge to a discontinuous steady state.




FIGURE 1. Convergence of three initial data to three of the infinitely many steady states of (1.2) for  $L = 2.5$  and  $\lambda = 5$ .

For details of the numerics, please consult [3]. Note from this figure that these solutions have some stability properties (see [2] for a discussion of the right notion of stability for this case.)

There are also similarly stable non-monotone solutions as in Figure 2 arising from non-monotone initial data

$$u_0(x) = \frac{4x(L-x)}{L^2} \sin\left(\frac{10\pi x^2}{L^2}\right),$$

if  $\epsilon$  is taken to be sufficiently small in (1.2).




FIGURE 2. Convergence of a non-monotone initial condition to a non-monotone steady solution of (1.2) with  $\epsilon = 0.001$  and  $L = 2.5$

This indicates that the structure of patterns that the bistable quasilinear equation gives rise to is much richer than in the semilinear case, in which only the constant solutions  $\pm 1$  attract all initial conditions with probability one.

Finally, given a continuum of stationary solutions, it is interesting to know which has the lowest energy. It turns out that it is the most asymmetric of the possible stationary solutions that minimize the energy over the continuum.




FIGURE 3. Plot of position of interface  $x_0$  against energy  $E[u]$  of stationary solutions to (1.2) corresponding to  $L = 2.5$  and  $\lambda = 5$ .

## 5. CONCLUDING REMARKS

We have presented a model for solid-solid phase transitions and have proved the existence of weak (variational inequality) solutions on all  $[0, T]$ ,  $T > 0$ . We have also presented some results on discontinuous stationary solutions for the model, which have some stability properties in stark contrast to the semilinear situation. Much work remains to be done, in particular proving stabilisation of orbits. We expect that nonlinear semigroup techniques of [1] together with a Simon-Lojasiewicz inequality type result [4] will be required for that.

## REFERENCES

- [1] F. Andreu-Vaillio, V. Caselles & J. M. Mazón, Parabolic quasilinear equations minimizing linear growth functionals, vol. 223 of Progress in Mathematics, Birkhäuser Verlag, Basel, 2004
- [2] M. Burns, & M. Grinfeld, Steady state solutions of a bistable quasilinear equation with saturating flux. (*to appear*)
- [3] Burns, M. Reaction-Diffusion Equations with Saturating Flux, Ph.D. Thesis. University of Strathclyde, in preparation.
- [4] R. Chill, On the Lojasiewicz-Simon gradient inequality, *J. Funct. Anal.* **201**, (2003), 572–601.
- [5] L. Dascal, S. Kamin & N. Sochen, A variational inequality for discontinuous solutions of degenerate parabolic equations., *RACSAM* **99**, (2005), 243–256.
- [6] F. Demengel & R. Temam, Convex functions of a measure and applications. *Indiana Univ. Math. J.* **33**, (1984), 673–709
- [7] P. Rosenau, Extension of Landau-Ginzburg free-energy functionals to high-gradient domains. *Phys. Rev. A* **39**, (1989), 614–6617.
- [8] P. Rosenau, Free-energy functionals at the high-gradient limit. *Phys. Rev. A* **41**, (1990), 2227–2230.
- [9] J. Smoller & A. Wasserman, Global bifurcation of steady state solutions. *J. Differ. Equ* **39**, 1981, 269–290