

Total Syntheses of Conformationally Locked Difluorinated Pentopyranose Analogues and a Pentopyranosyl Phosphate Mimetic

Jonathan A. L. Miles,[†] Lisa Mitchell,^{†,‡} Jonathan M. Percy,^{*,‡} Kuldeep Singh,[†] and E. Uneyama[†]

Department of Chemistry, University of Leicester, University Road, Leicester LE1 7RH, U.K.,
and Department of Pure and Applied Chemistry, WestCHEM, University of Strathclyde,
Thomas Graham Building, 295 Cathedral Street, Glasgow G1 1XL, U.K.

jonathan.percy@strath.ac.uk

Received September 30, 2006



Trifluoroethanol has been elaborated, via a telescoped sequence involving a metalated difluoroenol, a difluoroallylic alcohol, [2,3]-Wittig rearrangement, and ultimately an RCM reaction and requiring minimal intermediate purification, to a number of cyclooctenone intermediates. Epoxidation of these intermediates followed by transannular ring opening or dihydroxylation, then transannular hemiacetalization delivers novel bicyclic analogues of pentopyranoses, which were elaborated (in one case) to an analogue of a glycosyl phosphate.

Introduction

Saccharide recognition is a key event in a wide range of biological processes. Sugars present arrays of hydroxyl groups in a spatially defined manner, and proteins bind to and distinguish between different saccharides by forming complex

networks of hydrogen bonds to them.¹ The core six-membered oxacycle in pyranose sugars also exerts a significant effect on a range of conformational properties via the anomeric effect.² The linkages between sugars in di- and higher saccharides are acetals, and a wide range of glycosyltransferases and glycosidases exist to synthesize and cleave those linkages, respectively. Saccharide mimetics,³ which present hydroxyl groups in a useful manner but cannot be cleaved from their sites of attachment by glycosidases, could be useful probes of sugar-processing enzymes. If the mimetics lack the pyranose oxygen and therefore the anomeric effect, different conformers could become available.

Many groups have described the synthesis of five-, six-, and seven-membered carbocyclic analogues of saccharides. More recently, highly functionalized cyclooctane derivatives have attracted attention as ring-expanded analogues (Chart 1). The Sinay group combined the ideas of the stability of carbasugars with the potential for occupying uncharted conformational space, synthesizing **1**, which was shown by NOE to occupy a boat-chair conformation related to that of the corresponding galac-


[†] University of Leicester.

[‡] University of Strathclyde.

(1) (a) Loganathan, D.; Aich, U. *Glycobiology* **2006**, *16*, 343. (b) Laederach, A.; Reilly, P. J. *Proteins: Struct., Funct., Bioinf.* **2005**, *60*, 591. (c) Carcabal, P.; Jockusch, R. A.; Hunig, I.; Snoek, L. C.; Kroemer, R. T.; Davis, B. G.; Gamblin, D. P.; Compagnon, I.; Oomens, J.; Simons, J. P. *J. Am. Chem. Soc.* **2005**, *127*, 11414. (d) Flint, J.; Bolam, D. N.; Nurizzo, D.; Taylor, E. J.; Williamson, M. P.; Walters, C.; Davies, G. J.; Gilbert, H. J. *J. Biol. Chem.* **2005**, *280*, 23718. (e) Fernandez, M. D.; Canada, F. J.; Jimenez-Barbero, J.; Cuevas, G. *J. Am. Chem. Soc.* **2005**, *127*, 7379. (f) Mitchell, E. P.; Sabin, C.; Snajdrova, L.; Pokorna, M.; Perret, S.; Gautier, C.; Hofr, C.; Gilboa-Garber, N.; Koca, J.; Wimmerova, M.; Imberty, A. *Proteins: Struct., Funct., Bioinf.* **2005**, *58*, 735. (g) Loris, R.; Imberty, A.; Beeckmans, S.; Van, Driessche, E.; Read, J. S.; Bouckaert, J.; De Greve, H.; Buts, L.; Wyns, L. *J. Biol. Chem.* **2003**, *278*, 16297. (h) Vyas, N. K.; Vyas, M. N.; Chervenak, M. C.; Bundle, D. R.; Pinto, B. M.; Quiocho, F. A. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 15023. (i) Pathiaseril, A.; Woods, R. J. *J. Am. Chem. Soc.* **2000**, *122*, 331. (j) Zacharias, M.; Straatsma, T. P.; McCammon, J. A.; Quiocho, F. A. *Biochemistry* **1993**, *32*, 7428. (k) Quiocho, F. A. *Biochem. Soc. Trans.* **1993**, *21*, 442. (l) Jimenez-Barbero, J.; Canada, F. J.; Cuevas, G.; Asensio, J. L.; Aboitiz, N.; Canales, A.; Chavez, M. I.; Fernandez-Alonso, M. C.; Garcia-Herrero, A.; Mari, S.; Vidal, P. *NMR Spectroscopy and Computer Modeling of Carbohydrates: Recent Advances*; ACS Symposium Series 930; American Chemical Society: Washington, DC, 2006; p 60. (m) Bryce, R. A.; Hillier, I. H.; Naismith, J. H. *Biophys. J.* **2001**, *81*, 1373.

(2) (a) Kirby, A. *The Anomeric Effect and Related Stereoelectronic Effects at Oxygen*; Springer-Verlag: Berlin, 1983. (b) For recent mechanistic studies, see: Bennet, A. J.; Kitos, T. E. *J. Chem. Soc., Perkin Trans.* **2002**, *2*, 1207. (c) For a computational examination of the mechanism, see: Dimelow, R. J.; Bryce, R. A.; Masters, A. J.; Hillier, I. H.; Burton, N. A. *J. Chem. Phys.* **2006**, *124*. (d) Mohr, M.; Bryce, R. A.; Hillier, I. H. *J. Phys. Chem. A* **2001**, *105*, 8216.


CHART 1



topyranose.⁴ The synthesis used carbohydrate starting materials to provide most of the functionality in the products and ensure stereocontrol, exploiting a Lewis acid mediated ring-expanding Claisen rearrangement (described originally by Paquette) as the key step. More recently, Paquette has shown that zirconocene-promoted ring contraction can be used to convert vinylfuranosides to cyclooctane polyols via vinyl cyclobutanones, and a [3,3]-sigmatropic rearrangement again secures the carbocycles.⁵ Using a quite different approach, Mehta and co-workers have shown that unsaturated eight-membered ring compounds such as cyclooctatetraene can be elaborated controllably to the corresponding polyols.⁶

The ring-expanding Claisen rearrangement also allows the synthesis of masked cyclooctanones **2**. Van Boom and the Leiden group exploited transannular strain-relieving nucleophilic attack upon a ketone carbonyl group to close a number of bicyclic systems, which provide conformationally locked analogues of sugars **3** and azasugars **4**.⁷ The bicyclic array is very interesting because of the extremely low reactivity of the pseudoglycosidic bond. Kirby and co-workers showed that **5** was 10^{13} times less reactive than **6** because of the way the bicyclic architecture opposes stabilization of developing positive charge at the pseudooanomeric carbon through oxacarbenium ion formation as the pseudoglycosidic bond stretches toward cleavage.⁸ The idea of using conformational locking to allow mimicry

SCHEME 1. Outline of Sequence Showing the Development of NDP Sugar Analogues



of conformations traversed at transition states has also been explored extensively.⁹

We¹⁰ and others¹¹ have developed a number of approaches for the synthesis of fluorinated analogues of the molecules of nature from commercial fluorinated starting materials, in which RCM forms a key step. One of our projects aimed to develop a de novo route via difluorinated cyclooctenones **7** to conformationally locked difluorinated analogues **8** of pentoses and their phosphates **9**, which we would advance ultimately to analogues of NDP sugars, with the global aim of the development of new chemical tools for the study of glycosyltransferase enzymes. The syntheses would start from sustainable fluorinated starting materials, avoiding the use of materials banned under the Montreal and Kyoto Protocols (Scheme 1). Although these protocols deal with large-scale production, they affect the availability and supply of materials for use on the laboratory scale; the development of a new methodology based on banned substances would therefore seem like nugatory effort.

The fluorine atoms would allow location of the compounds in vitro or in vivo by ¹⁹F NMR, and they could report conformational changes in the hydroxyl-bearing ring through ³J_{H-F} coupling constants.¹² In our preliminary publications, we showed how we could use metalated difluoroalkene chemistry to advance trifluoroethanol rapidly to precursors to eight-membered rings and then close them via RCM¹³ to afford difluorinated cyclooctenones, templates for stereoselective oxidation (dihydroxylation¹⁴ or epoxidation¹⁵), and transannular reactions. These preliminary findings delivered a number of model polyol systems which contain the distinctly unnatural *gem*-dimethyl and *N,N*-diethylcarbamoyloxy groups, which we now wished to delete. We wished to explore a route, which would avoid the use of strong base/low-temperature conditions

(3) (a) Haneda, T.; Goekjian, P. G.; Kim, S. H.; Kishi, Y. *J. Org. Chem.* **1992**, *57*, 490. (b) Goekjian, P. G.; Wu, T. C.; Kang, H. Y.; Kishi, Y. *J. Org. Chem.* **1991**, *56*, 6422. (c) Kuntz, D. A.; Ghavami, A.; Johnston, B. D.; Pinto, B. M.; Rose, D. R. *Tetrahedron: Asymmetry* **2005**, *16*, 25. (d) Yang, G. L.; Schmiege, J.; Tsuji, M.; Franck, R. W. *Angew. Chem., Int. Ed.* **2004**, *43*, 3818. (e) Mikkelsen, L. M.; Hernaiz, M. J.; Martin-Pastor, M.; Skrydstrup, T.; Jimenez-Barbero, J. *J. Am. Chem. Soc.* **2002**, *124*, 14940. For reviews, see: (f) Sears, P.; Wong, C. H. *Angew. Chem., Int. Ed.* **1999**, *38*, 2301. (g) Wong, C. H. *Acc. Chem. Res.* **1999**, *32*, 376.

(4) (a) Wang, W.; Zhang, Y. M.; Sollogoub, M.; Sinay, P. *Angew. Chem., Int. Ed.* **2000**, *39*, 2466. (b) Wang, W.; Zhang, Y. M.; Zhou, H. H.; Bleriot, Y.; Sinay, P. *Eur. J. Org. Chem.* **2001**, 1053.

(5) Paquette, L. A.; Zhang, Y. L. *J. Org. Chem.* **2006**, *71*, 4353.

(6) (a) Mehta, G.; Pallavi, K. *Tetrahedron Lett.* **2004**, *45*, 3865. (b) Mehta, G.; Pallavi, K. *Chem. Commun.* **2002**, 2828. (c) For a review of earlier work, see: Mehta, G.; Singh, V. *Chem. Rev.* **1999**, *99*, 881.

(7) van Hooft, P. A. V.; Litjens, R.; van der Marel, G. A.; van Boeckel, C. A. A.; van Boom, J. H. *Org. Lett.* **2001**, *3*, 731.

(8) Briggs, A. J.; Evans, C. M.; Glenn, R.; Kirby, A. J. *J. Chem. Soc., Perkin Trans. 2* **1983**, 1637.

(9) For recent examples, see: (a) Bleriot, Y.; Vadivel, S. K.; Herrera, A. J.; Greig, I. R.; Kirby, A. J.; Sinay, P. *Tetrahedron* **2004**, *60*, 6813. (b) Lorthiois, E.; Meyyappan, M.; Vasella, A. *Chem. Commun.* **2000**, 1829.

(10) (a) Audouard, C.; Fawcett, J.; Griffiths, G. A.; Percy, J. M.; Pintat, S.; Smith, C. A. *Org. Biomol. Chem.* **2004**, *2*, 528. (b) Audouard, C.; Fawcett, J.; Griffiths, G. A.; Kerouredan, E.; Miah, A.; Percy, J. M.; Yang, H. L. *Org. Lett.* **2004**, *6*, 4269. (c) Kariuki, B. M.; Owton, W. M.; Percy, J. M.; Pintat, S.; Smith, C. A.; Spencer, N. S.; Thomas, A. C.; Watson, M. *Chem. Commun.* **2002**, 228. (d) Butt, A. H.; Percy, J. M.; Spencer, N. S. *Chem. Commun.* **2000**, 1691.



(11) (a) Fustero, S.; Catalan, S.; Piera, J.; Sanz-Cervera, J. F.; Fernandez, B.; Acena, J. L. *J. Org. Chem.* **2006**, *71*, 4010. (b) Fustero, S.; Sanchez-Rosello, M.; Jimenez, D.; Sanz-Cervera, J. F.; del Pozo, C.; Acena, J. L. *J. Org. Chem.* **2006**, *71*, 2706. (c) Fustero, S.; Bartolome, A.; Sanz-Cervera, J. F.; Sanchez-Rosello, M.; Soler, J. G.; de Arellano, C. R.; Fuentes, A. S. *Org. Lett.* **2003**, *5*, 2523. (d) Yang, Y. Y.; Meng, W. D.; Qing, F. L. *Org. Lett.* **2004**, *6*, 4257. (e) You, Z. W.; Wu, Y. Y.; Qing, F. L. *Tetrahedron Lett.* **2004**, *45*, 9479. (f) De Matteis, V.; van Delft, F. L.; de Gelder, R.; Tiebes, J.; Rutjes, F. *Tetrahedron Lett.* **2004**, *45*, 959. (g) Marhold, M.; Buer, A.; Hiemstra, H.; van Maarseveen, J. H.; Haufe, G. *Tetrahedron Lett.* **2004**, *45*, 57.

(12) Thibaudeau, C.; Plavec, J.; Chattopadhyaya, J. *J. Org. Chem.* **1998**, *63*, 4967.

(13) Griffiths, G. A.; Percy, J. M.; Pintat, S.; Smith, C. A.; Spencer, N.; Uneyama, E. *Org. Biomol. Chem.* **2005**, *3*, 2701.

(14) Fawcett, J.; Griffiths, G. A.; Percy, J. M.; Pintat, S.; Smith, C. A.; Spencer, N.; Uneyama, E. *Chem. Commun.* **2004**, 302.

(15) Fawcett, J.; Griffiths, G. A.; Percy, J. M.; Uneyama, E. *Org. Lett.* **2004**, *6*, 1277.

SCHEME 2. Retrosynthetic Analysis of Key Cyclooctenone 14**SCHEME 3. Attempted Halodifluoromethyl Ketone Syntheses**

if possible and minimize the number of purifications in the sequences, to complete syntheses of a number of analogues of sugar-like and an illustrative phosphate monoester through stereoselective oxidation reactions and the development of an effective phosphorylation strategy. We report the results of these studies in this manuscript.


Results and Discussion

Attempted Development of a Scaleable Route Based on Reductive Dehalogenation. Scheme 2 shows the retrosynthetic analysis carried out for **14** from key intermediate α,α -difluoro- β -hydroxy ketone **13**. The literature describes a number of methods for the generation of difluorinated silyl enol ether **12**, which could make **13** available through aldol chemistry under potentially scaleable conditions (Scheme 2).

Ishihara¹⁶ and Uneyama¹⁷ have described methods for the synthesis of difluoroenol silyl ethers such as **12** from chlorodifluoromethyl and trifluoromethyl ketones, respectively, so we prepared **11a** and **11b** from commercial 1-bromopent-4-ene via reaction of the Grignard reagent with electrophiles **10** derived from chlorodifluoroacetic or trifluoroacetic acids (Scheme 3).¹⁸ These starting materials are available at low cost and appear to be sustainable.

Typical reaction conditions for this type of perhalomethylketone synthesis use ester electrophiles or, alternatively, an excess of Grignard reagent and the free acid.¹⁸ The latter procedure would waste a moderately expensive bromide, so we preformed the sodium or magnesium salts using NaH or *i*-PrMgCl.

A large number of unsuccessful experiments are summarized in the Scheme. The ketones were isolated by distillation after careful extraction of the product into pentane/diethyl ether, and hydration of the ketones was a distinct problem during these procedures. The poor yield of the trifluoromethyl ketone was comparable to that obtained by Laurent and co-workers.¹⁹ We were unable to synthesize **12** in more than trace amounts from either **11a** or **11b** under published (Zn, Me₃SiCl, MeCN, Δ or excess Mg/Me₃SiCl, DMF) conditions. Uneyama¹⁷ has reported

SCHEME 4. Cyclooctenone Syntheses Based on Metalated Difluoroenol Acetal Chemistry

that aliphatic trifluoromethyl ketones undergo slow reductive defluorination under the latter conditions; for **11a**, the reaction was prohibitively slow. We also attempted to use the conditions described by Ishihara¹⁸ for direct aldol reaction between **11b** and acrolein or cinnamaldehyde. The best result (ca. 10% conversion) was obtained with the latter electrophile despite considerable efforts to optimize the reactions, which stopped at very low conversion, and we were not able to isolate any of **13**. We therefore decided to use our metalated difluoroenol chemistry to develop a working strategy to the target aldol.

Successful Dehydrofluorination/Metalation Route. Known²⁰ difluoroallylic alcohol **15** was synthesized in good yield (82%, 20 g scale) using our published procedure from the MEM-ether of trifluoroethanol and 4-pentenal (Scheme 4).²¹ The Kugelrohr-distilled alcohol was allylated under phase-transfer conditions²² to ether **16** (91%) and progressed without further purification (Scheme 4).

The rearrangement of **16** took place over 4 h on warming from -100 to -30 °C, and chromatography of the product returned a disappointing yield of pure **17** (ca. 30%) initially. We were able to purify hydroxyketone **13** (following enol ether cleavage²³) by Kugelrohr distillation improving the yield to 50% over the two steps and removing two chromatographic purifications from the sequence if the enol ether was cleaved directly from crude [2,3]-Wittig product. In the RCM reaction (5 mol % of **21**, 30 mol % of Ti(O-*i*Pr)₄, 5 mM in DCM), starting material appeared to be consumed completely within 2 h but the volatile cyclooctenone product **14** was difficult to isolate. Careful removal of the dichloromethane solvent by distillation at atmospheric pressure and then eluting the residue through a polymer-bound thiol SPE tube with methanol afforded the product in modest (estimated 60%) yield. This first-generation synthesis provides proof of concept but delivered a rather

(16) Yamana, M.; Ishihara, T.; Ando, T. *Tetrahedron Lett.* **1983**, 24, 507.

(17) (a) Amii, H.; Kobayashi, T.; Hatamoto, Y.; Uneyama, K. *Chem. Commun.* **1999**, 1323. (b) For a review, see: Uneyama, K.; Amii, H. *J. Fluorine Chem.* **2002**, 114, 127.

(18) Kuroboshi, M.; Ishihara, T. *Bull. Chem. Soc. Jpn.* **1990**, 63, 428.

(19) Felix, C.; Laurent, A.; Mison, P. *J. Fluorine Chem.* **1995**, 70, 71.


(20) Griffith, G. A.; Hillier, I. H.; Percy, J. M.; Roig, R.; Vincent, M. A. *J. Org. Chem.* **2006**, 71, 8250.

(21) Patel, S. T.; Percy, J. M.; Wilkes, R. D. *Tetrahedron* **1995**, 51, 9201.

(22) Patel, S. T.; Percy, J. M.; Wilkes, R. D. *J. Org. Chem.* **1996**, 61, 166.

(23) Broadhurst, M. J.; Brown, S. J.; Percy, J. M.; Prime, M. E. *J. Chem. Soc., Perkin Trans. 1* **2000**, 3217.

CHART 2



volatile product, which was not easy to characterize fully, in ca. 20% overall yield from trifluoroethanol.

We therefore sought to optimize the sequence. The [2,3]-Wittig step afforded good estimated yields (ca. 90%) of up to 24 mmol of homoallylic alcohol **17** which was sufficiently pure by ^1H and ^{19}F NMR spectra and GC-MS to take on directly. Material of the same quality could also be obtained by taking crude undistilled difluoroallylic alcohol through the allylation/rearrangement sequence (after checking the purity from the ^1H and ^{19}F NMR spectra). We esterified **17** by shaking with benzoic anhydride and polymer-supported base poly(4-vinylpyridine) in dichloromethane to afford **18a** and then cleaved the MEM group directly from the crude material. Benzoate **19a** was purified before RCM in the first round of reactions and underwent RCM (second-generation Grubbs' catalyst **21**, $\text{Ti}(\text{O}i\text{-Pr})_4$ cocatalyst) to afford **20a** in acceptable (46%) yield after chromatography. We subsequently found that the unpurified precursor also underwent RCM in good yield. Trace contaminants of Ru could be removed by eluting the columned material through thiol SPE tubes with MeOH. The entire sequence from the starting trifluoroethyl acetal therefore finally involves only a single purification, which is a column after RCM, and the overall yield of **20a** is ca. 30% from trifluoroethanol. Rearrangement product **17** was also benzylated (92%, using an excess of sodium hydride to ensure complete alkoxide formation), subjected to enol ether methanolysis (94%), and taken through the RCM without purification to afford **20b** in good (75%) yield.

Though superficially less attractive than the Ishihara and Uneyama aldol syntheses because it contains two low-temperature steps, our route delivers gram-scale quantities of the RCM product reproducibly with 30 mmol of material coming through the [2,3]-Wittig rearrangement (the most demanding step) and with minimal purification.


Developing Conditions for the RCM Reaction. Many of the RCM reactions²⁴ which form medium rings²⁵ use relatively high loadings of ruthenium catalysts, suggesting that either catalytic efficiency is low or cyclizations are very slow,²⁶ or both. We screened commercial precatalysts **21**–**25** (Chart 2) for the cyclization of **19a** and **19b** at an initial loading of 5 mol %.

(24) For general reviews of RCM issues, see: (a) Conrad, J. C.; Fogg, D. E. *Curr. Org. Chem.* **2006**, *10*, 185. (b) Nicolaou, K. C.; Bulger, P. G.; Sarlah, D. *Angew. Chem., Int. Ed.* **2005**, *44*, 4490. (c) Furstner, A. *Angew. Chem., Int. Ed.* **2000**, *39*, 3013.

(25) For reviews of medium ring forming RCM, see: (a) Deiters, A.; Martin, S. F. *Chem. Rev.* **2004**, *104*, 2199. (b) Maier, M. E. *Angew. Chem., Int. Ed.* **2000**, *39*, 2073. (c) Yet, L. *Chem. Rev.* **2000**, *100*, 2963. (d) Michaut, A.; Rodriguez, J. *Angew. Chem., Int. Ed.* **2006**, *45*, 5740. For recent examples, see: (e) Bleriot, Y.; Giroult, A.; Mallet, J. M.; Rodriguez, E.; Vogel, P.; Sinay, P. *Tetrahedron: Asymmetry* **2002**, *13*, 2553. (f) Bourgeois, D.; Mahuteau, J.; Pancrazi, A.; Nolan, S. P.; Prunet, J. *Synthesis* **2000**, 869. (g) Boyer, F. D.; Hanna, I.; Nolan, S. P. *J. Org. Chem.* **2001**, *66*, 4094. (h) Buono, F.; Tenaglia, A. *J. Org. Chem.* **2000**, *65*, 3869. (i) Furstner, A.; Langemann, K. *J. Org. Chem.* **1996**, *61*, 8746. (j) Buszek, K. R.; Sato, N.; Jeong, Y. *Tetrahedron Lett.* **2002**, *43*, 181.

(26) Illuminati, G.; Mandolini, L. *Acc. Chem. Res.* **1981**, *14*, 95.

SCHEME 5. Attempted RCM with the Neolyst Catalyst



The reaction with Neolyst catalyst **25**²⁷ (10 mol %) returned significant quantities of homodimeric cross-metathesis products from **19a** as a mixture of C_2 -symmetric racemic *syn* and achiral and *meso anti* diastereoisomers **26a** and **27a**, respectively, in moderate (34% isolated from a possible 50%) yield, along with recovered starting material (42%) and traces of RCM product **20a** (Scheme 5).

A similar result was observed from benzyl ether **19b** with the isolation of **26b** and **27b** (38% combined) along with recovered starting material (36%) and a small amount of **20b** (4%). The reaction was started with a lower loading (5%) of catalyst, followed by the addition of a further portion (5%) when the reaction did not appear to proceed. The homodimerization was observed with and without the Ti(IV) cocatalyst.


The isolation of terminal alkene open-chain homodimers from an RCM reaction run at high dilution is relatively unusual.²⁸ The ^1H NMR spectra obtained for the mixtures of **26** and **27** are relatively simple and highly symmetrical, consistent only with the formation of these dimers. The COSYs show the relationship between the vinylic proton of the symmetrical internal alkene and the allylic methylene protons clearly. The connectivity between the terminal alkene and the allylic methine is also established unambiguously. The 1D ^1H and ^{13}C NMR spectra fail to show the presence of the different stereoisomers clearly, though some of the methylene carbon signals are doubled. The 1D $\{^1\text{H}\}^{19}\text{F}$ NMR spectrum appears to show the only difference between the diastereoisomers; formally, there are four ^{19}F environments arising from two pairs of enantiotopic fluorines (two environments) in the achiral and *meso* stereoisomers and two pairs of homotopic fluorines (two environments) in the C_2 -symmetric species (similar considerations apply to various protons, but the higher chemical shift range of fluorine nuclei reveals more subtle differences in environment). Though we observed four environments for **26b/27b**, there were seven for **26a/27a**; given the rather flexible nature of these extended

(27) Furstner, A.; Guth, O.; Duffels, A.; Seidel, G.; Liebl, M.; Gabor, B.; Mynott, R. *Chem.—Eur. J.* **2001**, *7*, 4811.

(28) (a) Creighton and co-workers identified an open-chain dimer by LC-MS from a mixture of products generated during a cyclooctannulation reaction: Creighton, C. J.; Du, Y. M.; Reitz, A. B. *Bioorg. Med. Chem.* **2004**, *12*, 4375. For recent applications, see: (b) Creighton, C. J.; Du, Y. M.; Santulli, R. J.; Toung, B. A.; Reitz, A. B. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 3971. (c) Abell, A. D.; Brown, K. M.; Coxon, J. M.; Jones, M. A.; Miyamoto, S.; Neffe, A. T.; Nikkel, J. M.; Stuart, B. G. *Peptides* **2005**, *26*, 251.

molecules, conformational erosion of symmetry is perhaps unsurprising.

We were surprised by the homodimerization given the reports that this ruthenium indenylidene complex is particularly suitable for the formation of medium-sized rings by RCM. Indeed this (pre)catalyst is more effective at closing the eight-membered E-ring of Nakadomarin A **28** than either Grubbs' catalyst **21** or **22**.²⁷




The effectiveness of Neolyst **25** for slow RCM reactions is attributed to higher stability in solution, though it is not clear if this refers to slow initial dissociation of phosphane, slow recapture of the catalytically active 14e complex, or slow second-order decomposition of the latter reactive species.²⁹ It appears that the RCM is slow for **19a** and **19b** with this precatalyst and that **26** and **27**, once formed, either are too unreactive to form η^2 -complexes with the methyldiene 14e complex which must now carry the chain or cannot progress to metallocyclobutanes from the η^2 -complexes. The presence of phosphane, rather than the NHC ligand, is known to make metallocyclobutane formation less favorable. Relatively rapid consumption and recycling of **26** and **27** would be expected under these conditions. The internal alkene derives from cross metathesis between type I alkenes,³⁰ which is normally fast, so we have observed an interruption of the normally rapid retro-cross metathesis. Exposure of a purified **26b/27b** mixture to 5 mol % of **21** in the presence of the Ti(IV) cocatalyst resulted in 50% conversion to **20b** after 18 h as the sole product by ¹⁹F NMR (the remainder was unreacted starting material). Mixing the crude product with a sample of authentic **20b** and running the GC of the mixture resulted in a single peak being observed. Clearly, the retro-cross metathesis is not rapid under these conditions, even with the more reactive precatalyst **21**.

Catalyst **22** alone will not catalyze RCM of any of our substrates; the presence of the Ti(IV) cocatalyst is required to achieve slow cyclization. Second-generation catalyst **21** is the most effective catalyst explored for the cyclooctannulation reaction, and despite the generally lower Lewis acidity of the alkylidene when the NHC ligand replaces phosphane, the Ti(IV) cocatalyst still exerted a small positive qualitative effect on the reaction outcome. Second-generation Grubbs–Hoveyda catalyst³¹ **24** showed comparable reactivity to **21** (in terms of apparent rate of consumption of **19a**) but afforded a lower yield of eight-membered cyclic products under any given conditions due to oligomer formation, whereas the corresponding first-generation catalyst **23** was ineffective.

Scale-up requires reactions that can be run at concentrations approaching 0.01 M or better, or the volumes of solvent required

SCHEME 6. Dioxirane Oxidations



20a, R = Bz;

20b, R = Bn.

29a, 75%;

29b, 50%.

become prohibitive for normal laboratory equipment. We therefore sought to explore the effects of concentration upon the metathesis outcomes qualitatively. The RCM of benzoate **19a** could be carried out successfully up to 20 mM, with dimerization and other side reactions starting at higher concentrations. This is a relatively high concentration compared to most of those used in the literature, and its use has allowed the preparation of gram quantities of material. Higher dilution was required for benzyl ether **19b** (2.5 mM) and alcohol **13** (1 mM); significant quantities of side products were formed at 20 mM in these cases. Quantitative aspects of these investigations will be presented elsewhere.

Elaboration of RCM Products to Protected Pentopyranose Analogues via Oxidation Transformations. Epoxides **29a** and **29b** were made using the dioxirane method of Yang³² which was diastereofacially selective (Scheme 6). In the case of benzyl ether **20b**, a trace of benzoate **29a** was observed in the product consistent with oxidation at the benzylic methylene by the reactive dioxirane.³³ The sharp ¹⁹F NMR spectra of the crude products showed the formation of a single diastereoisomer in each case with yield loss occurring on purification and removal of the hydrate of the trifluoroacetone. Neither product formed crystals of sufficient quality for the sense of stereoselection to be confirmed directly, but the size of *J* between H-1 and H-2 provides a strong indication; at ca. 9 Hz, it is more likely to arise from a *trans* pseudodiaxial relationship between these protons in electron-deficient environments, consistent with a *trans* relationship between the benzyloxy (or benzyloxy) and epoxide C–O bonds rather than a *cis* arrangement. This conclusion is supported strongly by precedent from previous systems which crystallized well¹⁴ and the outcome of epoxide-opening reactions (*vide infra*).

The results also appear to be consistent with the sense of diastereofacial selection reported by Curci and co-workers³⁴ during the acetone/potassium caroate epoxidation of cyclooctenol and in our preliminary studies; dioxirane attack occurs on the more sterically open face of the alkene (assuming the solution conformer type or population is not modified significantly by the presence of the protecting group).¹³ Hydrogen bond formation to a hydroxyl group clearly cannot play a part in controlling this reaction because the outcome is the same when the hydroxyl group is protected.

Epoxide hydrolysis was carried out under microwave conditions in most cases. We used *N*-methylimidazole as a base

(32) Yang, D.; Wong, M.-K.; Yip, Y.-C. *J. Org. Chem.* **1995**, *60*, 3887.

(33) For examples of related oxidations, see: (a) Csuk, R.; Dorr, P. *Tetrahedron* **1994**, *50*, 9983. (b) Angibeaud, P.; Defaye, J.; Gabelle, A.; Utille, J. P. *Synthesis* **1985**, 1123. (c) Hayes, C. J.; Sherlock, A. E.; Selby, M. D. *Org. Biomol. Chem.* **2006**, *4*, 193. (d) Ochiai, M.; Ito, T.; Takahashi, H.; Nakanishi, A.; Toyonari, M.; Sueda, T.; Goto, S.; Shiro, M. *J. Am. Chem. Soc.* **1996**, *118*, 7716. (e) Arnone, A.; Bernardi, R.; Cavicchioli, M.; Resnati, G. *J. Org. Chem.* **1995**, *60*, 2314.


(34) Cicala, G.; Curci, R.; Fiorentino, M.; Laricchiuta, O. *J. Org. Chem.* **1982**, *47*, 2670.

(29) Sanford, M. S.; Love, J. A.; Grubbs, R. H. *J. Am. Chem. Soc.* **2001**, *123*, 6543.

(30) Chatterjee, A. K.; Choi, T. L.; Sanders, D. P.; Grubbs, R. H. *J. Am. Chem. Soc.* **2003**, *125*, 11360.

(31) (a) Kingsbury, J. S.; Harrity, J. P. A.; Bonitatebus, P. J.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1999**, *121*, 791. (b) For a review, see: Hoveyda, A. H.; Gillingham, D. G.; Van Veldhuizen, J. J.; Kataoka, O.; Garber, S. B.; Kingsbury, J. S.; Harrity, J. P. A. *Org. Biomol. Chem.* **2004**, *2*, 8.

SCHEME 7. Epoxide-Opening Reactions



catalyst in our preliminary study¹⁴ but found subsequently that sodium hydroxide works just as well. In the case of **29a**, microwave irradiation in water alone consumed starting material to afford a mixture of products, which was saponified to afford a single crude triol. The crude product was then treated with an excess (5 equiv) of acetic anhydride in DCM containing poly(vinylpyridine) to afford bisacetate **30** (49% over three steps), for which a crystal structure was obtained, confirming the stereochemistry of the ring-opened product. The pseudoanomeric hydroxyl group is expected to be the least nucleophilic in the triol and was unaffected under these neutral acetylation conditions (Scheme 7).

The hydrolysis of epoxy benzyl ether **29b** was more straightforward. Treatment with a dilute (5%) aqueous solution of NaOH in the microwave resulted in the formation of **31** in good (74%) yield. The structure and stereochemistry of this educt also was confirmed by X-ray crystallography. We note that although the two $^3J_{\text{H-F}}$ couplings visible in the ^{19}F NMR spectrum are consistent with the presence of a pseudoaxial proton next to the CF_2 center (and therefore a chair-type conformation for this ring) one of the couplings is smaller than the values reported for H-C-C-F dihedrals approaching the antiperiplanar angle. These species are now highly oxygenated, which will lower $^3J_{\text{H-F}}$ values generally.

The hydrolytic ring opening of the epoxide is believed to take place via initial reversible attack on the carbonyl followed by irreversible transannular epoxide opening³⁵ with strain relief. Though the addition of nucleophiles to the carbonyl group of cyclooctanone is opposed by the development of additional transannular strain (an *I*-strain effect), the equilibrium constant for cyanohydrin formation from cyclooctanone is still as high as 1.2.³⁶ Fluorination next to a ketonic carbonyl group favors the addition of nucleophiles,³⁷ so there seems to be no difficulty in postulating enough of the hydroxide adduct to trigger a

subsequent strain-relieving reaction,³⁸ if we assume that hydroxide addition is at least as favorable as cyanohydrin formation. The mechanism is supported strongly by the methanolysis of **29b** carried out with catalytic sodium methoxide in methanol under microwave conditions. An inseparable mixture containing two products and starting material (ratio 39:10:1 by ^{19}F NMR) was obtained, but acetylation allowed separation, to afford **32** and **33** (obtained as a mixture with **32**). The presence of the methoxy group at the pseudoanomeric position in **32** is easily detectable in the HMBC spectrum by the presence of a strong $^3J_{\text{C-H}}$ cross-peak between the methoxy protons and C-1. The structure of this major product was also confirmed by the elucidation of the molecular structure in the crystal. The only credible route to major product **32** from **29b** involves methoxide attack at the ketone carbonyl of **29b** followed by transannular ring opening which confirms the sense of diastereofacial selection assigned in Scheme 6.

The formation of **33** was unexpected. We detected a similar $^3J_{\text{C-H}}$ cross-peak between the methoxy protons and C-1 in the HMBC spectrum, which proves that direct epoxide opening by methoxide does not account for the formation of this second product. The 1D NMR spectra showed smaller $^3J_{\text{H-F}}$ coupling constants than usual (the biggest one visible was 9.9 Hz, compared to 17.5 Hz for **32**) suggesting a nonchairlike conformation. One of the ^{19}F NMR signals was very highly split, indicating extended or through-space couplings involving the methoxy group which appears as a doublet in the coupled spectrum and simplifies to a singlet in the $\{^{19}\text{F}\}^1\text{H}$ spectrum. The structure **33** is assigned on the basis of this evidence; it must be formed by transannular nucleophilic ring opening of the epoxide from C-8 rather than from C-7. We were unable to grow suitable crystals of this material. Unfortunately, hydrogenolysis of the benzyl ether to afford **34** failed to deliver crystalline material, but the HMBC spectrum showed a much stronger $\text{CH}_3\text{O}/\text{C-1}$ cross-peak. The full scope of this reaction and its mechanism will be discussed elsewhere.

Dihydroxylation of benzyl ether **20b** under UpJohn conditions³⁹ afforded a mixture of separable diols; both afforded crystals of suitable quality for X-ray crystallographic analysis allowing the identification of the major and minor products as **37** and **38** (3:1), arising from diols **35** and **36** which undergo transannular collapse with relief of strain (Scheme 8).

We were surprised to note that the major product **37** arose from oxidant attack *cis* to the benzyloxy group. In our communication of related work, we advanced an explanation which involved delivery of the osmium reagent to the less accessible concave face of a boat-chair conformer by coordination to the Lewis basic carbonyl group oxygen. In the preliminary case,¹³ we were confident of the identity of the major conformer in solution; however, the ^{19}F NMR spectra of **20a** and **20b** merely broaden at temperatures as low as 213 K preventing conformational insight.

This product is still formed under the conditions developed by Donohoe⁴⁰ which involve the stoichiometric $\text{OsO}_4\cdot\text{TMEDA}$

(35) White, J. D.; Hrciar, P. *J. Org. Chem.* **2000**, *65*, 9129.

(36) Eliel, E. L.; Wilen, S. H.; Mander, L. N. *Stereochemistry of Organic Compounds*; Wiley-Interscience: New York, 1994; p 769.


(37) Gelb, M. H.; Svaren, J. P.; Abeles, R. H. *Biochemistry* **1985**, *24*, 1813.

(38) (a) For a computational study of the prototypical transannular reaction, see: Parveen; Singh, H.; Singh, T. V.; Bharatam, P. V.; Venugopalan, P. *Theochem: J. Mol. Struct.* **2004**, *685*, 139. For a recent elegant synthetic application of a transannular reaction, see: (b) Nicolaou, K. C.; Carenzi, G. E. A.; Jeso, V. *Angew. Chem., Int. Ed.* **2005**, *44*, 3895. (c) Park, C. M. *J. Org. Chem.* **2006**, *71*, 413.


(39) VanRheenen, V.; Kelly, R. C.; Cha, D. Y. *Tetrahedron Lett.* **1976**, 1973.

(40) Donohoe, T. J. *Synlett* **2002**, 1223.


SCHEME 8. Dihydroxylation and Transannular Collapse



SCHEME 9. Benzyl Ether Hydrogenolyses



SCHEME 10. Attempted Selective Phosphorylation



- (i) *n*-BuLi, THF, -78 to 0 °C; or,
(ii) NaHMDS, 15-crown-5, THF, -78 to 0 °C.

complex. In an NMR experiment, we observed an ca. 1:1 mixture of osmate esters⁴¹ before acidic methanolysis and a 1:1 mixture of 37 and 38 after workup. The presence of the chelating TMEDA ligand appears to have made the unexpected pathway less favorable but has not prevented its operation. Donohoe and co-workers have suggested that the diamine and the metal oxide appear to stay bound together under the dihydroxylation conditions, so control via coordination to the carbonyl group seems extremely unlikely under these conditions.

(41) Donohoe, T. J.; Johnson, P. D.; Pye, R. J.; Keenan, M. *Org. Lett.* **2005**, *7*, 1275. This paper describes a well-characterized cyclic osmate ester complexed to TMEDA.




FIGURE 1. Calculated lowest-energy conformations for 39–41. The conformational descriptors refer to the carbon skeleton.

TABLE 1. Energies for RHF 6-31G* Optimized Structures and Calculated Energies (RHF 6-31+G**) for the Lowest-Energy Conformers of 39–41

triol	conformer	$E_{rel}/\text{kcal mol}^{-1}$ (6-31G*)	$E_{rel}/\text{kcal mol}^{-1}$ (6-31+G**)
39	boat-boat	0.594	0.000
39	boat-chair	3.028	2.310
40	boat-boat	0.000	0.276
40	boat-chair	2.118	2.397
41	boat-boat	2.742	3.150
41	boat-chair	4.774	4.126

We also examined the ruthenium-based dihydroxylation described recently by Tiwari and Misra⁴² and found it most effective with rapid conversion of 29b to the bicyclic products after a short reaction time, though with lower stereoselectivity (37/38 1.7:1) than under the UpJohn conditions. The elucidation of the controlling factors in these oxidations will require electronic structure calculations which lie outside the scope of this manuscript.

Unprotected Pentopyranose Analogues and Conformational Analysis. Diols 31, 37, and 38 were debenzylated smoothly under conventional conditions to afford triols 39–41 completing the syntheses of the first three examples of this new class of difluorinated and conformationally locked sugar analogues (Scheme 9). As drawn, 39 represents a locked 2-deoxy-2,2-difluoro analogue of the β -L-lyxo- or β -L-xylopyranoside in the ¹C₄ conformation whereas 41 is the α -anomer in the ⁴C₁ conformation. As drawn, triol 40 is the locked 2-deoxy-2,2-difluoro analogue of β -D-arabino- or ribopyranoside.


The line formula representations of 39–41 conceal a number of close contacts between atoms which will result in Van der Waals strain, notably between 40 and 41 where there are pseudo-1,3-diaxial heavy atoms.

The consequences were explored by carrying out a Monte Carlo conformational search (MMFF94 force field) in Spartan 04.⁴³ The geometries of all the conformers generated in this way were optimized by ab initio calculations (RHF 6-31G*), and energies were calculated using the 6-311+G** basis set. Table 1 shows the relative energies of the lowest-energy

(42) Tiwari, P.; Misra, A. K. *J. Org. Chem.* **2006**, *71*, 2911.

(43) *Spartan 04*, version 1.0.3, Wavefunction Inc.: 18401 Von Karman Avenue, Suite 370, Irvine, CA 92612, 2005.

SCHEME 11. Phosphorylation Following Secondary Hydroxyl Protection



conformers; for all three sugar analogues, these have the functionalized pyran ring in a chair conformation (Figure 1).

Formally, the eight-membered *carbon* skeleta of the bicycles occupy oxygen-bridged boat-boat conformations in the lowest-energy conformers, and the oxygen-bridged boat-chair conformations are both ca. 2 kcal mol⁻¹ higher in energy for **39** and **40** and ca. 1 kcal mol⁻¹ higher in energy for **41**. The boat-boat conformer brings two groups (substituents on C-3 and C-7) very close together. In **39**, for example, H-3 and an H-7 proton are within 2.1 Å, but it also allows each bridging C–O bond to be close to antiperiplanar to a C–H and the exocyclic C–O bond to be antiperiplanar to a C–C bond. In the boat-chair, H-2 and an H-7 proton are within 2.25 Å, and one of the fluorine atoms is within 2.25 Å of H-2. The antiperiplanar relationships with the bridging C–O bonds in the boat-chair are lost.

There is some distortion of structure **40** to allow the pseudo-1,3-diaxial oxygen and fluorine atoms to relieve some Van der Waals strain. The largest ³J_{H–F}s observed were for **40** and the intermediates leading to and derived from it; a similar value was observed for **39**. These values are consistent with the calculated low-energy conformations. Though **41** has the potential to flip the functionalized ring into a boat and exchange the two axial hydroxyl groups into equatorial environments, the ³J_{H–F} suggests that the methine is equatorial and bisects the F–C–F angle, a conclusion supported by the ab initio calculations (the alternate conformer is 1 kcal mol⁻¹ higher in energy).

Synthesis of a Pentopyranosyl Phosphate Analogue. Initially, phosphorylation was attempted without further protection steps. Diol **37** contains two hydroxyl groups which should differ significantly in acidity.⁴⁴ Treating diol **31** with one equivalent of a strong base and allowing the system to equilibrate should favor the formation of **42** at the expense of **43**. The pK_a of the pseudoanomeric hydroxyl group should be lower than the secondary hydroxyl because of the combined inductive effects of the bridging oxygen and the two fluorine atoms; phosphorylation of **42** was therefore anticipated as the kinetic pathway (Scheme 10).⁴⁵

Deprotonation with *n*-BuLi at –78 °C, slow warming to room temperature, and equilibration overnight, followed by the addition of tetrabenzyl pyrophosphate, afforded a mixture of **44** (14%), **45** (3%), and recovered **37** (35%). A higher yield (43%) of the monophosphate **44** could be achieved by changing the base to NaHMDS and adding Na-selective 15-crown-5, consistent with the formation of a more ionic and dissociated

Na salt. However, the poor conversion of **37** and loss of material in **45** directed us toward protection of the secondary hydroxyl group. Acetylation to **46** occurred selectively (89%), and the regiochemistry was confirmed by an HMBC experiment which shows a cross-peak (³J_{C–H}) between the acetate carbonyl carbon and H-4. Exposure of **46** to NaHMDS followed by the addition of tetrabenzyl pyrophosphate allowed the isolation of **47** in good yield, though deacetylated **44** was an occasional contaminant after chromatography. Hydrogenolysis of the benzyl groups was followed by deacetylation, lyophilization, and column chromatography allowing the isolation of deprotected material (Scheme 11). No intermediates were purified during this sequence, but we did check for ¹⁹F and ³¹P NMR spectral changes, obtaining accurate ion masses of the crude materials at each stage. Particularly broad (and rather uninformative) ³¹P NMR spectra were obtained for the free phosphomonoester diacid intermediates.

The final structure **48** is assigned as the ammonium sodium salt on the basis of the presence of ammonia in the chromatographic eluent; chromatography over silica can lead to the formation of monosodium salts, consistent with the results of combustion analysis.

The successful development of a phosphorylation and deprotection strategy sets the stage for the preparation of a wider range of analogues and the chemical or enzymatic synthesis of the NDP sugar analogues themselves.

Conclusions

Telescoped syntheses of locked difluorinated analogues of pentopyranosides have been achieved using metalated difluoroenol acetal chemistry and RCM as key steps in high-yielding multistep sequences which contain minimal purifications. Dihydroxylation allows triol synthesis though with modest stereoselectivity in the case of the all-*cis* triol and subsequent bicyclic hemiacetal. The toxic osmium(VIII) reagent can be replaced by Ru when stereoselectivity is not an issue. Epoxidation and hydrolysis provide a highly stereoselective route to a stereocomplementary class of analogue. A viable phosphorylation strategy involving blocking of the secondary hydroxyl group followed alkoxide formation from the pseudoanomeric hydroxyl group, and subsequent phosphorylation has been developed. Conversion of the sensitive triester to the more robust monoester then allows base-catalyzed deacetylation-completing syntheses of a representative example of a new class of analogue of important biomolecules.

Experimental Section

Representative procedures only are described in this section. Full preparative and characterization details for all other compounds are in the electronic Supporting Information (SI). 2-(2'-Methoxyethoxymethoxy)-1,1,1-trifluoroethane²¹ and 1,1-difluoro-2-(2'-methoxyethoxymethoxy)-4,4-dimethyl-hepta-1,6-dien-3-ol **15**²⁰ were prepared according to published procedures.

(44) (a) For calculations of hemiacetal pK_a (the values are only ca. 2–3 pK_a units lower than those of the analogous alcohols), see: Guthrie, J. P.; Pitchko, V. *J. Am. Chem. Soc.* **2000**, *122*, 5520. (b) An estimate used by Jencks suggests a larger difference of 5 pK_a units: Jencks, W. P. *Catalysis in Chemistry and Enzymology*; McGraw-Hill: New York, 1969; p 522.

(45) The rates of nucleophilic oxyanion attack at phosphotriesters were shown to depend linearly on nucleophile pK_a with a Brønsted coefficient (β_N) of 0.3–0.48 (the size depends on the leaving group). See: Khan, S. A.; Kirby, A. J. *J. Chem. Soc. (B)* **1970**, 1172. We therefore anticipated a competition. Though **42** could be the major species present, **43** could be as much as 2.5 log units more nucleophilic than **42** (assuming a maximum β_N of 0.5). This may explain the relatively low selectivity.

3-Allyloxy-1,1-difluoro-2-(2'-methoxy-ethoxymethoxy)-hepta-1,6-diene 16. A mixture of **15** (68.24 mmol, 17.40 g) and allyl bromide (81.90 mmol, 6.87 mL) was added over 1 min to a vigorously stirred solution of tetra-*n*-butylammonium hydrogen-sulfate (3.45 mmol, 1.16 g) and sodium hydroxide (488 mmol, 24.3 mL of a 50% aqueous solution) at 0 °C. The mixture was stirred at this temperature for 30 min, allowed to warm to room temperature, and stirred for a further 16 h. The yellow white solution was diluted with water (40 mL), and the layers were separated. The aqueous phase was extracted with diethyl ether (4 × 100 mL). The combined organic extracts were washed with brine (2 × 30 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure to afford **16** as a pale yellow oil (18.24 g, 91%, 100% by GC), which was used without any further purification: *R*_f (10% diethyl ether in hexane) 0.42; δ_H (300 MHz, CDCl₃) 5.96–5.83 (m, 1H), 5.81 (ddt, *J* = 16.1, 10.1, 6.6, 1H), 5.28 (ddd, *J* = 17.2, ²*J* = 3.0, ⁴*J* = 1.8, 1H), 5.19 (ddd, *J* = 10.2, ²*J* = 3.0, ⁴*J* = 1.2, 1H), 5.09–4.94 (env., 4H), 4.10 (ddt, *J* = 12.7, 5.1, 1.5, 1H), 4.06–4.00 (m, 1H), 3.94–3.74 (m, 3H), 3.61–3.57 (m, 2H), 3.41 (s, 3H), 2.13–2.06 (m, 2H), 1.95–1.70 (m, 2H); δ_C (75 MHz, CDCl₃) 156.0 (dd, ¹*J*_{C-F} = 291.8, 282.8), 137.7, 134.3, 117.3, 115.1, 112.4 (dd, ²*J*_{C-F} = 36.8, 9.8), 97.1, 73.8, 71.6, 69.3, 68.3, 59.0, 31.0, 29.6; δ_F (282 MHz, CDCl₃) –97.7 (dd, ²*J*_{F-F} = 63.5, ⁴*J*_{F-H} = 1.9, 1F), –109.5 (dd, ²*J*_{F-F} = 63.5, ⁴*J*_{F-H} = 3.8, 1F); ν_{max}(film)/cm⁻¹ 2880s, 1748m, 1642w; *m/z* (FAB) 293 (20%, [M + H]⁺) 215 (24), 165 (74), 137 (90), 89 (100), 59 (84); HRMS (FAB, [M + H]⁺) calcd for C₁₄H₂₃F₂O₄ 293.15634, found 293.15634.

4,4-Difluoro-5-(2-methoxy-ethoxymethoxy)-deca-1,5,9-trien-3-ol 17. A solution of ether **16** (34.25 mmol, 10.00 g) in THF (60 mL) was added dropwise over 15 min to a stirred solution of LDA (prepared from *n*BuLi (68.50 mmol, 28.30 mL of a 2.42 M solution in hexane) and diisopropylamine (75.30 mmol, 10.58 mL) in THF (342 mL)) at –100 °C under a nitrogen atmosphere. The pale pink solution was stirred at this temperature for 30 min before being allowed to warm to –30 °C over 4 h and stirred for a further 3 h at this temperature. The solution changed color during warming, from yellow through orange to brown and finally black. The reaction was quenched with ammonium chloride (10 mL of a saturated aqueous solution), whereupon the black color disappeared and an orange/red solution was observed. The layers were separated, and the aqueous phase was extracted with diethyl ether (3 × 150 mL). The combined organic extracts were washed with brine (2 × 50 mL), dried (MgSO₄), and concentrated under reduced pressure to give the homoallylic alcohol **17** (8.71 g, 95% conversion by NMR and GC) as a red–brown oil, which was used without further purification: *R*_f (30% ether in hexane) 0.12; δ_H (300 MHz, CDCl₃) 5.93 (ddd, *J* = 17.2, 10.5, 5.6, 1H), 5.80 (ddt, *J* = 17.1, 10.1, 6.4, 1H), 5.57 (td, *J* = 7.3, 1.3, 1H), 5.47 (dt, *J* = 17.2, ²*J* = 1.5, 1H), 5.35 (dt, *J* = 10.5, ²*J* = 1.5, 1H), 5.07–4.95 (m, 4H), 4.50 (broad s, 1H), 3.86 (t, *J* = 4.6, 1H), 3.85 (t, *J* = 4.6, 1H), 3.58 (t, *J* = 4.6, 2H), 3.38 (s, 3H), 2.92 (1H, broad s), 2.35–2.25 (m, 2H), 2.20–2.08 (m, 2H); δ_C (75 MHz, CDCl₃) 145.4 (dd, ²*J*_{C-F} = 27.5, ²*J*_{C-F} = 25.2), 135.5, 131.5, 119.8 (t, *J*_{C-F} = 5.4), 118.8, 118.2 (dd, ¹*J*_{C-F} = 250.1, ¹*J*_{C-F} = 247.1), 115.4, 97.2, 72.4 (dd, ²*J*_{C-F} = 30.5, ²*J*_{C-F} = 27.5), 70.5, 67.9, 58.0, 32.0, 23.6; δ_F (282 MHz, CDCl₃) –109.5 (dd, ²*J*_{F-F} = 253.2, *J*_{H-F} = 8.3, 1F), –115.9 (dd, ²*J*_{F-F} = 253.5, *J*_{H-F} = 14.7, 1F); ν_{max}(film)/cm⁻¹ 3434br, 2928w, 1682w, 1641w, 1452w, 1252w, 1170s, 1112s, 1006s, 933s; *m/z* (FAB) 293 (44%, [M + H]⁺), 137 (100), 89 (90), 59 (100); HRMS (FAB, MH⁺) calcd for C₁₄H₂₃F₂O₄ 293.15648, found 293.15648.

3-Benzoyloxy-4,4-difluoro-5-(2'-methoxy-ethoxymethoxy)-deca-1,5Z,9-triene 18a. Alcohol **17** (6.3 g, 21.7 mmol), benzoic anhydride (4.90 g, 21.7 mmol), and DMAP (0.53 g, 4.3 mmol) were dissolved in DCM (217 mL). Poly(vinylpyridine) (10.6 g, 10.6 mmol, 0.5 equiv) was added, and the reaction mixture was swirled gently at room temperature overnight. The resin was collected at the pump and washed with water (50 mL) and saturated NaHCO₃ solution (50 mL). The aqueous layer was extracted with diethyl

ether (3 × 50 mL), and the combined organic extracts were washed with brine, dried, and concentrated in vacuo to yield **18a** as a brown oil (6.85 g, 80%). The crude product was taken on without purification: *R*_f (30% diethyl ether in hexane) 0.27; δ_H (300 MHz, CDCl₃) 8.09–8.05 (m, 2H), 7.58 (apparent tt, *J* = 7.4, ⁴*J* = 1.4, 1H), 7.48–7.42 (m, 2H), 6.01–5.89 (m, 2H), 5.73 (ddd, *J* = 17.0, 10.2, 6.4, 1H), 5.62 (t, *J* = 7.5, 1H), 5.53 (dd, *J* = 16.0, ²*J* = 1.3, 1H), 5.43 (dd, *J* = 10.3, ²*J* = 1.3, 1H), 5.00 (s, 2H), 5.00–4.90 (m, 2H), 3.94–3.80 (m, 2H), 3.60–3.55 (m, 2H), 3.38 (s, 3H), 2.33–2.23 (m, 2H), 2.13–2.04 (m, 2H); δ_C (75 MHz, CDCl₃) 164.8, 144.8 (t, ²*J*_{C-F} = 25.7), 137.3, 133.4, 129.9, 129.5, 128.9 (dd, *J*_{C-F} = 3.6, 1.8), 128.4, 121.4, 120.2 (t, *J*_{C-F} = 5.1), 117.3 (dd, ¹*J*_{C-F} = 251.2, 247.4), 115.4, 98.3, 72.8 (dd, ²*J*_{C-F} = 31.7, 26.9), 71.6, 68.9, 59.0, 33.9, 24.5; δ_F (282 MHz, CDCl₃) –110.8 (dd, ²*J*_{F-F} = 253.2, *J*_{H-F} = 10.9, 1F), –112.4 (dd, ²*J*_{F-F} = 253.2, *J*_{H-F} = 12.3, 1F); ν_{max}(film)/cm⁻¹ 2887w, 1730s, 1602w, 1452w, 1264s, 1096s, 988s, 709s; *m/z* (EI) 396 (3%, M⁺) 355 (7), 220 (39), 205 (100), 145 (21), 106 (68); HRMS (EI, M⁺) calcd for C₂₁H₂₆F₂O₅ 396.17473, found 396.17473.

3-Benzoyloxy-4,4-difluoro-deca-1,9-dien-5-one 19a. Thionyl chloride (15.6 mmol, 1.12 mL) was added to a stirred solution of benzoate **18a** (15.6 mmol, 6.16 g) in methanol (156 mL) at 0 °C. The solution was allowed to warm to room temperature and stirred for 18h, and then the solvent was removed in vacuo. The resulting paste was taken up in water (120 mL), and the mixture was extracted with diethyl ether (5 × 100 mL). The combined organic extracts were washed with NaHCO₃ (200 mL) and brine (200 mL), dried (MgSO₄), filtered, and concentrated in vacuo to give ketone **19a** (100 mg, 75%, 88% by GC) as a brown oil, which could be used crude or purified on silica gel eluted with 10% ethyl acetate in hexane to afford a clear oil (3.60 g, 75%, 97% by GC): *R*_f (10% diethyl ether in hexane) 0.56; δ_H (300 MHz, CDCl₃) 8.06–8.01 (m, 2H), 7.60 (tt, *J* = 7.5, ⁴*J* = 1.3, 1H), 7.49–7.42 (m, 2H), 6.05–5.88 (m, 2H), 5.71 (ddt, *J* = 17.0, *J* = 10.2, *J* = 6.7, 1H), 5.57 (dd, *J* = 16.0, ⁴*J* = 0.9, 1H), 5.50 (dd, *J* = 9.4, ⁴*J* = 0.9, 1H), 5.02–4.94 (m, 2H), 2.74 (t, *J* = 7.3, 2H), 2.05 (q, *J* = 7.3, 2H), 1.71 (pentet, *J* = 7.3, 2H); δ_C (75 MHz, CDCl₃) 199.7 (dd, ²*J*_{C-F} = 28.7, 28.7), 164.4, 137.3, 133.7, 129.9, 128.9, 128.6, 127.7, 122.8, 115.7, 114.1 (dd, ¹*J*_{C-F} = 260.9, 256.1), 72.4 (dd, ²*J*_{C-F} = 29.9, 25.1), 36.7, 32.6, 21.5; δ_F (282 MHz, CDCl₃) –113.7 (dd, ²*J*_{F-F} = 293.9, *J*_{F-H} = 9.0, 1F), –118.9 (dd, ²*J*_{F-F} = 273.9, *J*_{F-H} = 14.2, 1F); ν_{max}(film)/cm⁻¹ 3076w, 2937w, 1733s, 1642w; *m/z* (EI) 308 (37%, M⁺), 105 (85); HRMS (EI, M⁺) calcd for C₁₇H₁₈F₂O₃ 308.12242, found 308.12240.

3-Benzoyloxy-2,2-difluoro-cyclooct-4Z-en-1-one 20a. A solution of diene **19a** (4.07 mmol, 1.26 g) and Ti(O^{*i*}Pr)₄ (1.21 mmol, 0.365 mL) in freshly degassed DCM (407 mL) was refluxed under nitrogen for 30 min, and then Grubbs' catalyst **21** (0.203 mmol, 173 mg, 5 mol %) was added via syringe in DCM (5 mL). Reflux was maintained until the ¹⁹F NMR spectrum of an aliquot showed that starting material had been consumed completely (after 18 h). The solvent was then removed in vacuo, and the residue was taken up in diethyl ether (30 mL) then filtered and concentrated to give crude cyclooctenone as a brown oil which was purified by flash column chromatography (silica gel, 10% diethyl ether in hexane) to give cyclooctenone **20a** which crystallized (0.531 g, 46%, 96% by GC): *R*_f (10% diethyl ether in hexane) 0.20; mp 91–90 °C; δ_H (300 MHz, CDCl₃) 8.14–8.10 (m, 2H), 7.64–7.58 (m, 1H), 7.52–7.45 (m, 2H), 6.36 (dddd, *J*_{H-F} = 21.3, 1.5, *J* = 7.8, ⁴*J* = 3.8, 1H), 6.07–5.96 (m, 1H), 5.65 (ddd, *J* = 11.0, 7.8, 2.5, 1H), 2.82 (dddd, ²*J* = 12.6, *J* = 10.4, 3.9, 2.0, 1H), 2.68 (ddt, ²*J* = 12.6, *J* = 7.2, ⁴*J* = 3.7, 1H), 2.44–2.27 (m, 2H), 2.13–2.02 (m, 1H), 1.89–1.74 (m, 1H); δ_C (75 MHz, CDCl₃) 199.5 (t, ²*J*_{C-F} = 25.5), 165.2, 135.6, 133.7, 130.0, 129.0, 128.6, 125.3, 116.6 (dd, ¹*J*_{C-F} = 262.6, 260.0), 68.2 (dd, ²*J*_{C-F} = 24.2, 18.9), 36.8, 27.5, 27.1; δ_F (282 MHz, CDCl₃) –111.0 (d, ²*J*_{F-F} = 239.8, 1F), –130.9 (dd, ²*J*_{F-F} = 239.8, ³*J*_{F-H} = 21.3, 1F); ν_{max} (solid)/cm⁻¹ 2968m, 2919m, 1725w, 1743w; *m/z* (ES⁺) 281 (M + H⁺, 42%) 121 (PhCOO, 100%). Anal. calcd for C₁₅H₁₄F₂O₃: C, 64.28; H, 5.03.

Found: C, 64.31; H, 5.16. Colorless or almost colorless material of improved quality for further use can be obtained by taking the crude oil (ca. 0.5 g, 2 mmol) up in a minimum volume of methanol and eluting through a preconditioned (3 mL of MeOH per tube) thiol SPE tube with MeOH (3 mL).

3-Benzoyloxy-4,4-difluoro-5-(2'-methoxy-ethoxymethoxy)-deca-1,5Z,9-triene 18b. A solution of **17** (20.2 mmol, 5.9 g) in THF (50 mL) was added cautiously to a suspension of NaH (101 mmol, 4.04 g of a 60% suspension in mineral oil, prewashed with hexane 3 × 30 mL) in THF at 0 °C under nitrogen. The mixture was stirred at this temperature for 45 min as hydrogen evolved. TBAI (2.86 mmol, 1.05 g) then benzyl bromide (19.19 mmol, 2.28 mL) were added, and the mixture was allowed to warm to room temperature over 1 h and then stirred for a further 12 h. The reaction was quenched by the cautious addition of water (150 mL), and the layers were separated. The aqueous layer was extracted with ethyl acetate (3 × 150 mL), and the combined organic extracts were washed with brine (100 mL), dried (MgSO₄), and concentrated in vacuo to give benzyl ether **18b** as a brown oil (7.10 g, 92%, 97% by GC) which was used without further purification: *R_f* (30% ether in hexane) 0.38; δ_H (400 MHz, CDCl₃) 7.25–7.17 (m, 5H), 5.77 (ddd, *J* = 17.5, 10.1, 7.4, 1H), 5.72 (ddt, *J* = 17.1, 10.2, 6.5, 1H), 5.50 (td, *J* = 7.4, ⁴*J*_{H-F} = 1.4, 1H), 5.38–5.36 (m, 1H), 5.35–5.32 (m, 1H), 4.98–4.92 (dq, *J* = 17.1, ⁴*J* = 1.7, 1H), 4.89–4.87 (dq, *J* = 10.2, ⁴*J* = 1.9, 1H), 4.87 (d, ²*J* = 5.9), 4.86 (d, ²*J* = 5.9), 4.58 (d, ²*J* = 11.9, 1H), 4.44 (d, ²*J* = 11.9, 1H), 4.14 (dddd, *J*_{H-F} = 14.2, 8.5, *J* = 7.4, ⁴*J* = 0.9), 3.71 (t, *J* = 4.8, 1H), 3.69 (t, *J* = 4.4, 1H), 3.46 (dd, *J* = 5.1, 4.3, 2H), 3.29 (s, 3H), 2.26–2.18 (m, 2H), 2.09–2.03 (q, *J* = 7.1, 2H); δ_C (100 MHz, CDCl₃) 145.4 (dd, ²*J*_{C-F} = 27.2, 24.8), 137.6, 137.5, 131.1 (dd, *J*_{C-F} = 3.6, 1.8), 128.3, 127.8, 127.7, 121.5, 119.7 (t, *J*_{C-F} = 5.1), 118.1 (dd, ¹*J*_{C-F} = 251.0, 246.0), 115.3, 98.2, 79.0 (dd, ²*J*_{C-F} = 31.7, 25.7), 71.6, 71.4, 68.8, 59.0, 33.1, 24.6; δ_F (282 MHz, CDCl₃) –108.1 (dd, ²*J*_{F-F} = 254.8, *J*_{H-F} = 8.5, 1F), –114.7 (dd, ²*J*_{F-F} = 154.8, *J*_{H-F} = 14.2, 1F); ν_{max}(film)/cm⁻¹ 2923s, 1743w, 1678w, 1637w, 1454s, 1113s, 938s, 850w, 732w, 699s. A satisfactory mass spectrum could not be obtained for this compound (ES-MS, EI, CI, FAB).

3-Benzoyloxy-4,4-difluoro-deca-1,9-dien-5-one 19b. Thionyl chloride (18.6 mmol, 1.34 mL) was added to a stirred solution of enol ether **18b** (18.6 mmol, 7.10 g) in methanol (180 mL) at 0 °C. The solution was allowed to warm to room temperature over 1 h and stirred for 15 h. The methanol was removed in vacuo, and the resulting paste was taken up in water (40 mL) and extracted with ethyl acetate (3 × 150 mL). The combined organic extracts were washed with NaHCO₃ (70 mL) and brine (2 × 100 mL), dried (MgSO₄), and concentrated in vacuo to give ketone **19b** as a yellow oil (5.13 g, 94%, 99% by GC) which was used in the next step without purification. The following data were obtained from a purified sample (flash chromatography, 10% diethyl ether in hexane): *R_f* (30% ether in hexane) 0.68; δ_H (400 MHz, CDCl₃) 7.37–7.21 (m, 5H), 5.85 (ddd, *J* = 17.2, 10.5, 7.6, 1H), 5.72 (ddt, *J* = 17.1, 10.2, 6.7, 1H), 5.53 (ddd, *J* = 10.5, ²*J* = 1.4, ⁴*J* = 0.9, 1H), 5.48 (dt, *J* = 17.2, ²*J* = 1.4, 1H), 4.99 (ddd, *J* = 17.1, ²*J* = 3.4, ⁴*J* = 1.6, 1H), 4.96 (dddd, *J* = 10.2, ²*J* = 3.4, ⁴*J* = 2.0, 1.2, 1H), 4.61 (d, ²*J* = 11.5, 1H), 4.38 (d, ²*J* = 11.5, 1H), 4.26 (dddt, *J*_{H-F} = 16.6, 6.6, *J* = 7.5, ⁴*J* = 0.9, 1H), 2.68 (tt, *J* = 7.3, 1.7, 2H), 2.03 (tdd, *J* = 7.3, 6.7, ⁴*J* = 1.2, 2H), 1.69 (pentet, *J* = 7.3, 2H); δ_C (75 MHz, CDCl₃) 201.7 (dd, ²*J*_{C-F} = 31.1, 25.1), 137.6, 136.8, 129.6 (dd, *J*_{C-F} = 3.6, 1.2), 128.4, 128.1, 128.0, 123.2, 115.4, 115.0 (dd, ¹*J*_{C-F} = 261.5, 253.7), 79.4 (dd, ²*J*_{C-F} = 30.5, 23.9), 71.4, 37.6, 32.7, 21.5; δ_F (282 MHz, CDCl₃) –110.6 (dd, ²*J*_{F-F} = 263.0, *J*_{H-F} = 6.6, 1F), –124.1 (dd, ¹*J*_{F-F} = 263.0, *J*_{H-F} = 16.6, 1F); ν_{max}(film)/cm⁻¹ 1941w, 1740s (C=O), 1642w, 1455w, 1372w, 1217s, 1091s, 913s, 736s, 698s; *m/z* (CI⁺) 312 (100%, [M + NH₄]⁺) 294 (4), 216 (33), 186 (20), 170 (9), 126 (8), 108 (11), 84 (8); HRMS (CI⁺, [M + NH₄]⁺) calcd for C₁₇H₂₄F₂O₂N 312.1770, found 312.1769. Anal. calcd for C₁₇H₂₀F₂O₂: C, 69.37; H, 6.85. Found: C, 69.49; H, 6.98.

3-Benzoyloxy-2,2-difluoro-cyclooct-4Z-en-1-one 20b. A solution of **19b** (2.21 mmol, 0.650 g) and Ti(OⁱPr)₄ (0.66 mmol, 0.198 mL) in freshly degassed DCM (1000 mL) was refluxed for 30 min under nitrogen, and then a solution of Grubbs' catalyst **21** (0.11 mmol, 94 mg, 5 mol %) in DCM (5 mL) was added via syringe. Reflux was maintained until the ¹⁹F NMR spectrum of an aliquot showed that starting material had been consumed completely (after 18h). The solvent was removed in vacuo, and the residue was taken up in diethyl ether (50 mL), then filtered, and concentrated to give crude cyclooctenone **20b** as a brown oil which was purified by flash column chromatography (silica gel, 10% diethyl ether in hexane) to afford cyclooctenone **20b** as a yellow solid (440 mg, 75%, 96% by GC): *R_f* (30% diethyl ether in hexane) 0.40; mp 32–35 °C; δ_H (300 MHz, CDCl₃) 7.40–7.22 (m, 5H), 5.93 (app. q, *J* = 9.2, 1H), 5.60 (ddd, *J* = 11.1, 9.2, 1.2, 1H), 4.76 (d, ²*J* = 12.0, 1H), 4.66 (d, ²*J* = 12.0, 1H), 4.66 (ddd, ³*J*_{H-F} = 20.0, *J* = 8.0, 1.3, 1H), 2.60–2.48 (m, 2H), 2.28 (dddd, *J* = 13.7, 11.1, 5.6, 3.1, 1H), 2.04–1.86 (m, 2H), 1.80–1.46 (m, 1H); δ_C (75 MHz, CDCl₃) 200.4 (dd, ²*J*_{C-F} = 26.6, 24.8), 137.0, 135.4, 128.6, 128.1, 128.0, 127.7 (d, ³*J*_{C-F} = 6.0), 117.8 (dd, ¹*J*_{C-F} = 263.9, 258.5), 173.0 (dd, ²*J*_{C-F} = 23.3, 19.7), 72.1, 36.7, 27.4, 27.1; δ_F (376 MHz, CDCl₃) 200.4 (dd, ²*J*_{F-F} = 240.4, 1F), –130.8 (dd, ²*J*_{F-F} = 240.4, ³*J*_{H-F} = 19.5, 1F); ν_{max}(solid)/cm⁻¹ 2866s, 1743s, 1648w, 1497w, 1455s, 1185s, 1100s, 1070s, 992w, 845s, 812s, 737s, 698s; *m/z* (CI⁺) 284 (100%, [M + NH₄]⁺) 270 (5), 220 (3), 158 (7), 140 (11), 123 (6), 90 (8); HRMS (ES⁺, [M + NH₄]⁺) calcd for C₁₅H₁₆F₂O₂: C, 67.66; H, 6.06. Found: C, 67.59; H, 6.15.

Carrying out the RCM with 7.12 mmol of **19b** at 0.005 M afforded **20b** in 50% yield after purification. The crude material contained significant quantities of higher molecular weight material, so use of the lower concentration is therefore recommended.

2R*-Benzoyloxy-3,3-difluoro-9-oxa-(1S*,8S*)-bicyclo[6.1.0]nonan-4-one 29a. Disodium EDTA (7.2 mL of a 4 × 10⁻⁴ M aqueous solution, 2.9 μmol) followed by trifluoroacetone (3.6 mL of a 60% aqueous solution, 25 mmol from a precooled syringe) was added to a solution of **20a** (0.5 g, 1.8 mmol) in acetonitrile (18 mL) at 0 °C. Sodium hydrogen carbonate (2.3 g, 31.9 mmol) and oxone (5.5 g, 8.6 mmol) were added in one portion. The mixture was stirred for 6 h and allowed to warm to room temperature. The solids were removed by filtration and washed at the pump with DCM (25 mL). The aqueous phase was extracted with DCM (3 × 20 mL), and the combined organic extracts and filter washings were washed with brine (20 mL), dried (MgSO₄), and concentrated in vacuo to yield **29a** as a white solid (0.40 g, 75%) which was used without purification: *R_f* (30% ethyl acetate in hexane) 0.22; mp 75–78 °C; δ_H (300 MHz, CDCl₃) 8.18–8.13 (m, 2H), 7.62 (dt, *J* = 7.3, ⁴*J* = 1.5, 1H), 7.49 (t, *J* = 7.3, 2H), 5.59 (ddd, *J*_{H-F} = 20.9, *J* = 8.9, *J*_{H-F} = 4.4, 1H), 3.24 (ddt, *J* = 8.9, 4.3, ⁴*J* = 1.2, 1H), 3.05 (dt, *J* = 10.6, 4.3, 1H), 2.86–2.65 (m, 2H), 2.49 (dddd, ²*J* = 14.7, *J* = 4.5, 4.3, 3.0, 1H), 2.20–2.10 (m, 1H), 2.05–1.90 (m, 1H), 1.37 (dddd, ²*J* = 14.7, *J* = 13.6, 10.6, 3.4, 1H); δ_C NMR (75 MHz, CDCl₃) 198.9, 164.0, 132.8, 129.2, 127.6, 127.6, 116.7 (t, ¹*J*_{C-F} = 260.9), 68.9, 53.9, 52.0, 35.0, 27.5, 23.2; δ_F (282 MHz, CDCl₃) –113.2 (dd, ²*J*_{F-F} = 247.4, *J*_{H-F} = 4.4, 1F), –127.3 (dd, ²*J*_{F-F} = 247.4, *J*_{H-F} = 20.9, 1F); ν_{max}(solid)/cm⁻¹ 2954w, 1736s, 1275, 1262, 1250 all m, 1081m, 1070m, 707s; *m/z* (EI⁺) 296 (30%, M⁺) 224 (40), 174 (100, M – BzOH); HRMS (EI, M⁺) calcd for C₁₅H₁₄F₂O₄ 296.08602, found 296.08601. Anal. calcd for C₁₅H₁₄F₂O₄: C, 60.8; H, 4.8. Found: C, 60.9; H, 4.8.

2R*-Benzoyloxy-3,3-difluoro-9-oxa-(1S*,8S*)-bicyclo[6.1.0]nonan-4-one 29b. Prepared as for **29a**, from **20b** (0.620 g, 2.33 mmol), Na₂EDTA (9.31 mL of a 4 × 10⁻⁴ M aqueous solution), acetonitrile (23.30 mL), trifluoroacetone (4.66 mL of a 60% aqueous solution, 25 mmol), NaHCO₃ (2.94 g, 34.95 mmol), and oxone (7.16 g, 11.65 mmol) for 6 h at 0 °C. Purification (flash chromatography, silica gel, 20% diethyl ether in hexane) gave epoxide **29b** as a white solid (495 mg, 50%, 98% by GC): *R_f* (30% diethyl ether in hexane) 0.20; mp 70–73 °C; δ_H (400 MHz, CDCl₃) 7.44–7.29 (m, 5H),

4.89 (d, $^2J = 11.9$, 1H), 4.85 (d, $^2J = 11.9$, 1H), 3.68 (ddd, $J_{H-F} = 20.3$, $J = 8.6$, 5.1, 1H), 3.07 (ddt, $J = 8.6$, 4.3, $^4J = 1.6$, 1H), 2.90 (dt, $J = 10.7$, 4.3, 1H), 2.76–2.68 (m, 1H), 2.48 (dddd, $^2J = 13.4$, $J = 9.2$, 4.1, $^4J = 2.2$, 1H), 2.36 (dtd, $J = 14.7$, 4.5, $^2J = 3.1$, 1H), 2.09–1.99 (m, 1H), 1.97–1.85 (m, 1H), 1.02 (dddd, $J = 14.5$, 13.7, 10.6, 3.3, 1H); δ_C (75 MHz, $CDCl_3$) 201.1 (t, $J_{C-F} = 25.7$), 136.8, 128.5, 128.2, 128.1, 115.6 (dd, $^1J_{C-F} = 161.5$, 157.9), 75.3 (dd, $^2J_{C-F} = 22.4$, 18.7), 73.2 (d, $^4J_{C-F} = 1.2$), 54.6 (d, $J_{C-F} = 10.8$), 53.8, 35.8, 28.3, 24.0; δ_F (282 MHz, $CDCl_3$) –113.6 (d, $^2J_{F-F} = 245.9$, 1F), –128.8 (dd, $^2J_{F-F} = 245.9$, $J_{H-F} = 20.3$, 1F); $\nu_{max}(\text{solid})/\text{cm}^{-1}$ 3465w, 2923w, 1744s, 1499w, 1452s, 1340w, 1236w, 1193s, 1105s, 1083s, 1019s, 1028s, 969s, 862s, 830s, 755s, 702s; m/z (CI^+) 300 (100%, $[M + NH_4]^+$) 282 (6), 262 (2), 174 (3), 125 (3), 108 (3); HRMS (ES^+ , $[M + NH_4]^+$) calcd for $C_{15}H_{20}F_2O_3N$ 300.1406, found 300.1409. Anal. calcd for $C_{15}H_{16}F_2O_3$: C, 63.82; H, 5.71. Found: C, 63.77; H, 5.78.

3R*-Benzyloxy-2,2-difluoro-9-oxa-1S*,5R*-bicyclo[3.3.1]nonane-1S*,4R*-diol 31. Epoxide **29b** (0.44 mmol, 125 mg) and NaOH (2.2 mL of a 0.5% aqueous solution, 0.28 mmol) were sealed in a microwave vial containing a stirrer bead. The solution was irradiated in the cavity of a CEM Discover instrument (30 W power to maintain a temperature of 100 °C for 10 min, with a 10 min heating ramp, no cooling). The vial was vented and opened, and the solution was made just acidic (pH 6 to indicator paper) by the addition of a few drops of HCl (0.3 mL of a 3 M aqueous solution), which caused a white solid to precipitate. The aqueous solution was extracted with ethyl acetate (3 × 20 mL), and the combined organic extracts were washed with $NaHCO_3$ (20 mL) and brine (30 mL), dried ($MgSO_4$), and then filtered. The solvent was removed in vacuo to give a white solid which was purified (flash silica, 50% ethyl acetate in hexane) to give **31** as a white solid (98 mg, 74%): R_f (50% ethyl acetate in hexane) 0.28; mp 127–130 °C; δ_H (300 MHz, $CDCl_3$) 7.45–7.32 (m, 5H), 5.01 (dd, $^2J = 11.3$, $^4J = 1.1$, 1H), 4.67 (d, $^2J = 11.3$, 1H), 4.22 (t, $J = 5.1$, 1H), 4.06–3.98 (m, 2H), 3.39 (d, $J = 5.4$, 1H), 2.23 (br. s, 1H), 2.11–2.02 (m, 1H), 1.95–1.52 (m, 5H); δ_C (75 MHz, CD_3OD) 138.0, 127.9, 127.8, 127.4, 118.7 (dd, $^1J_{C-F} = 257.6$, 252.8), 93.6 (dd, $^2J_{C-F} = 26.9$, 20.3), 80.2 (dd, $^2J_{C-F} = 19.1$, 19.1), 74.4 (d, $^4J_{C-F} = 2.4$), 73.0, 71.4 (d, $^3J_{C-F} = 8.4$), 28.7 (d, $^3J_{C-F} = 2.4$), 20.0, 17.9; δ_F (282 MHz, CD_3OD) –115.8 (dd, $^2J_{F-F} = 247.8$, $J_{H-F} = 7.6$, 1F), (–128.7)–(–129.8) (m, incl. app. d, $^2J_{F-F} = 247.8$, 1F); $\nu_{max}(\text{solid})/\text{cm}^{-1}$ 3364br, 2949w, 1350w, 1213, 1080s, 1022s, 907s, 735s, 695s; m/z (CI^+) 318 (100%, $[M + NH_4]^+$) 228 (3), 210 (3), 108 (10), 91 (5), 52 (52); HRMS (ES^+ , $[M + NH_4]^+$) calcd for $C_{15}H_{22}F_2O_4N$ 318.1511, found 318.1510. Anal. calcd for $C_{15}H_{18}F_2O_4$: C, 59.99; H, 6.04. Found: C, 60.13; H, 6.10.

Crystal data: $C_{15}H_{18}F_2O_4$, crystal size 0.19 × 0.10 × 0.04 mm³, $M = 300.29$, triclinic, $a = 9.9359(15)$ Å, $b = 11.1641(17)$ Å, $c = 13.130(2)$ Å, $\alpha = 78.181(3)^\circ$, $\beta = 86.925(3)^\circ$, $\gamma = 88.750(3)^\circ$, $U = 1423.4(4)$ Å³, $T = 150(2)$ K, space group $P1$, $Z = 4$, $\mu(\text{Mo K}\alpha) = 0.117$ mm^{–1}, 10 395 reflections measured, 4966 [R(int) = 0.0818] which were used in all calculations. Final R indices [$F^2 > 2\sigma(F^2)$] R1 = 0.0571, wR2 = 0.0924; R indices (all data) R1 = 0.1002, wR2 = 0.1055.

4R*-Acetoxy-3R*-benzyloxy-2,2-difluoro-1S*-methoxy-9-oxa-1S*,5R*-bicyclo[3.3.1]nonane 32 and 5S*-Acetoxy-3R*-benzyloxy-2,2-difluoro-1S*-methoxy-9-oxa-1S*,4R*-bicyclo[4.2.1]nonane 33. A solution of epoxide **29b** (0.32 mmol, 90 mg) in methanolic sodium methoxide (3.2 mL of a 0.1 M solution in methanol) was irradiated in the microwave as for the hydrolysis of **29b** (30 W, 100 °C for 20 min, 10 min heating ramp, no cooling). The solvent was removed in vacuo, and the residue was taken up in DCM (70 mL) and washed with cold HCl (10 mL of a 1 M solution) and brine (20 mL), dried ($MgSO_4$), then filtered. Poly(vinylpyridine) (340 mg) and acetic anhydride (1.17 mmol, 0.160 mL) were added to the filtrate, and the mixture was swirled at room temperature for 64 h. The poly(vinylpyridine) was removed by filtration and washed with DCM (70 mL). The combined organic extracts and washings were washed with $NaHCO_3$ (10 mL), brine

(20 mL), dried ($MgSO_4$), filtered, and concentrated in vacuo to give a gray paste which was purified (flash chromatography, silica, 20% ethyl acetate in hexane) to give (in order of elution) **33** as a gray paste (20 mg, ca. 18%, 83% of a mixture with **32** by ^{19}F NMR) followed by **32**. **33**: R_f (20% ethyl acetate in hexane) 0.30; δ_H (400 MHz, $CDCl_3$) 7.38–7.30 (m, 5H), 5.15 (dd, $J = 5.7$, 5.3, 1H), 4.84 (d, $^2J = 11.9$, 1H), 4.57 (d, $^2J = 11.9$, 1H), 4.42–4.38 (m, 1H), 4.33 (ddd, $J_{H-F} = 9.9$, 8.5, $J = 3.7$, 1H), 3.49 (d, $^5J_{H-F} = 1.8$, 3H), 2.11–1.96 (m, 2H), 1.92 (s, 3H), 1.70–1.28 (m, 4H); δ_C (100 MHz, $CDCl_3$) 169.6, 136.9, 128.5, 128.1, 128.1, 123.7 (dd, $^1J_{C-F} = 267.2$, 261.6), 105.8 (dd, $^2J_{C-F} = 29.2$, 18.0), 77.9 (d, $J_{C-F} = 9.6$), 77.3 (dd, $^2J_{C-F} = 16.0$, 10.4), 72.4 (d, $^4J_{C-F} = 2.4$), 71.0, 51.8 (d, $^4J_{C-F} = 5.6$), 34.3, 30.1, 20.9, 18.0; δ_F (376 MHz, $CDCl_3$) –114.3 (dd, $^2J_{F-F} = 235.0$, $J_{H-F} = 9.9$, 1F), (–126.0)–(–126.9) (m incl. apparent d, $^2J_{F-F} = 235.0$, 1F); $\nu_{max}(\text{film})/\text{cm}^{-1}$ 2948s, 2359s, 1742s, 1454m, 1372m, 1238s, 1062s, 739w, 699w; m/z (CI^+) 374 (100%, $[M + NH_4]^+$) 284 (5), 208 (6), 106 (9), 77 (18), 52 (79); HRMS (ES^+ , $[M + NH_4]^+$) calcd for $C_{18}H_{26}F_2O_5N$ 374.1774, found 374.1773. In the $\{^{19}F\}^1H$ NMR spectrum, the signal at 4.42–4.38 simplified to 4.40 (dd, $J = 6.5$, 4.0, 1H) and the signal at 3.49 simplified to 3.49 (s, 3H). **32** (45 mg, 39%): R_f (20% ethyl acetate in hexane) 0.16; mp 92–93 °C; δ_H (400 MHz, $CDCl_3$) 7.41–7.32 (m, 5H), 5.21 (ddd, $J = 9.9$, 6.5, 1.2, 1H), 4.96 (d, $^2J = 12.0$, 1H), 4.70 (d, $^2J = 12.0$, 1H), 4.46–4.42 (m, 1H), 4.16 (ddd, $J_{H-F} = 17.4$, 7.6, $J = 9.9$, 1H), 3.51 (d, $^5J_{H-F} = 1.5$, 3H), 2.02 (s, 3H), 1.96–1.60 (m, 6H); δ_C (100 MHz, $CDCl_3$) 169.5, 137.4, 128.4, 128.4, 128.0, 119.2 (dd, $^1J_{C-F} = 259.2$, 257.6), 95.8 (dd, $^2J_{C-F} = 25.6$, 17.6), 77.2 (dd, $^2J_{C-F} = 21.6$, 19.2), 74.1 (d, $^4J_{C-F} = 2.4$), 72.5 (d, $J_{C-F} = 8.8$), 69.6, 50.7 (d, $^4J_{C-F} = 4.0$), 25.9 (d, $J_{C-F} = 2.4$), 21.1, 20.7, 18.1; δ_F (376 MHz, $CDCl_3$) –114.8 (dd, $^2J_{F-F} = 248.3$, $J_{H-F} = 7.1$), –128.0 (dddq, $^2J_{F-F} = 248.3$, $J_{H-F} = 17.5$, $J = 3.8$, $^5J_{H-F} = 1.4$, 1F); $\nu_{max}(\text{solid})/\text{cm}^{-1}$ 2955w, 1737s, 1440w, 1363s, 1239s, 1029s, 892s, 758s; m/z (CI^+) 374 (100%, $[M + NH_4]^+$) 284 (13), 208 (9), 108 (21), 77 (19); HRMS (ES^+ , $[M + NH_4]^+$) calcd for $C_{18}H_{26}F_2O_5N$ 374.1774, found 374.1777. Anal. calcd for $C_{18}H_{22}F_2O_5$: C, 60.67; H, 6.22. Found: C, 60.76; H, 6.30. In the $\{^{19}F\}^1H$ NMR spectrum, the signal at 3.51 simplified to 3.51 (s, 3H).

Crystal data: $C_{18}H_{22}F_2O_5$, crystal size 0.35 × 0.24 × 0.20 mm³, $M = 356.36$, triclinic, $a = 7.240(2)$ Å, $b = 9.607(3)$ Å, $c = 13.226(4)$ Å, $\alpha = 98.732(5)^\circ$, $\beta = 102.590(5)^\circ$, $\gamma = 100.713(5)^\circ$, $U = 864.2(4)$ Å³, $T = 150(2)$ K, space group $P1$, $Z = 2$, $\mu(\text{Mo K}\alpha) = 0.113$ mm^{–1}, 6306 reflections measured, 3024 [R(int) = 0.0485] which were used in all calculations. Final R indices [$F^2 > 2\sigma(F^2)$] R1 = 0.0455, wR2 = 0.1179; R indices (all data) R1 = 0.0554, wR2 = 0.1240.

5S*-Acetoxy-2,2-difluoro-1S*-methoxy-9-oxa-1S*,4R*-bicyclo[4.2.1]nonane-3R*-ol 34. Acetate **33** (0.07 mmol, 25 mg) was dissolved in ethanol (1 mL) containing 10% palladium on activated carbon (10 mg). The atmosphere was removed and replaced with hydrogen from a balloon. The solution was stirred at room temperature for 72 h, and then the hydrogen atmosphere was removed and replaced with air. The solution was filtered through celite, then concentrated in vacuo, and purified by flash chromatography (silica gel, 10–30% ethyl acetate/hexane) to give alcohol **34** as a gray paste (10 mg, 59%): R_f (30% ethyl acetate/hexane) 0.25; $\nu_{max}(\text{film})/\text{cm}^{-1}$ 3447br, 2952w, 1736s, 1441w, 1374w, 1232s, 1036s, 973w, 787w; δ_H (400 MHz, $CDCl_3$) 5.20–5.15 (m, 1H), 4.54 (ddd, $^3J_{H-F} = 11.0$, 8.1, $J = 3.5$, 1H), 4.32 (ddd, $J = 6.4$, 3.5, $^4J_{H-F} = 1.7$, 1H), 3.49 (d, $^5J_{H-F} = 1.4$, 3H), 2.12–1.94 (env., 6H), 1.74–1.60 (m, 3H); δ_C (100 MHz, $CDCl_3$) 170.0, 122.8 (t, $^1J_{C-F} = 262.8$), 105.5 (dd, $^2J_{C-F} = 28.8$, 18.4), 80.0 (d, $^3J_{C-F} = 9.6$), 72.0 (dd, $^2J_{C-F} = 28.0$, 17.6), 71.2, 51.6 (d, $^4J_{C-F} = 4.8$), 33.7–33.6 (m), 29.75–29.70 (m), 21.0, 18.1; δ_F (376 MHz, $CDCl_3$) –119.6 (dd, $^2J_{F-F} = 236.5$, $^3J_{H-F} = 11.1$, 1F), (–126.6)–(–127.5) (m, incl. app. d, $^2J_{F-F} = 236.5$, 1F); m/z (ES^+) 289 (33%, $[M + Na]^+$) 155 (5), 136 (6), 73 (22), 51 (100); HRMS (ES^+ , $[M + NH_4]^+$) calcd for $C_{11}H_{20}F_2O_5N$ 374.1774, found 374.1778. In the $\{^{19}F\}^1H$ NMR spectrum, the signal at 4.32 collapses to 4.32 (dd,

$J = 6.4, 3.5, 1\text{H}$) and the signal at 3.49–3.49 (s, 3H). The signals reported as multiplets in the ^{13}C NMR spectrum are weak and significantly broadened.

3R*-Benzyloxy-2,2-difluoro-9-oxa-1S*,5R*-bicyclo[3.3.1]nona-1S*,4S*-diol 37 and 3R*-Benzyloxy-2,2-difluoro-9-oxa-1R*,5S*-bicyclo[3.3.1]nona-1R*,4R*-diol 38. NMO (295 mg, 2.52 mmol) was added to a solution of cyclooctenone **20b** (336 mg, 1.26 mmol) in acetone (3.16 mL) and H_2O (1.58 mL) at 0 °C. Osmium tetroxide (0.790 mL of a 2.5% by wt. solution in *t*-BuOH, 0.063 mmol) was added, and the black solution was stirred at 0 °C for 6 h. Solid Na_2SO_3 (0.4 g) was added, and the suspension was stirred for 1 h, then diluted with water (3 mL) and extracted with ethyl acetate (3 × 30 mL). The combined organic extracts were washed with brine (30 mL), dried (MgSO_4), filtered, and concentrated in vacuo. Flash chromatography (silica gel, 50–70% ethyl acetate in hexane) gave (in order of elution) minor diastereoisomer **38** (44 mg, 12%) followed by a mixture of **37** and **38** (106 mg, 28%) and then **37** as a white solid. **38**: R_f (50% ethyl acetate in hexane) 0.24; mp 128–131 °C; δ_{H} (300 MHz, CDCl_3) 7.40–7.29 (m, 5H), 4.89 (d, $^2J = 11.5, 1\text{H}$), 4.66 (d, $^2J = 11.5, 1\text{H}$), 4.21 (d, $J = 6.7, 1\text{H}$), 3.97 (dddd, $^3J_{\text{H-F}} = 13.2, J = 6.4, 4.4, 0.9, 1\text{H}$), 3.78–3.71 (m, 1H), 3.44 (d, $^4J_{\text{H-F}} = 6.3, 1\text{H}$), 2.46 (dd, $^4J = 7.9, 1, 2, 1\text{H}$), 2.24–2.06 (m, 2H), 1.92–1.81 (m, 1H), 1.78–1.68 (m, 3H); δ_{C} (75 MHz, CDCl_3) 137.2, 128.6, 128.2, 127.9, 118.2 (dd, $^1J_{\text{C-F}} = 268.1, 251.9$), 94.7 (dd, $^2J_{\text{C-F}} = 29.9, 20.3$), 80.2 (dd, $^2J_{\text{C-F}} = 30.4, 18.0$), 77.0, 75.0 (d, $^4J_{\text{C-F}} = 1.8$), 71.5 (dd, $^3J = 3.0, 3.0$), 29.6, 24.4, 15.9; δ_{F} (282 MHz, CDCl_3) –109.5 (ddd, $^2J_{\text{F-F}} = 259.6, J_{\text{H-F}} = 13.3, 5.5, 1\text{F}$), (–121.8)–(–122.8) (m incl. app. d, $^2J_{\text{F-F}} = 259.6, 1\text{F}$); $\nu_{\text{max}}(\text{solid})/\text{cm}^{-1}$ 3364br, 3166br, 2913s, 1470m, 1351s, 1210s, 1150s, 1072s, 1034s, 943s, 915s, 882m, 819w, 750s, 699s; m/z (CI^+) 318 (100%, $[\text{M} + \text{NH}_4]^+$), 288 (8), 258 (3), 241 (14), 228 (5), 212 (4), 192 (5), 163 (10), 108 (3); HRMS (ES^+ , $[\text{M} + \text{NH}_4]^+$) calcd for $\text{C}_{15}\text{H}_{22}\text{F}_2\text{O}_4\text{N}$ 318.1511, found 318.1512. Anal. calcd for $\text{C}_{15}\text{H}_{18}\text{F}_2\text{O}_4$: C, 59.99; H, 6.04. Found: C, 59.86; H, 5.90.

Crystal data: $\text{C}_{15}\text{H}_{18}\text{F}_2\text{O}_4$, crystal size $0.16 \times 0.13 \times 0.08 \text{ mm}^3$, $M = 300.29$, monoclinic, $a = 10.3624(14) \text{ \AA}$, $b = 6.6363(9) \text{ \AA}$, $c = 11.0444(15) \text{ \AA}$, $\alpha = 90^\circ$, $\beta = 112.613(2)^\circ$, $\gamma = 90^\circ$, $U = 701.11(16) \text{ \AA}^3$, $T = 150(2) \text{ K}$, space group $P2(1)$, $Z = 2$, $\mu(\text{Mo K}\alpha) = 0.119 \text{ mm}^{-1}$, 5059 reflections measured, 2397 $[R(\text{int}) = 0.0672]$ which were used in all calculations. Final R indices $[F^2 > 2\sigma(F^2)]$ $R1 = 0.0465$, $wR2 = 0.0618$; R indices (all data) $R1 = 0.0684$, $wR2 = 0.0683$.

37 (158 mg, 42%): R_f (50% ethyl acetate in hexane) 0.12; mp 109–112 °C; δ_{H} (300 MHz, CDCl_3) 7.41–7.33 (m, 5H), 4.91 (d, $^2J = 11.7, 1\text{H}$), 4.78 (d, $^2J = 11.7, 1\text{H}$), 4.32 (d, $^3J = 6.6, 1\text{H}$), 4.01 (ddd, $J_{\text{H-F}} = 20.1, 7.8, J = 4.8, 1\text{H}$), 3.86–3.80 (m, 1H), 3.64 (d, $^4J = 5.5, 1\text{H}$), 2.95 (s, 1H), 2.08–2.00 (m, 1H), 1.96–1.70 (m, 3H), 1.50–1.37 (m, 2H); δ_{C} (75 MHz, CDCl_3) 136.6, 128.6, 128.5, 128.2, 118.0 (dd, $^1J_{\text{C-F}} = 258.8, 254.6$), 94.1 (dd, $^2J_{\text{C-F}} = 26.9, 20.3$), 74.8, 74.6 (dd, $^2J_{\text{C-F}} = 20.0, 17.6$), 73.3 (d, $^4J_{\text{C-F}} = 1.8$), 71.0 (dd, $^3J_{\text{C-F}} = 7.8, 1.2$), 27.9 (d, $^3J_{\text{C-F}} = 1.8$), 23.0, 18.3; δ_{F} (282 MHz, CDCl_3) (–114.0)–(–114.9) (m incl. app. d, $^2J_{\text{F-F}} = 287.8, 1\text{F}$), –124.1 (dddd, $^2J_{\text{F-F}} = 287.8, ^3J_{\text{H-F}} = 20.1, ^4J_{\text{H-F}} = 5.5, 2.4, 1\text{F}$); $\nu_{\text{max}}(\text{solid})/\text{cm}^{-1}$ 3364br, 3180br, 2902w, 1737s, 1453w, 1343w, 1155s, 1089s, 933s, 867w, 728s, 693s; m/z (CI^+) 318 (100%, $[\text{M} + \text{NH}_4]^+$) 302 (4), 228 (17), 212 (3), 121 (6), 52 (10); HRMS (ES^+ , $[\text{M} + \text{NH}_4]^+$) calcd for $\text{C}_{15}\text{H}_{22}\text{F}_2\text{O}_4\text{N}$ 318.1511, found 318.1515. Anal. calcd for $\text{C}_{15}\text{H}_{18}\text{F}_2\text{O}_4$: C, 59.99; H, 6.04. Found: C, 59.86; H, 5.95.

Crystal data: $\text{C}_{15}\text{H}_{18}\text{F}_2\text{O}_4$, crystal size $0.14 \times 0.09 \times 0.06 \text{ mm}^3$, $M = 300.29$, monoclinic, $a = 15.5415(19) \text{ \AA}$, $b = 6.6332(8) \text{ \AA}$, $c = 13.9404(17) \text{ \AA}$, $\alpha = 90^\circ$, $\beta = 106.126(2)^\circ$, $\gamma = 90^\circ$, $U = 1380.6(3) \text{ \AA}^3$, $T = 150(2) \text{ K}$, space group $P2(1)/c$, $Z = 4$, $\mu(\text{Mo K}\alpha) = 0.121 \text{ mm}^{-1}$, 9657 reflections measured, 2430 $[R(\text{int}) = 0.0967]$ which were used in all calculations. Final R indices $[F^2 > 2\sigma(F^2)]$ $R1 = 0.0514$, $wR2 = 0.0661$; R indices (all data) $R1 = 0.1050$, $wR2 = 0.0785$. Crude cyclooctenone **20b** could be used in the dihydroxylation reaction to give diols **37** and **38** in 61% combined yield over two steps from purified RCM precursor **19b**.

2,2-Difluoro-9-oxa-1S*,5R*-bicyclo[3.3.1]nona-1S*,3R*,4R*-triol 39. Hemiacetal **31** (0.080 mmol, 24 mg) was dissolved in ethanol (1 mL) containing 10% Pd–C (5 mg). The atmosphere was removed and replaced several times by hydrogen from a double balloon, and then the reaction was stirred at room temperature for 23 h. The hydrogen atmosphere was removed in vacuo and replaced with air, and then the catalyst was removed by filtration through celite. Concentration of the filtrate in vacuo afforded **39** (16 mg, 95%): R_f (100% ethyl acetate) 0.31; mp 156–158 °C; δ_{H} (300 MHz, CD_3OD) 4.06–3.92 (m, 2H), 3.76 (dd, $^3J = 9.5, 6.4, 1\text{H}$), 1.92–1.48 (env., 6H); δ_{C} (100 MHz, CD_3OD) 117.9 (dd, $^1J_{\text{C-F}} = 156.4, 150.9$), 93.5 (dd, $^2J_{\text{C-F}} = 26.8, 20.4$), 73.0, 72.9 (dd, $^2J_{\text{C-F}} = 20.4, 20.4$), 72.0 (d, $^3J_{\text{C-F}} = 8.0$), 28.7 (d, $^3J_{\text{C-F}} = 2.4$), 20.0, 18.9; δ_{F} (282 MHz, CD_3OD) –118.2 (dd, $^2J_{\text{F-F}} = 246.7, J_{\text{H-F}} = 8.2, 1\text{F}$), –129.4 (ddd, $^2J_{\text{F-F}} = 246.7, J_{\text{H-F}} = 19.4, ^4J_{\text{H-F}} = 4.3, 1\text{F}$); $\nu_{\text{max}}(\text{solid})/\text{cm}^{-1}$ 3296br, 2964w, 1440w, 1345w, 1207m, 1116m, 1034s, 996s, 929s, 823s; m/z (CI^-) 209 (30%), $[\text{M} - \text{H}]^-$ 191 (11), 170 (18), 152 (15), 79 (22); HRMS (CI^- , $[\text{M} - \text{H}]^-$) calcd for $\text{C}_8\text{H}_{11}\text{F}_2\text{O}_4$ 209.0631, found 209.0630. Anal. calcd for $\text{C}_8\text{H}_{12}\text{F}_2\text{O}_4$: C, 45.72; H, 5.75. Found: C, 45.68; H, 5.70.

2,2-Difluoro-9-oxa-1S*,5R*-bicyclo[3.3.1]nona-1S*,3R*,4S*-triol 40. From **37** (0.077 mmol, 23 mg), 10% Pd–C (5 mg) in ethanol (1 mL) over 72 h. Filtration through celite and concentration afforded triol **40** (36 mg, 100%): R_f (100% ethyl acetate) 0.15; mp 153–155 °C; δ_{H} (300 MHz, CD_3OD) 4.17–4.04 (env. 2H, containing 4.10 (ddd, $J_{\text{H-F}} = 21.6, ^3J_{\text{H-F}} = 8.6, J = 4.8, 1\text{H}$) and 4.11–4.09 (m, 1H)), 3.72 (broad s, 1H), 1.90 (broad d, $J = 9.6, 1\text{H}$), 1.80–1.44 (m, 5H); δ_{C} (100 MHz, CD_3OD) 117.9 (dd, $^1J_{\text{C-F}} = 254.0, 254.0$), 93.8 (dd, $^2J_{\text{C-F}} = 26.8, 20.4$), 75.9, 72.0 (dd, $^3J_{\text{C-F}} = 8.0, 1.6$), 68.8 (dd, $^2J_{\text{C-F}} = 21.6, 19.2$), 29.0 (d, $^3J_{\text{C-F}} = 2.4$), 23.0, 17.7; δ_{F} (282 MHz, CD_3OD) (–119.3)–(–120.2) (m incl. app. d, $^2J_{\text{F-F}} = 247.4, 1\text{F}$), –127.3 (ddd, $^2J_{\text{F-F}} = 247.4, ^3J_{\text{H-F}} = 21.3, ^4J_{\text{H-F}} = 3.8, 1\text{F}$); $\nu_{\text{max}}(\text{solid})/\text{cm}^{-1}$ 3346br, 2951w, 1647w, 1444w, 1353w, 1204s, 1076s, 1037s, 928s; m/z (CI^+) 228 (100%, $[\text{M} + \text{NH}_4]^+$) 123 (6); HRMS (ES^+ , $[\text{M} + \text{NH}_4]^+$) calcd for $\text{C}_8\text{H}_{16}\text{F}_2\text{O}_4\text{N}$ 228.1042, found 228.1038. Anal. calcd for $\text{C}_8\text{H}_{12}\text{F}_2\text{O}_4$: C, 45.72; H, 5.75. Found: C, 45.79; H, 5.80.

2,2-Difluoro-9-oxa-1R*,5S*-bicyclo[3.3.1]nona-1R*,3R*,4R*-triol 41. From **38** (0.067 mmol, 20 mg), 10% palladium-on-carbon (5 mg) in ethanol (1 mL) over 23 h. Filtration through celite and concentration in vacuo afforded triol **41** (25 mg, 91%): R_f (100% ethyl acetate) 0.29; mp 57–60 °C; δ_{H} (300 MHz, CD_3OD) 3.97 (broad s, 1H), 3.86 (ddd, $J_{\text{H-F}} = 13.4, 12.5, J = 6.9, 1\text{H}$), 3.53 (dt, $J = 6.9, 2.7, 1\text{H}$), 2.00–1.90 (m, 1H), 1.72–1.38 (m, 5H); δ_{C} (400 MHz, CD_3OD) 118.1 (dd, $^1J_{\text{C-F}} = 258.0, 258.0$), 94.2 (dd, $^2J_{\text{C-F}} = 31.2, 20.0$), 77.0, 72.2 (dd, $^2J_{\text{C-F}} = 28.8, 19.2$), 70.6 (dd, $^3J_{\text{C-F}} = 6.0, 2.0$), 30.2, 24.8, 15.4; δ_{F} (282 MHz, CD_3OD) –112.7 (ddd, $^2J_{\text{F-F}} = 253.5, J = 12.5, ^4J_{\text{H-F}} = 3.8, 1\text{F}$), –125.1 (dd, $^2J_{\text{F-F}} = 253.5, J_{\text{H-F}} = 13.4, 1\text{F}$); $\nu_{\text{max}}(\text{solid})/\text{cm}^{-1}$ 3289br, 2963w, 1351m, 1205m, 1092s, 1000s, 958s, 894s; m/z (CI^+) 228 (100%, $[\text{M} + \text{NH}_4]^+$); HRMS (ES^+ , $[\text{M} + \text{NH}_4]^+$) calcd for $\text{C}_8\text{H}_{16}\text{F}_2\text{O}_4\text{N}$ 228.1042, found 228.1038. Anal. calcd for $\text{C}_8\text{H}_{12}\text{F}_2\text{O}_4$: C, 45.72; H, 5.75. Found: C, 45.84; H, 5.88.

4S*-Acetoxy-3R*-benzyloxy-2,2-difluoro-9-oxa-1S*,5R*-bicyclo[3.3.1]nonan-1S*-ol 46. Acetic anhydride (0.214 mL, 2.26 mmol), DMAP (16.5 mg, 0.14 mmol), and poly(vinylpyridine) (0.9 mmol, 450 mg at 2.0 mmol per gram of loading) were added to a solution of diol **37** (136 mg, 0.45 mmol) in DCM (4.5 mL). The suspension was shaken at room temperature for 75 h. TLC analysis showed the reaction was incomplete, so additional acetic anhydride (200 μL , 2.11 mmol), DMAP (20 mg, 0.17 mmol), and poly(vinylpyridine) (320 mg) were added. The reaction was shaken at room temperature for a further 48 h until consumption of starting material was observed by TLC. Workup as before afforded **46** (138 mg, 89%) as a white solid: R_f (50% ethyl acetate in hexane) 0.34; mp 28–30 °C; δ_{H} (300 MHz, CDCl_3) 7.39–7.30 (m, 5H), 5.20 (ddd, $J = 4.9, ^4J_{\text{H-F}} = 3.8, J = 1.5, 1\text{H}$), 4.76 (d, $^2J = 12.4, 1\text{H}$), 4.73 (d, $^2J = 12.4, 1\text{H}$), 4.22 (d, $J = 6.7, 1\text{H}$), 4.04 (ddd, $J_{\text{H-F}} = 21.8, 7.3, J = 5.0, 1\text{H}$), 3.61 (d, $J = 6.4, 1\text{H}$), 2.15 (s, 3H), 2.06–1.38

(m, 6H); δ_C (100 MHz, $CDCl_3$) 170.6, 136.7, 128.6, 128.3, 128, 117.4 (dd, $^1J_{C-F} = 256.4, 255.6$), 94.3 (dd, $^2J_{C-F} = 26.8, 20.4$), 73.8–73.2 (m), 70.8 (dd, $^3J_{C-F} = 9.2, 1.6$), 27.8 (d, $^3J_{C-F} = 1.6$), 23.0, 21.0, 18.2; δ_F (282 MHz, $CDCl_3$) (–116.8)–(–117.8) (m, incl. app. d, $^2J_{F-F} = 245.9, 1F$), –127.0 (dd, $^2J_{F-F} = 245.8, J_{H-F} = 21.8, ^4J_{H-F} = 5.2, 1F$); ν_{max} (film)/ cm^{-1} 3434br, 2951s, 1730s, 1367s, 1074s, 908s, 733s; m/z (ES^+) 342 (2%, $[M + H]^+$) 176 (13), 116 (83), 91 (100), 43 (61); HRMS (ES^+ , $[M + H]^+$) calcd for $C_{17}H_{21}F_2O_5$ 343.1352, found 343.1356. Anal. calcd for $C_{17}H_{20}F_2O_5$: C, 59.64; H, 5.89. Found: C, 59.50; H, 5.73.

The ^{13}C NMR spectrum contained a number of weak signals in the 73.8–73.2 ppm region which could not be resolved well, hence the recording of this signal as a multiplet.

4R*-Acetoxy-3R*-benzyloxy-1R*-(dibenzylphosphoryloxy)-2,2-difluoro-9-oxa-1R*,5S*-bicyclo[3.3.1]nonane 47 and 3R*-Benzyloxy-1R*-(dibenzylphosphoryloxy)-2,2-difluoro-9-oxa-1R*,5S*-bicyclo[3.3.1]nonan-4R*-ol 44. NaHMDS (0.54 mmol, 317 μL of a 1.7 M solution in THF) was added dropwise to a solution of **46** (0.49 mmol, 168 mg) in THF (10 mL) at 0 °C and stirred at this temperature for 1 h. Tetrabenzyl pyrophosphate (0.54 mmol, 290 mg) was added, and the reaction was allowed to warm to room temperature over 2 h, then stirred for 18 h after which a white precipitate was observed. The reaction was quenched with pH 7 buffer (10 mL) and extracted with ethyl acetate (2 \times 50 mL). The combined organic extracts were washed with brine (20 mL), dried ($MgSO_4$), filtered, and concentrated in vacuo to give a gray paste which was purified (flash chromatography, silica, 50% ethyl acetate in hexane) to afford **47** (188 mg, 64%): R_f (ethyl acetate) 0.60; δ_H (300 MHz, $CDCl_3$) 7.38–7.28 (m, 15H), 5.22–5.18 (m, 1H), 5.17–5.05 (m, 4H), 4.75 (s, 2H), 4.35 (d, $^3J = 6.6, 1H$), 4.05 (ddd, $^3J_{H-F} = 20.5, J = 7.0, 5.0, 1H$), 2.56–2.40 (m, 1H), 2.16–1.79 (envelope, 6H), 2.06 (s, 3H), 1.52–1.36 (m, 2H); δ_C (100 MHz, $CDCl_3$) 170.5, 136.6, 136.0 (d, $^3J_{C-P} = 8.8$), 135.7 (d, $^3J_{C-P} = 8.0$), 128.6, 128.5, 128.5, 128.4, 128.4, 128.3, 128.1, 128.0, 127.9, 116.0 (ddd, $^2J_{F-F} = 264.7, 256.9, ^3J_{C-P} = 6.8$), 99.8 (ddd, $^2J_{C-F} = 27.2, 18.4, ^2J_{C-P} = 7.2$), 75.8, 73.3, 73.0 (ddd, $^2J_{C-F} = 21.6, 17.6, ^4J_{C-P} = 1.6$), 70.4 (d, $^3J_{C-F} = 9.6$), 69.7 (dd, $^2J_{C-P} = 6.4, ^6J_{C-F} = 1.6$), 69.4 (d, $^2J_{C-P} = 6.4$), 27.8, 22.6, 20.9, 18.4; δ_F (282 MHz, $CDCl_3$) (–117.7)–(–118.5) (m, 1F), –124.3 (ddd, $^2J_{F-F} = 245.9, J_{H-F} = 20.5, J = 5.2, 1F$); δ_P (121 MHz, $CDCl_3$) –8.8 (quintet, $^3J_{H-P} = 7.3$); ν_{max} (film)/ cm^{-1} 3472w, 2955s, 1738s, 1496s, 1455s, 1371s, 1243s, 1017s, 873w, 738s; m/z (CI^+) 603 (28%, $[M + H]^+$) 513 (5), 360 (7), 125 (13), 108 (51), 106 (100); HRMS (ES^+ , $[M + H]^+$) calcd for $C_{31}H_{34}F_2O_8P$ 603.1954, found 603.1955. Traces of deprotected **44** (16 mg, 6%) were also produced, as reported previously.

2,2-Difluoro-3R*,4S*-dihydroxy-9-oxa-1S*,5R*-bicyclo[3.3.1]nonanyl-1S*-phosphate Ammonium Sodium Salt 48. Hydrogenolysis. 10% Palladium-on-carbon (64 mg) was added to a solution of **47** (0.28 mmol, 170 mg) in ethanol (5.6 mL). The apparatus was pump-purged with hydrogen from a double balloon, and the reaction was stirred under hydrogen at room temperature

for 90 h. The solution was filtered through celite, and the filtrate was concentrated in vacuo (98 mg, 100%): $\{^1H\}\delta_F$ (282 MHz, CD_3OD) –116.3 (d, $^2J_{F-F} = 243.2, 1F$), –121.2 (d, $^2J_{F-F} = 243.2, 1F$); δ_P (121 MHz, CD_3OD) (+8)–(–12) (br. m); m/z (ES^-) 331 (100%, $[M - H]^-$) 289 (15), 167 (17), 89 (46), 75 (44); HRMS (ES^- , $[M - H]^-$) calcd for $C_{10}H_{14}F_2O_8$ 331.0400, found 331.0396.

Acetate Cleavage and Bis(triethylammonium) Salt Formation. The crude acid (0.28 mmol, 98 mg) was taken up in a mixture of methanol, water, and triethylamine (5.9 mL, 5:2:1) and stirred at room temperature for 22 h. The organic solvents were removed in vacuo, then the residue was freeze-dried to afford the crude bis-(triethylammonium) salt: δ_F (282 MHz, CD_3OD) –119.7 (d, $^2J_{F-F} = 245.0, 1F$), –124.2 (dd, $^2J_{F-F} = 245.0, J_{H-F} = 20.8, 1F$); δ_P (121 MHz, CD_3OD) –3.9 (s); m/z (ES^+) 493 (58%, $[M + H]^+$) 392 (97), 242 (5), 102 (100), 74 (35); HRMS (ES^+ , $[M + H]^+$) calcd for $C_{20}H_{43}F_2N_2O_7P$ 493.2849, found 493.2851.

Purification and Ammonium Sodium Salt Formation. Flash chromatography (silica, ethanol/water/35% aqueous ammonia (5:3:1)) afforded ammonium sodium salt **48** (60 mg, 66%): R_f (ethanol/water/35% aqueous ammonia (5:3:1)) 0.13; mp 137–139 °C; δ_H (400 MHz, D_2O) 4.35 (ddd, $J_{H-F} = 21.5, 8.1, J = 4.8, 1H$), 4.33–4.31 (m, 1H), 3.98–3.92 (m, 1H), 2.28 (ddd, $^2J = 14.2, J = 14.0, 7.3, 1H$), 2.06 (dd, $^2J = 14.2, J = 5.4, 1H$), 1.95–1.75 (m, 2H), 1.71–1.60 (m, 1H), 1.57–1.50 (dd, $J = 14.0, 5.3, 1H$); δ_C (100 MHz, D_2O) 117.4 (dd, $^1J_{C-F} = 256.4, 250.1$), 97.2 (ddd, $^2J_{C-F} = 25.6, 18.4, ^2J_{C-P} = 7.2$), 76.8, 71.4 (dd, $^3J_{C-F} = 8.0, 1.6$), 68.4 (dd, $^2J_{C-F} = 20.8, 18.4$), 27.6, 22.3, 17.5; δ_F (282 MHz, D_2O) (–118.4)–(–119.4) (m, incl. app. d, $^2J_{F-F} = 243.1, 1F$), –123.7 (ddd, $^2J_{F-F} = 243.1, J_{H-F} = 21.5, ^4J_{H-F} = 5.9, 1F$); δ_P (121 MHz, D_2O) –4.1 (s); ν_{max} (film)/ cm^{-1} 2952br, 1444w, 1361w, 1167m, 1078s, 1040s, 910s, 808s, 751s, 685w; m/z (ES^-) 289 (4%, $[M - H]^-$) 273 (5), 183 (3), 125 (56), 97 (100); HRMS (ES^- , $[M - H]^-$) calcd for $C_8H_{12}F_2O_7P$ 289.0294, found 289.0297. Anal. calcd for $C_8H_{15}F_2NO_7NaP$: C, 29.18; H, 4.56; N, 4.26. Found: C, 29.48; H, 4.32; N, 4.27.

Acknowledgment. We thank the EPSRC for studentships (J.A.L.M., L.M.), the EPSRC Mass Spectrometry Service (Swansea) for accurate mass measurements, Dr. Roland Wende (Umicore) for a donation of the Neolyst catalyst, and Dr. A. Caravano (Sanofi-Aventis) and Professor R. A. Field (University of East Anglia) for helpful discussions.

Supporting Information Available: Experimental procedures for **11a**, **11b**, **13**, **14**, **26a–27b**, **30**, **44**, and **45**; NMR spectra (1H , ^{13}C , ^{19}F , ^{31}P) for **11a**, **11b**, **13**, **14**, **16–19a**, **26a–27b**, **33**, **34**, **44**, **45**, and **47**; Cartesian coordinates and energies for RHF 6-31G* optimized structures for lowest-energy conformers of **39–41**, and calculated energies (RHF 6-31+G**) for **39–41**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO0620258