

# Bivariate Spline Interpolation with Optimal Approximation Order\*

O. Davydov,<sup>†</sup> G. Nürnberger and F. Zeilfelder

## Abstract

Let  $\Delta$  be a triangulation of some polygonal domain  $\Omega \subset \mathbb{R}^2$  and let  $S_q^r(\Delta)$  denote the space of all bivariate polynomial splines of smoothness  $r$  and degree  $q$  with respect to  $\Delta$ . We develop the first Hermite type interpolation scheme for  $S_q^r(\Delta)$ ,  $q \geq 3r + 2$ , whose approximation error is bounded above by  $Kh^{q+1}$ , where  $h$  is the maximal diameter of the triangles in  $\Delta$ , and the constant  $K$  only depends on the smallest angle of the triangulation and is independent of near-degenerate edges and near-singular vertices. Moreover, the fundamental functions of our scheme are minimally supported and form a locally linearly independent basis for a superspline subspace of  $S_q^r(\Delta)$ . This shows that the optimal approximation order can be achieved by using minimally supported splines. Our method of proof is completely different from the quasi-interpolation techniques for the study of the approximation power of bivariate splines developed in [7, 18].

---

\*1991 *Mathematics Subject Classification*. Primary 41A25, 41A63; Secondary 41A05, 41A15, 65D05, 65D07.

*Key words and phrases*. Bivariate spline, triangulation, interpolation, approximation order, minimally supported basis, locally linearly independent basis.

<sup>†</sup>Supported in part by a Research Fellowship from the Alexander von Humboldt Foundation

# 1 Introduction

Let  $\Omega \subset \mathbb{R}^2$  be a polygonal domain, and let  $\Delta$  denote a regular triangulation of  $\Omega$ . The space of bivariate polynomial splines of degree  $q$  and smoothness  $r$  with respect to  $\Delta$  is defined by

$$S_q^r(\Delta) := \{s \in C^r(\Omega) : s|_T \in \Pi_q \text{ for all } T \in \Delta\}, \quad 0 \leq r < q,$$

where

$$\Pi_q := \text{span} \{x^i y^j : i \geq 0, j \geq 0, i + j \leq q\}$$

is the space of bivariate polynomials of total degree  $q$ .

In the literature, point sets that admit unique Lagrange and Hermite interpolation by spaces  $S_q^r(\Delta)$  of splines of degree  $q$  and smoothness  $r$  were constructed for crosscut partitions  $\Delta$ , in particular for  $\Delta^1$  and  $\Delta^2$ -partitions [1, 6, 17, 22, 23, 24, 27, 28]. Results on the approximation order of these interpolation methods were given in [6, 12, 17, 21, 22, 25, 27, 28].

In the case of an arbitrary triangulation  $\Delta$ , standard finite-element techniques [9] provide a tool to construct Hermite type interpolation schemes for  $S_q^r(\Delta)$  with optimal approximation order  $\mathcal{O}(h^{q+1})$ , where  $h$  is the maximal diameter of the triangles in  $\Delta$ . However, as shown in [30], this method only works if  $q \geq 4r + 1$ .

On the other hand, the approximation power of the spline space  $S_q^r(\Delta)$  for  $q \geq 3r + 2$  was studied in [3, 7, 8, 18]. It was shown in [7] that for a sufficiently smooth function  $f$ ,

$$(1.1) \quad \text{dist}(f, S_q^r(\Delta)) \leq K h^{q+1},$$

where  $K$  is a constant that depends only on  $f$ ,  $r$ ,  $q$  and the smallest angle  $\theta_\Delta$  in  $\Delta$ . In the earlier constructions in [3, 8] the constant  $K$  is bounded under additional assumption that the triangulation has no near-singular vertices. Another proof of (1.1) was recently suggested in [18]. If  $q < 3r + 2$ , then the optimal approximation order fails for certain triangulations (see [4]).

The problem of constructing a Hermite interpolation operator satisfying (1.1) was already brought up in [8]. However, the question if the constant  $K$  for such an operator could be bounded independently of near-singularities remained open.

It is the purpose of this paper to develop the first Hermite type interpolation scheme for  $S_q^r(\Delta)$ ,  $q \geq 3r + 2$ , that possesses optimal approximation order in the sense that the corresponding constant  $K$  in (1.1) is independent of near-degenerate edges and near-singular vertices. The details of our construction are given in Section 2, whereas the main result of the paper, Theorem 3.1 about the approximation order,

as well as its proof are presented in Section 3. Finally, Section 4 is devoted to an analysis of the fundamental functions of our scheme that form a basis for a superspline subspace of  $S_q^r(\Delta)$ .

We emphasize that our approach is completely different from that of [7] and [18]. In each of these papers a stable local basis for a superspline subspace of  $S_q^r(\Delta)$  was constructed first by using Bernstein-Bézier techniques, and then the basis functions were used to build up a quasi-interpolation operator that yielded the optimal approximation order. In contrast to this, we argue directly with nodal functionals and develop a new approach that has its roots in the idea of “weak interpolation” introduced in [21] and further developed in [25] and [12]. Furthermore, we need a new description of  $C^r$  smoothness across edges in terms of nodal functionals (see Lemma 3.2). As a result, we manage to exploit linear dependencies between directional derivatives of smooth functions to suppress the instability of the minimally supported fundamental functions in the neighborhood of a near-singular vertex, see Lemma 3.3 and the discussion at the end of Section 4. Note that in [7, 18] much more complicated basis functions with larger supports were constructed in order to achieve stability.

Among other advantages of our approach is that the estimation of the approximation error (3.2) is completely local, *i.e.*, it involves only the local Sobolev norm of  $f$  on the triangle  $T$  on which the error is measured. Moreover, the result remains valid for any nonconvex polygonal domain  $\Omega$  with the constant  $K$  independent of the Lipschitz constant of the boundary of  $\Omega$ .

As a by-product of our construction, we get a nodal basis for the space of supersplines

$$S_q^{r,\rho}(\Delta) := \{s \in S_q^r(\Delta) : s \in C^\rho(v) \text{ for all vertices } v \text{ of } \Delta\},$$

where  $\rho = r + \lceil \frac{r+1}{2} \rceil$  and  $q \geq 3r + 2$ . The basis consists of the fundamental functions  $s_1, \dots, s_n$  of our interpolation scheme. Some properties of this basis are studied in Section 4. Namely, it is shown that  $\{s_1, \dots, s_n\}$  is *locally linearly independent* and thus *least supported* [5], *i.e.*, the supports of the basis functions  $s_i$  are as small as possible, which is not the case for the basis functions constructed in [7, 18]. Moreover, we show that  $\{s_1, \dots, s_n\}$  is *stable* if  $\Delta$  does not contain near-degenerate edges.

We note that the support of each fundamental function  $s_i$  has at most one vertex of the triangulation in its interior, similar to the minimally supported basis functions for bivariate splines of degree  $q \geq 3r + 2$  constructed in [8, 15, 16]. Particularly, the supports of  $s_1, \dots, s_n$  coincide with the supports of the basis splines for  $S_q^{r,\rho}(\Delta)$  given in [16]. However, there are important differences between our basis and the

bases in [8, 15, 16]. Since the Bernstein-Bézier minimal determining sets used in [15, 16] cannot be directly transformed into a Hermite interpolation scheme, it does not seem possible to employ those bases in a proof of optimal approximation order in the presence of near-singularities. The main difference between our interpolation operator and the one presented in [8] is that we do not avoid using normal derivatives to the edges as nodal functionals. Therefore, we assign the nodal functionals to every vertex, edge and triangle in a more symmetric fashion.

## 2 Nodal Functionals

Given a regular triangulation  $\Delta$ , we denote by  $N$  the number of triangles, by  $V$  the number of vertices, by  $V_I$  and  $V_B$  the number of interior and boundary vertices respectively,  $V_I + V_B = V$ , by  $E$  the number of edges, and by  $E_I$  and  $E_B$  the number of interior and boundary edges respectively,  $E_I + E_B = E$ . It is well known that

$$(2.1) \quad \begin{aligned} E_B &= V_B, \\ E_I &= 3V_I + V_B - 3, \\ N &= 2V_I + V_B - 2. \end{aligned}$$

In [16] it was shown that

$$(2.2) \quad \begin{aligned} \dim S_q^{r,\rho}(\Delta) &= \binom{\rho+2}{2}V + \left( \binom{q-3r-1}{2} - 3\binom{2r-\rho+1}{2} \right) N \\ &+ \frac{1}{2}(r+1)(2q-4\rho+r-2)E + \binom{2r-\rho+1}{2}\sigma, \end{aligned}$$

with  $\sigma$  being the number of singular vertices of  $\Delta$ , where a *singular vertex*  $v$  is a vertex which is formed by two lines which cross at  $v$ . It is easy to see that a vertex  $v$  is singular if and only if at least three edges are degenerate at  $v$ , where the degeneracy of an edge is defined as follows.

**Definition 2.1** Suppose  $e_1, e_2, e_3$  are three consecutive edges attached to a vertex  $v$ . The edge  $e_2$  is said to be *degenerate* at  $v$  whenever the edges  $e_1$  and  $e_3$  are collinear. An edge  $e$  attached to  $v$  is said to be *nondegenerate* at  $v$  if it is either a boundary edge or an interior edge which fails to be degenerate.

In the finite element method piecewise polynomial trial functions are usually determined by their values and derivatives at some points, so-called nodal values (see,

e.g., [29, p. 101]). In [19, 11] and [26] this technique was applied to the study of spline spaces  $S_q^1(\Delta)$ ,  $q \geq 5$ , and supersplines  $S_q^{r,\rho}(\Delta)$  with  $\rho \geq 2r$  and  $q \geq 2\rho + 1$ , respectively.

We set

$$C^\mu(\Delta) := \{f \in C(\Omega) : f|_T \in C^\mu(T) \text{ for all } T \in \Delta\}, \quad \mu = 0, 1, \dots,$$

and denote by  $D_\tau$  the derivative operator in the direction of a unit vector  $\tau = (\tau_x, \tau_y)$  in the plane, so that

$$D_\tau f := \tau_x D_x f + \tau_y D_y f, \quad D_x f := \frac{\partial f}{\partial x}, \quad D_y f := \frac{\partial f}{\partial y}.$$

**Definition 2.2** Given  $f \in C^{\alpha+\beta}(\Delta)$ ,  $\alpha, \beta \geq 0$ , any number

$$(2.3) \quad \nu f = D_{\tau_1}^\alpha D_{\tau_2}^\beta (f|_T)(z),$$

where  $T \in \Delta$ ,  $z \in T$ , and  $\tau_1, \tau_2$  are some unit vectors in the plane, is said to be a *nodal value* of  $f$ , and the linear functional  $\nu : C^{\alpha+\beta}(\Delta) \rightarrow \mathbb{R}$  defined by (2.3) is a *nodal functional*, with  $d(\nu) := \alpha + \beta$  being the *degree* of  $\nu$ .

For some special choices of  $z, \tau_1, \tau_2$  it is convenient to use the following simplified notation which goes back to [19]. 1) If  $v$  is a vertex of  $\Delta$  and  $e$  is an edge attached to  $v$ , we set

$$D_e^\alpha f(v) := D_\tau^\alpha (f|_T)(v), \quad \alpha \geq 1,$$

where  $\tau$  is the unit vector in the direction of  $e$  away from  $v$ , and  $T \in \Delta$  is one of the triangles with edge  $e$ . The notation is correct since in the case when there are two different triangles  $T_1, T_2$  attached to  $e$ ,  $f|_{T_1}$  and  $f|_{T_2}$  coincide along  $e$ , and hence

$$D_\tau^\alpha (f|_{T_1})(v) = D_\tau^\alpha (f|_{T_2})(v).$$

2) If  $v$  is a vertex of  $\Delta$  and  $e_1, e_2$  are two consecutive edges attached to  $v$ , we set

$$D_{e_1}^\alpha D_{e_2}^\beta f(v) := D_{\tau_1}^\alpha D_{\tau_2}^\beta (f|_T)(v), \quad \alpha, \beta \geq 1,$$

where  $T \in \Delta$  is the triangle with vertex  $v$  and edges  $e_1, e_2$ , and  $\tau_i$  is the unit vector in the  $e_i$  direction away from  $v$ . 3) For every edge  $e$  of the triangulation  $\Delta$  we choose a unit vector  $\tau^\perp$  (one of two possible) orthogonal to  $e$  and set

$$D_{e^\perp}^\alpha f(z) := D_{\tau^\perp}^\alpha f(z), \quad z \in e, \quad \alpha \geq 1,$$

provided  $f \in C^\alpha(z)$ .

We now associate with the superspline space  $S_q^{r,\rho}(\Delta)$ , with  $q \geq 3r + 2$  and

$$(2.4) \quad \rho = r + \left\lceil \frac{r+1}{2} \right\rceil,$$

a set  $\mathcal{N}$  of nodal functionals, as follows.

For every vertex  $v$  of  $\Delta$ , let  $T_v^1, \dots, T_v^{n(v)}$  be all triangles attached to  $v$  and numbered counterclockwise (starting from a boundary triangle if  $v$  is a boundary vertex). Denote by  $e_i$  the common edge of  $T_v^{i-1}$  and  $T_v^i$ ,  $i = 2, \dots, n(v)$ . If  $v$  is an interior vertex,  $e_1 = e_{n(v)+1}$  denote the common edge of  $T_v^1$  and  $T_v^{n(v)}$ . Otherwise,  $e_1$  and  $e_{n(v)+1}$  are the boundary edges (attached to  $v$ ) of  $T_v^1$  and  $T_v^{n(v)}$  respectively.

We define  $\mathcal{N}(v)$  to be the set of nodal functionals assigning to every function  $f \in C^\rho(v) \cap C^{2r}(\Delta)$  the following nodal values:

(v1)  $D_x^\alpha D_y^\beta f(v)$  for all  $(\alpha, \beta) \in A_1$ , where

$$A_1 := \{(\alpha, \beta) \in \mathbb{Z}^2 : \alpha \geq 0, \beta \geq 0, \alpha + \beta \leq \rho\},$$

(v2)  $D_{e_i}^\alpha D_{e_{i+1}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$ , where

$$A_2 := \{(\alpha, \beta) \in \mathbb{Z}^2 : \alpha \leq r, \beta \leq r, \alpha + \beta \geq \rho + 1\},$$

and for each  $i \in \{1, \dots, n(v)\}$  such that  $e_i$  is nondegenerate at  $v$ ,


(v3)  $D_{e_i}^\alpha D_{e_{i+1}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_3$ , where

$$A_3 := \{(\alpha, \beta) \in \mathbb{Z}^2 : \alpha \geq r + 1, 2\alpha + \beta \leq 3r + 1, \alpha + \beta \geq \rho + 1\},$$

and for each  $i \in \{1, \dots, n(v)\}$  such that  $e_i$  is degenerate at  $v$ ,

(v4)  $D_{e_1}^\alpha D_{e_2}^\beta f(v)$  and  $D_{e_{n(v)+1}}^\alpha D_{e_{n(v)}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_3$  if  $v$  is a boundary vertex, and

(v5)  $D_{e_1}^\alpha D_{e_2}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$  if  $v$  is a singular vertex.



**Fig. 2.1.** The sets  $A_1$ ,  $A_2$  and  $A_3$ .

On every edge  $e$  of  $\Delta$ , with vertices  $v_1$  and  $v_2$ , we take points

$$(2.5) \quad z_e^{\mu, i} := v_1 + \frac{i}{\kappa_\mu + 1}(v_2 - v_1), \quad i = 1, \dots, \kappa_\mu, \quad \mu = 0, \dots, r,$$

where

$$(2.6) \quad \kappa_\mu := q - 3r - 1 - (r - \mu) \bmod 2 = q - 2r - 1 - \mu - 2 \left\lfloor \frac{r+1-\mu}{2} \right\rfloor,$$

and define  $\mathcal{N}(e)$  to be the set of nodal functionals assigning to every function  $f \in C^r(\Omega)$  the following nodal values:

$$(e) \quad D_{e^\perp}^\mu f(z_e^{\mu, 1}), \dots, D_{e^\perp}^\mu f(z_e^{\mu, \kappa_\mu}) \text{ for all } \mu = 0, \dots, r.$$

In every triangle  $T \in \Delta$ , with vertices  $v_1$ ,  $v_2$  and  $v_3$ , we take uniformly spaced points

$$(2.7) \quad z_T^{i, j, k} := (iv_1 + jv_2 + kv_3)/q, \quad i + j + k = q,$$

and define  $\mathcal{N}(T)$  to be the set of nodal functionals assigning to every function  $f \in C(\Omega)$  the following nodal values:

$$(t) \quad f(z_T^{i, j, k}) \text{ for all } i, j, k \text{ such that } i + j + k = q \text{ and } r < i, j, k < q - 2r.$$

We set

$$\mathcal{N} := \bigcup_v \mathcal{N}(v) \cup \bigcup_e \mathcal{N}(e) \cup \bigcup_T \mathcal{N}(T).$$

**Lemma 2.3** *We have*

$$(2.8) \quad \text{card } \mathcal{N} = \dim S_q^{r,\rho}(\Delta).$$

**Proof.** It is easy to see that

$$(2.9) \quad \begin{aligned} \text{card } A_1 &= \binom{\rho+2}{2}, \quad \text{card } A_2 = \text{card } A_3 = \binom{2r-\rho+1}{2}, \\ \text{card } \mathcal{N}(e) &= (r+1)(q-3r-1) - \left\lfloor \frac{r+1}{2} \right\rfloor, \quad \text{card } \mathcal{N}(T) = \binom{q-3r-1}{2}. \end{aligned}$$

Therefore,

$$\begin{aligned} \text{card } \mathcal{N} &= \binom{\rho+2}{2}V + \left( \binom{q-3r-1}{2} + 3\binom{2r-\rho+1}{2} \right)N + 2\binom{2r-\rho+1}{2}V_B \\ &+ \left( (r+1)(q-3r-1) + r - \rho \right)E + \binom{2r-\rho+1}{2}\sigma. \end{aligned}$$

The lemma now follows from (2.1), (2.2) and a simple computation. ■

### 3 Hermite Type Interpolation

**Theorem 3.1** *Let  $r \geq 1$ ,  $q \geq 3r + 2$  and  $\rho = r + \left\lfloor \frac{r+1}{2} \right\rfloor$ . Given  $f \in C^{2r}(\Omega)$ , there exists a unique spline  $s_f \in S_q^{r,\rho}(\Delta)$  satisfying the following Hermite type interpolation conditions*

$$(3.1) \quad \nu s_f = \nu f \quad \text{for all } \nu \in \mathcal{N},$$

where  $\mathcal{N}$  is defined above. Moreover, if  $f \in C^m(\Omega)$  ( $m \in \{2r, \dots, q+1\}$ ) and  $T \in \Delta$ , then

$$(3.2) \quad \|D_x^\alpha D_y^\beta (f - s_f)\|_{L_\infty(T)} \leq K h_T^{m-\alpha-\beta} \max_{0 \leq \mu \leq m} \|D_x^\mu D_y^{m-\mu} f\|_{C(T)},$$

for all  $\alpha, \beta \geq 0$ ,  $\alpha + \beta \leq m$ , where  $h_T$  is the diameter of  $T$ , and  $K$  is a constant which depends only on  $r, q$  and the smallest angle  $\theta_\Delta$  in  $\Delta$ .


We will prove Theorem 3.1 at the end of this section, after establishing several lemmas.

In the first two lemmas we consider a simple triangulation consisting of two triangles and establish some relations between nodal values of two polynomials defined on each triangle and joined together with  $C^r$  smoothness across a common edge of the triangles.

Let  $T_1$  and  $T_2$  be two triangles sharing a common edge  $e = [v_1, v_2]$ , and let  $e_i \neq e$  be the other edge of  $T_i$  with endpoint  $v_1$ ,  $i = 1, 2$ . Denote by  $\tau, \tau_1, \tau_2$  the unit



vectors applied at  $v_1$  in the direction of edges  $e$ ,  $e_1$ ,  $e_2$  respectively, and by  $\theta_i$  the angle between  $\tau$  and  $\tau_i$ ,  $i = 1, 2$ . (See Fig. 3.1.)



**Fig. 3.1.**

Furthermore, let  $s$  be a piecewise polynomial function on  $T_1 \cup T_2$  such that

$$s|_{T_i} = p_i \in \Pi_q, \quad i = 1, 2.$$

Our first lemma characterizes  $C^r$  smoothness of  $s$  across  $e$  in terms of its nodal values. (See Lemma 2 in [8] for a similar characterization using barycentric coordinates.)

**Lemma 3.2** *Let  $r \leq q$ .*

1) *If  $\theta_1 + \theta_2 \neq \pi$ , then  $s \in C^r(T_1 \cup T_2)$  if and only if*

$$(3.3) \quad \sin^\alpha \theta_1 D_{\tau_2}^\alpha D_\tau^{\gamma-\alpha} p_2(v_1) = \sum_{\beta=0}^{\alpha} (-1)^\beta \binom{\alpha}{\beta} \sin^{\alpha-\beta}(\theta_1 + \theta_2) \sin^\beta \theta_2 D_{\tau_1}^\beta D_\tau^{\gamma-\beta} p_1(v_1),$$

for all  $\alpha = 0, \dots, r$  and  $\gamma = \alpha, \dots, q$ .

2) *If  $\theta_1 + \theta_2 = \pi$ , then  $s \in C^r(T_1 \cup T_2)$  if and only if*

$$(3.4) \quad D_{\tau_2}^\alpha D_\tau^{\gamma-\alpha} p_2(v_1) = (-1)^\alpha D_{\tau_1}^\alpha D_\tau^{\gamma-\alpha} p_1(v_1), \quad \alpha = 0, \dots, r, \quad \gamma = \alpha, \dots, q.$$

**Proof.** Evidently,  $s \in C^r(T_1 \cup T_2)$  if and only if for some unit vector  $\tau'$  noncollinear with  $\tau$ ,

$$D_\tau^\mu D_{\tau'}^\alpha p_2(z) = D_\tau^\mu D_{\tau'}^\alpha p_1(z), \quad \text{for all } \alpha, \mu \geq 0, \alpha + \mu \leq r, \text{ and all } z \in e.$$

Since  $(D_{\tau'}^\alpha p_i)|_e$ ,  $i = 1, 2$ , is a univariate polynomial of degree at most  $q - \alpha$ , this is equivalent to the condition

$$D_\tau^{\gamma-\alpha} D_{\tau'}^\alpha p_2(v_1) = D_\tau^{\gamma-\alpha} D_{\tau'}^\alpha p_1(v_1), \quad \alpha = 0, \dots, r, \quad \gamma = \alpha, \dots, q.$$

We now choose  $\tau' = \tau_2$ . If  $\theta_1 + \theta_2 = \pi$ , then  $\tau_1 = -\tau_2$ , and we immediately get (3.4). Otherwise, if  $\theta_1 + \theta_2 \neq \pi$ , then the vectors  $\tau$ ,  $\tau_1$  and  $\tau_2$  stay in the relation

$$\tau \sin(\theta_1 + \theta_2) = \tau_1 \sin \theta_2 + \tau_2 \sin \theta_1,$$

which implies

$$\sin^\alpha \theta_1 D_{\tau_2}^\alpha D_\tau^{\gamma-\alpha} p_1(v_1) = \sum_{\beta=0}^{\alpha} (-1)^\beta \binom{\alpha}{\beta} \sin^{\alpha-\beta}(\theta_1 + \theta_2) \sin^\beta \theta_2 D_{\tau_1}^\beta D_\tau^{\gamma-\beta} p_1(v_1),$$

and the first statement of the lemma follows. ■

Thus, the nodal values of  $s \in C^r(T_1 \cup T_2)$  stay in relations (3.3). The same relations hold for every sufficiently smooth function  $f$ . By solving a linear system we can estimate some of the nodal values of  $f - s$  at  $v_1$  involved in (3.3) in terms of the others.

**Lemma 3.3** *Suppose that  $s$ , as defined above, is in  $C^r(T_1 \cup T_2)$ , and let  $f \in C^k(T_1 \cup T_2)$  for some  $k \in \{\rho + 1, \dots, 2r\}$ . If  $\theta_1 + \theta_2 \neq \pi$ , then for every  $\beta = 2k - 3r - 1, \dots, k - r - 1$ ,*

$$(3.5) \quad \begin{aligned} |D_{\tau_1}^\beta D_\tau^{k-\beta}(f - p_1)(v_1)| &\leq K \left( \max_{0 \leq \alpha \leq 2k-3r-2} |D_{\tau_1}^\alpha D_\tau^{k-\alpha}(f - p_1)(v_1)| \right. \\ &\quad \left. + |\sin^{-r}(\theta_1 + \theta_2)| \max_{\substack{k-r \leq \alpha \leq r \\ i=1,2}} |D_{\tau_i}^\alpha D_\tau^{k-\alpha}(f - p_i)(v_1)| \right), \end{aligned}$$

where  $K$  depends only on  $r$  and  $\theta_2$ .

**Proof.** Since  $f \in C^k(v_1)$ , we have

$$\sin^\alpha \theta_1 D_{\tau_2}^\alpha D_\tau^{k-\alpha} f(v_1) = \sum_{\beta=0}^{\alpha} (-1)^\beta \binom{\alpha}{\beta} \sin^{\alpha-\beta}(\theta_1 + \theta_2) \sin^\beta \theta_2 D_{\tau_1}^\beta D_\tau^{k-\beta} f(v_1)$$

for all  $\alpha = 0, \dots, k$ . This, together with (3.3), imply that

$$(3.6) \quad a_{2,\alpha} = \sum_{\beta=0}^{\alpha} (-1)^{\beta} \binom{\alpha}{\beta} a_{1,\beta}, \quad \alpha = 2k - 3r - 1, \dots, r,$$

where

$$\begin{aligned} a_{1,\beta} &:= \sin^{-\beta}(\theta_1 + \theta_2) \sin^{\beta} \theta_2 D_{\tau_1}^{\beta} D_{\tau}^{k-\beta}(f - p_1)(v_1), \\ a_{2,\beta} &:= \sin^{-\beta}(\theta_1 + \theta_2) \sin^{\beta} \theta_1 D_{\tau_2}^{\beta} D_{\tau}^{k-\beta}(f - p_2)(v_1). \end{aligned}$$

Consider (3.6) as a system

$$Ax = b$$

of  $4r - 2k + 2$  linear equations in  $4r - 2k + 2$  unknowns

$$a_{i,\beta}, \quad \beta = 2k - 3r - 1, \dots, k - r - 1, \quad i = 1, 2.$$

Thus, we have

$$x = (a_{1,2k-3r-1}, \dots, a_{1,k-r-1}, a_{2,2k-3r-1}, \dots, a_{2,k-r-1})^t,$$

$b$  is a  $(4r - 2k + 2)$ -vector whose components are some linear combinations of

$$\begin{aligned} a_{1,\beta}, \quad \beta = 0, \dots, 2k - 3r - 2, \quad \text{and} \\ a_{i,\beta}, \quad \beta = k - r, \dots, r, \quad i = 1, 2, \end{aligned}$$

and

$$A = \begin{pmatrix} B & -I \\ C & O \end{pmatrix},$$

where

$$C = \left( (-1)^{r+j} \binom{n+i}{n-m+j} \right)_{i,j=1}^m, \quad \text{with } n := k - r - 1, \quad m := 2r - k + 1,$$

$I$  is an  $m \times m$  identity matrix,  $O$  is an  $m \times m$  zero matrix, and  $B$  is a certain  $m \times m$  matrix. Since the determinant of  $C$  is a nonzero constant multiple of

$$\det \left( \frac{1}{(m+i-j)!} \right)_{i,j=1}^m \neq 0,$$

$A$  is nonsingular. Therefore,

$$\|x\|_\infty \leq \|A^{-1}\|_\infty \|b\|_\infty,$$

where  $\|A^{-1}\|_\infty$  is bounded by a constant dependent only on  $r$ . Particularly, for all  $\beta = 2k - 3r - 1, \dots, k - r - 1$ ,

$$|a_{1,\beta}| \leq K_1 \left( \max_{0 \leq \alpha \leq 2k-3r-2} |a_{1,\alpha}| + \max_{k-r \leq \alpha \leq r} |a_{1,\alpha}| + \max_{k-r \leq \alpha \leq r} |a_{2,\alpha}| \right),$$

where  $K_1$  depends only on  $r$ .

Recalling the definition of  $a_{i,\beta}$ ,  $i = 1, 2$ , we obtain

$$\begin{aligned} & |D_{\tau_1}^\beta D_\tau^{k-\beta}(f - p_1)(v_1)| = |a_{1,\beta} \sin^\beta(\theta_1 + \theta_2) \sin^{-\beta}\theta_2| \\ & \leq K_1 \left( \max_{0 \leq \alpha \leq 2k-3r-2} |\sin^{\beta-\alpha}(\theta_1 + \theta_2) \sin^{\alpha-\beta}\theta_2| |D_{\tau_1}^\alpha D_\tau^{k-\alpha}(f - p_1)(v_1)| \right. \\ & \quad + \max_{k-r \leq \alpha \leq r} |\sin^{\beta-\alpha}(\theta_1 + \theta_2) \sin^{\alpha-\beta}\theta_2| |D_{\tau_1}^\alpha D_\tau^{k-\alpha}(f - p_1)(v_1)| \\ & \quad \left. + \max_{k-r \leq \alpha \leq r} |\sin^{\beta-\alpha}(\theta_1 + \theta_2) \sin^\alpha\theta_1 \sin^{-\beta}\theta_2| |D_{\tau_2}^\alpha D_\tau^{k-\alpha}(f - p_2)(v_1)| \right), \end{aligned}$$

and (3.5) follows.  $\blacksquare$

We also need the following univariate ‘‘weak interpolation’’ lemma (compare [21, Remark 5ii] and [12, Lemma 4]).

**Lemma 3.4** *Let  $e \subset \mathbb{R}^2$  be an interval with endpoints  $v_1, v_2$ , and let  $\mu \in \{0, \dots, r\}$  and  $m \in \{r + \lceil \frac{r+1-\mu}{2} \rceil, \dots, q + 1 - \mu\}$ . Then for any  $f \in C^m(e)$ , any  $p \in \Pi_{q-\mu}$  and every  $\gamma = 0, \dots, m$ ,*

$$(3.7) \quad \begin{aligned} \|D_\tau^\gamma(f - p)\|_{C(e)} & \leq Kh^{-\gamma} \left( h^m \|D_\tau^m f\|_{C(e)} + \max_{1 \leq i \leq \kappa_\mu} |(f - p)(z_e^{\mu,i})| \right. \\ & \quad \left. + \max_{\substack{0 \leq \alpha \leq r + \lceil \frac{r+1-\mu}{2} \rceil \\ i=1,2}} h^\alpha |D_\tau^\alpha(f - p)(v_i)| \right), \end{aligned}$$

where  $h$  is the length of  $e$ ,  $\tau$  denotes the unit vector in the direction of  $e$ ,  $z_e^{\mu,i}$  and  $\kappa_\mu$  are defined in (2.5) and (2.6), respectively, and  $K$  is a constant which depends only on  $q$ .

**Proof.** It is sufficient to consider the case  $e = [0, h]$ , i.e.,  $v_1 = (0, 0)$ ,  $v_2 = (0, h)$ . Then  $\tau = (1, 0)$ ,  $D_\tau = D_x$ ,  $z_e^{\mu, i} = (\frac{ih}{\kappa_\mu + 1}, 0)$ ,  $i = 1, \dots, \kappa_\mu$ .

Since  $f \in C^m[0, h]$ , we have

$$(3.8) \quad \|D_x^\gamma(f - \tilde{p})\|_{C[0, h]} \leq \frac{h^{m-\gamma}}{(m-\gamma)!} \|D_x^m f\|_{C[0, h]}, \quad \gamma = 0, \dots, m,$$

where  $\tilde{p}$  is the (univariate) Taylor polynomial,

$$\tilde{p}(x) := \sum_{\nu=0}^{m-1} \frac{D_x^\nu f(0)}{\nu!} x^\nu.$$

Therefore,

$$\begin{aligned} \|D_x^\gamma(f - p)\|_{C[0, h]} &\leq \|D_x^\gamma(f - \tilde{p})\|_{C[0, h]} + \|D_x^\gamma(\tilde{p} - p)\|_{C[0, h]} \\ &\leq h^{m-\gamma} \|D_x^m f\|_{C[0, h]} + \|D_x^\gamma(\tilde{p} - p)\|_{C[0, h]}, \end{aligned}$$

and we only need to estimate  $\|D_x^\gamma(\tilde{p} - p)\|_{C[0, h]}$ .

Let

$$\lambda_\mu := r + \left[ \frac{r+1-\mu}{2} \right].$$

Since  $\kappa_\mu + 2(\lambda_\mu + 1) = q - \mu + 1$ , the following Hermite interpolation problem

$$g(z_e^{\mu, i}) = a_i, \quad i = 1, \dots, \kappa_\mu, \quad D_x^\alpha g(v_j) = a_{j, \alpha}, \quad \alpha = 0, \dots, \lambda_\mu, \quad j = 1, 2,$$

has a unique solution  $g$  among univariate polynomials of degree at most  $q - \mu$ , for any given data  $a_i$ ,  $i = 1, \dots, \kappa_\mu$ , and  $a_{j, \alpha}$ ,  $\alpha = 0, \dots, \lambda_\mu$ ,  $j = 1, 2$ . Then

$$(\tilde{p} - p)(t) = \sum_{i=1}^{\kappa_\mu} (\tilde{p} - p)(z_e^{\mu, i}) L_{i, h}(t) + \sum_{j=1, 2} \sum_{\alpha=0}^{\lambda_\mu} D_x^\alpha (\tilde{p} - p)(v_j) L_{j, \alpha, h}(t), \quad t \in [0, h],$$

where  $L_{i, h}$ ,  $i = 1, \dots, \kappa_\mu$ , and  $L_{j, \alpha, h}$ ,  $\alpha = 0, \dots, \lambda_\mu$ ,  $j = 1, 2$ , denote the fundamental polynomials of the above interpolation problem, i.e., they are univariate polynomials of degree at most  $q - \mu$ , uniquely determined by the conditions

$$\begin{aligned} L_{i, h}\left(\frac{jh}{\kappa_\mu + 1}\right) &= \delta_{i, j}, \quad i, j = 1, \dots, \kappa_\mu, \\ D_x^\alpha L_{i, h}(0) &= D_x^\alpha L_{i, h}(h) = 0, \quad \alpha = 0, \dots, \lambda_\mu, \quad i = 1, \dots, \kappa_\mu, \end{aligned}$$

and

$$\begin{aligned} L_{i,\alpha,h}\left(\frac{jh}{\kappa_\mu+1}\right) &= 0, \quad j = 1, \dots, \kappa_\mu, \quad \alpha = 0, \dots, \lambda_\mu, \quad i = 1, 2, \\ D_x^\nu L_{1,\alpha,h}(0) &= D_x^\nu L_{2,\alpha,h}(h) = \delta_{\alpha,\nu}, \quad \alpha, \nu = 0, \dots, \lambda_\mu, \\ D_x^\nu L_{1,\alpha,h}(h) &= D_x^\nu L_{2,\alpha,h}(0) = 0, \quad \alpha, \nu = 0, \dots, \lambda_\mu, \end{aligned}$$

respectively. By a uniqueness argument, it is easy to check that

$$\begin{aligned} D_x^\gamma L_{i,h}(t) &= h^{-\gamma} D_x^\gamma L_{i,1}\left(\frac{t}{h}\right), \quad t \in [0, h], \\ D_x^\gamma L_{j,\alpha,h}(t) &= h^{\alpha-\gamma} D_x^\gamma L_{j,\alpha,1}\left(\frac{t}{h}\right), \quad t \in [0, h], \end{aligned}$$

and, consequently,

$$\begin{aligned} \|D_x^\gamma L_{i,h}\|_{C[0,h]} &= h^{-\gamma} \|D_x^\gamma L_{i,1}\|_{C[0,1]}, \\ \|D_x^\gamma L_{j,\alpha,h}\|_{C[0,h]} &= h^{\alpha-\gamma} \|D_x^\gamma L_{j,\alpha,1}\|_{C[0,1]}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \|D_x^\gamma(\tilde{p} - p)\|_{C[0,h]} &\leq \sum_{i=1}^{\kappa_\mu} |(\tilde{p} - p)(z_e^{\mu,i})| h^{-\gamma} \|L_{i,1}\|_{C[0,1]} \\ &\quad + \sum_{j=1,2} \sum_{\alpha=0}^{\lambda_\mu} |D_x^\alpha(\tilde{p} - p)(v_j)| h^{\alpha-\gamma} \|L_{j,\alpha,1}\|_{C[0,1]}, \end{aligned}$$


Since  $\tilde{p} - p = (\tilde{p} - f) + (f - p)$ , (3.8) implies

$$\begin{aligned} |(\tilde{p} - p)(z_e^{\mu,i})| &\leq h^m \|D_x^m f\|_{C[0,h]} + |(f - p)(z_e^{\mu,i})|, \\ |D_x^\alpha(\tilde{p} - p)(v_j)| &\leq h^{m-\alpha} \|D_x^m f\|_{C[0,h]} + |D_x^\alpha(f - p)(v_j)|, \end{aligned}$$

and the lemma follows because  $\|L_{i,1}\|_{C[0,1]}$  and  $\|L_{j,\alpha,1}\|_{C[0,1]}$  are bounded by a constant dependent only on  $q$ . ■

Since our interpolation scheme is based on nodal values involving partial derivatives in various directions, we need a tool to recast the (weak) interpolation conditions in such a form that their interaction becomes tractable. As a “common unit” we will use derivatives of the type  $D_e^\gamma D_{e^\perp}^\mu(f - s)$ . The next two lemmas provide estimations of these derivatives in terms of nodal values of our scheme.

Consider first a single triangle  $T_1 \in \Delta$ , and let  $e$  be one of its edges, with vertices  $v_1$  and  $v_2$ . (Note that  $e$  may be a boundary edge of  $\Delta$ .) Denote by  $e_{1,1}$  and  $e_{1,2}$  two other edges of  $T_1$ , attached to  $v_1$  and  $v_2$ , respectively, and by  $\theta_{1,i}$  the angle between  $e$  and  $e_{1,i}$ ,  $i = 1, 2$ . (See Fig. 3.2.)



**Fig. 3.2.**

**Lemma 3.5** *Let  $s \in S_q^{r,\rho}(\Delta)$  and  $f \in C^m(\Omega)$  ( $m \in \{2r, \dots, q+1\}$ ) be given. Then for all  $\mu = 0, \dots, r$  and  $\gamma = 0, \dots, m - \mu$ ,*

$$\begin{aligned}
(3.9) \quad \|D_e^\gamma D_{e^\perp}^\mu (f - s)\|_{C(e)} &\leq Kh^{-\gamma-\mu} \left( h^m \max_{0 \leq \mu' \leq \mu} \|D_e^{m-\mu'} D_{e^\perp}^{\mu'} f\|_{C(e)} \right. \\
&\quad + \max_{\substack{0 \leq \mu' \leq \mu \\ 0 \leq i \leq \kappa_{\mu'}}} h^{\mu'} |D_{e^\perp}^{\mu'} (f - s)(z_e^{\mu', i})| \\
&\quad \left. + \max_{\substack{(\alpha, \beta) \in A_1 \cup A_2 \cup A_3 \\ \beta \leq \mu, i=1,2}} h^{\alpha+\beta} |D_e^\alpha D_{e_{1,i}}^\beta (f - s)(v_i)| \right),
\end{aligned}$$

where  $h$  is the length of  $e$ , the sets  $A_1$ – $A_3$  are defined above, and  $K$  depends only on  $q$  and  $\min\{\theta_{1,1}, \theta_{1,2}\}$ .

**Proof.** Since

$$(\alpha, \mu) \in A_1 \cup A_2 \cup A_3 \iff 0 \leq \alpha \leq r + \left\lceil \frac{r+1-\mu}{2} \right\rceil, \quad \mu = 0, \dots, r,$$

Lemma 3.4 shows that there exists a constant  $K_1$  dependent only on  $q$ , such that

$$\begin{aligned}
\|D_e^\gamma (f - s)\|_{C(e)} &\leq K_1 h^{-\gamma} \left( h^m \|D_e^m f\|_{C(e)} + \max_{0 \leq i \leq \kappa_\mu} |(f - s)(z_e^{0,i})| \right. \\
&\quad \left. + \max_{\substack{(\alpha, 0) \in A_1 \cup A_2 \cup A_3 \\ i=1,2}} h^\alpha |D_e^\alpha (f - s)(v_i)| \right), \quad \gamma = 0, \dots, m,
\end{aligned}$$

which proves (3.9) for  $\mu = 0$ . Proceeding by induction on  $\mu$ , we suppose that (3.9) holds for  $0, \dots, \mu-1$ . Again by Lemma 3.4, applied to  $D_{e^\perp}^\mu f \in C^{m-\mu}(e)$  and  $p = D_{e^\perp}^\mu s$ , we get for all  $\gamma = 0, \dots, m - \mu$ ,

$$\begin{aligned} \|D_e^\gamma D_{e^\perp}^\mu (f - s)\|_{C(e)} &\leq K_1 h^{-\gamma} \left( h^{m-\mu} \|D_e^{m-\mu} D_{e^\perp}^\mu f\|_{C(e)} + \max_{0 \leq i \leq \kappa_\mu} |D_{e^\perp}^\mu (f - s)(z_e^{\mu, i})| \right. \\ &\quad \left. + \max_{\substack{(\alpha, \mu) \in A_1 \cup A_2 \cup A_3 \\ i=1, 2}} h^\alpha |D_e^\alpha D_{e^\perp}^\mu (f - s)(v_i)| \right). \end{aligned}$$

Thus, we need to estimate  $D_e^\alpha D_{e^\perp}^\mu (f - s)(v_i)$  in terms of  $D_e^\alpha D_{e_{1,i}}^\beta (f - s)(v_i)$  with  $\beta \leq \mu$ . To this end, we use the relation

$$\tau_{1,i} = \pm \tau \cos \theta_{1,i} \pm \tau^\perp \sin \theta_{1,i}, \quad i = 1, 2,$$

where  $\tau_{1,i}$ ,  $\tau$  and  $\tau^\perp$  are the unit vectors in the directions of  $e_{1,i}$ ,  $e$  and  $e^\perp$  respectively, so that

$$(3.10) \quad D_e^\alpha D_{e_{1,i}}^\mu (f - s)(v_i) = \sum_{\mu'=0}^{\mu} \pm \binom{\mu}{\mu'} \cos^{\mu-\mu'} \theta_{1,i} \sin^{\mu'} \theta_{1,i} D_e^{\alpha+\mu-\mu'} D_{e^\perp}^{\mu'} (f - s)(v_i),$$

and hence,

$$\begin{aligned} |D_e^\alpha D_{e^\perp}^\mu (f - s)(v_i)| &\leq |D_e^\alpha D_{e_{1,i}}^\mu (f - s)(v_i)| \\ &\quad + K_2 \max_{0 \leq \mu' \leq \mu-1} |D_e^{\alpha+\mu-\mu'} D_{e^\perp}^{\mu'} (f - s)(v_i)|, \quad i = 1, 2, \end{aligned}$$


where  $K_2$  depends only on  $\mu$  and  $\min\{\theta_{1,1}, \theta_{1,2}\}$ . Furthermore, by the induction hypothesis,

$$\begin{aligned} |D_e^{\alpha+\mu-\mu'} D_{e^\perp}^{\mu'} (f - s)(v_i)| &\leq K h^{-\alpha-\mu} \left( h^m \max_{0 \leq \mu'' \leq \mu'} \|D_e^{m-\mu''} D_{e^\perp}^{\mu''} f\|_{C(e)} \right. \\ &\quad + \max_{\substack{0 \leq \mu'' \leq \mu' \\ 0 \leq i \leq \kappa_{\mu''}}} h^{\mu''} |D_{e^\perp}^{\mu''} (f - s)(z_e^{\mu'', i})| \\ &\quad \left. + \max_{\substack{(\alpha', \beta) \in A_1 \cup A_2 \cup A_3 \\ \beta \leq \mu', i=1, 2}} h^{\alpha'+\beta} |D_e^{\alpha'} D_{e_{1,i}}^\beta (f - s)(v_i)| \right), \end{aligned}$$

and (3.9) follows.  $\blacksquare$



Under the notations of Lemma 3.5, suppose that  $e$  is an interior edge of  $\Delta$  and denote by  $T_2$  the triangle in  $\Delta$  that share  $e$  with  $T_1$ . Let  $e_{2,1}$  and  $e_{2,2}$  be two other edges of  $T_2$ , attached to  $v_1$  and  $v_2$ , respectively, and let  $\theta_{2,i}$  be the angle between  $e$  and  $e_{2,i}$ ,  $i = 1, 2$ . (See Fig. 3.3.)



**Fig. 3.3.**

Furthermore, let  $h$  denote the length of  $e$ .

**Lemma 3.6** *Let  $s \in S_q^{r,\rho}(\Delta)$  and  $f \in C^m(\Omega)$  ( $m \in \{2r, \dots, q+1\}$ ) be given.*

1) *If  $\theta_{1,2} + \theta_{2,2} \neq \pi$ , then for all  $\mu = 0, \dots, r$  and  $\gamma = 0, \dots, m - \mu$ ,*

$$\begin{aligned}
(3.11) \quad & \|D_e^\gamma D_{e^\perp}^\mu (f - s)\|_{C(e)} \leq K h^{-\gamma-\mu} \left( h^m \max_{0 \leq \mu' \leq \mu} \|D_e^{m-\mu'} D_{e^\perp}^{\mu'} f\|_{C(e)} \right. \\
& + \max_{\substack{0 \leq \mu' \leq \mu \\ 0 \leq i \leq \kappa_{\mu'}}} h^{\mu'} |D_{e^\perp}^{\mu'} (f - s)(z_e^{\mu',i})| + \max_{\substack{(\alpha,\beta) \in A_1 \cup A_2 \cup A_3 \\ \beta \leq \mu}} h^{\alpha+\beta} |D_e^\alpha D_{e_{1,1}}^\beta (f - s)(v_1)| \\
& + \max_{(\alpha,\beta) \in A_1, \beta \leq \mu} h^{\alpha+\beta} |D_e^\alpha D_{e_{1,2}}^\beta (f - s)(v_2)| \\
& \left. + |\sin^{-r}(\theta_{1,2} + \theta_{2,2})| \max_{(\alpha,\beta) \in A_2, i=1,2} h^{\alpha+\beta} |D_e^\alpha D_{e_{i,2}}^\beta (f - s)(v_2)| \right),
\end{aligned}$$

where  $K$  depends only on  $q$  and  $\min\{\theta_{1,1}, \theta_{1,2}, \theta_{2,2}\}$ .

2) *If both  $\theta_{1,1} + \theta_{2,1} \neq \pi$  and  $\theta_{1,2} + \theta_{2,2} \neq \pi$ , then for all  $\mu = 0, \dots, r$  and  $\gamma = 0, \dots, m - \mu$ ,*

$$\begin{aligned}
(3.12) \quad & \|D_e^\gamma D_{e^\perp}^\mu (f - s)\|_{C(e)} \leq Kh^{-\gamma-\mu} \left( h^m \max_{0 \leq \mu' \leq \mu} \|D_e^{m-\mu'} D_{e^\perp}^{\mu'} f\|_{C(e)} \right. \\
& + \max_{\substack{0 \leq \mu' \leq \mu \\ 0 \leq i \leq \kappa_{\mu'}}} h^{\mu'} |D_{e^\perp}^{\mu'} (f - s)(z_e^{\mu', i})| + \max_{\substack{(\alpha, \beta) \in A_1, \beta \leq \mu \\ j=1,2}} h^{\alpha+\beta} |D_e^\alpha D_{e_{1,j}}^\beta (f - s)(v_j)| \\
& \left. + \max_{j=1,2} |\sin^{-r}(\theta_{1,j} + \theta_{2,j})| \max_{(\alpha, \beta) \in A_2, i=1,2} h^{\alpha+\beta} |D_e^\alpha D_{e_{i,j}}^\beta (f - s)(v_j)| \right),
\end{aligned}$$

where  $K$  depends only on  $q$  and  $\min\{\theta_{1,1}, \theta_{1,2}, \theta_{2,1}, \theta_{2,2}\}$ .

**Proof.** 1) The essential difference between (3.11) and the already established inequality (3.9) is that the terms

$$h^{\alpha+\beta} |D_e^\alpha D_{e_{1,2}}^\beta (f - s)(v_2)|, \quad (\alpha, \beta) \in A_3, \quad \beta \leq \mu,$$

in the right hand side of (3.9) are substituted by

$$|\sin^{-r}(\theta_{1,2} + \theta_{2,2})| h^{\alpha+\beta} |D_e^\alpha D_{e_{i,2}}^\beta (f - s)(v_2)|, \quad (\alpha, \beta) \in A_2, \quad i = 1, 2.$$

If  $\mu = 0$ , then  $\{(\alpha, \beta) \in A_3 : \beta \leq \mu\} = \emptyset$ , and (3.11) is a straightforward consequence of (3.9). Moreover, in order to perform induction on  $\mu$ , we only need an estimation of the form

$$\begin{aligned}
(3.13) \quad & \max_{\substack{(\alpha, \beta) \in A_3 \\ \beta \leq \mu}} h^{\alpha+\beta} |D_e^\alpha D_{e_{1,2}}^\beta (f - s)(v_2)| \leq K_1 \left( \max_{\substack{0 \leq \mu' \leq \mu-1 \\ 0 \leq \gamma \leq m-\mu'}} h^{\gamma+\mu'} \|D_e^\gamma D_{e^\perp}^{\mu'} (f - s)\|_{C(e)} \right. \\
& \left. + |\sin^{-r}(\theta_{1,2} + \theta_{2,2})| \max_{(\alpha, \beta) \in A_2, i=1,2} h^{\alpha+\beta} |D_e^\alpha D_{e_{i,2}}^\beta (f - s)(v_2)| \right).
\end{aligned}$$

To this end we employ Lemma 3.3, which gives for all  $(\alpha, \beta) \in A_3$ , with  $\beta \leq \mu$ ,

$$\begin{aligned}
& |D_e^\alpha D_{e_{1,2}}^\beta (f - s)(v_2)| \leq K_2 \left( \max_{0 \leq \beta' \leq 2(\alpha+\beta)-3r-2} |D_e^{\alpha+\beta-\beta'} D_{e_{1,2}}^{\beta'} (f - s)(v_2)| \right. \\
& \left. + |\sin^{-r}(\theta_{1,2} + \theta_{2,2})| \max_{\alpha+\beta-r \leq \beta' \leq r, i=1,2} |D_e^{\alpha+\beta-\beta'} D_{e_{i,2}}^{\beta'} (f - s)(v_2)| \right).
\end{aligned}$$

Since

$$\beta' \leq 2(\alpha + \beta) - 3r - 2 \implies \beta' \leq \mu - 1,$$

we obtain, by making use of (3.10),

$$\begin{aligned} & \max_{0 \leq \beta' \leq 2(\alpha + \beta) - 3r - 2} |D_e^{\alpha + \beta - \beta'} D_{e_{1,2}}^{\beta'} (f - s)(v_2)| \leq \max_{0 \leq \mu' \leq \mu - 1} |D_e^{\alpha + \beta - \mu'} D_{e_{1,2}}^{\mu'} (f - s)(v_2)| \\ & \leq K_3 \max_{0 \leq \mu' \leq \mu - 1} |D_e^{\alpha + \beta - \mu'} D_{e_\perp}^{\mu'} (f - s)(v_2)| \leq K_3 \max_{0 \leq \mu' \leq \mu - 1} \|D_e^{\alpha + \beta - \mu'} D_{e_\perp}^{\mu'} (f - s)\|_{C(e)}. \end{aligned}$$

Furthermore, since

$$\alpha + \beta - r \leq \beta' \leq r \implies (\alpha + \beta - \beta', \beta') \in A_2,$$

we have

$$\max_{\alpha + \beta - r \leq \beta' \leq r, i=1,2} |D_e^{\alpha + \beta - \beta'} D_{e_{i,2}}^{\beta'} (f - s)(v_2)| \leq \max_{\substack{(\alpha', \beta') \in A_2 \\ \alpha' + \beta' = \alpha + \beta, i=1,2}} |D_e^{\alpha'} D_{e_{i,2}}^{\beta'} (f - s)(v_2)|,$$

and (3.13) follows.

2) This part can be established by exactly the same arguments, the only difference being that the terms

$$h^{\alpha + \beta} |D_e^\alpha D_{e_{1,1}}^\beta (f - s)(v_1)|, \quad (\alpha, \beta) \in A_3, \quad \beta \leq \mu,$$

now also have to be estimated. ■

Let  $T \in \Delta$  and let  $v$  be a vertex of  $T$ . Then  $T = T_v^i$  for some  $i \in \{1, \dots, n(v)\}$ , where  $T_v^1, \dots, T_v^{n(v)}$  are all triangles attached to  $v$  and numbered counterclockwise, as in the definition of  $\mathcal{N}(v)$  (see Section 2). We are going to define various subsets of  $\mathcal{N}(v)$  and  $\mathcal{N}$  that will be instrumental in the proof of Theorem 3.1 and the key Lemma 3.8.

We define  $\mathcal{N}_T(v) \subset \mathcal{N}(v)$  to be the set of nodal functionals corresponding to the following nodal values:

(vt1)  $D_x^\alpha D_y^\beta f(v)$  for all  $(\alpha, \beta) \in A_1$ ,

(vt2)  $D_{e_i}^\alpha D_{e_{i+1}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$  if  $e_i$  is nondegenerate at  $v$ , or

$D_{e_{i-1}}^\alpha D_{e_i}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$  if  $e_i$  is degenerate at  $v$ , but  $e_{i-1}$  is nondegenerate at  $v$ , or

$D_{e_{i-2}}^\alpha D_{e_{i-1}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$  if both  $e_i$  and  $e_{i-1}$  are degenerate at  $v$ , but  $e_{i-2}$  is nondegenerate at  $v$ , or

$D_{e_1}^\alpha D_{e_2}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$  if  $v$  is a singular vertex,

- (vt3)  $D_{e_{i+1}}^\alpha D_{e_{i+2}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$  if  $e_{i+1}$  is a nondegenerate at  $v$  interior edge, or  
 $D_{e_{i+1}}^\alpha D_{e_{i+2}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_3$  if  $e_{i+1}$  is degenerate at  $v$ , or  
 $D_{e_{i+1}}^\alpha D_{e_i}^\beta f(v)$  for all  $(\alpha, \beta) \in A_3$  if  $e_{i+1}$  is a boundary edge, and
- (vt4)  $D_{e_i}^\alpha D_{e_{i+1}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_3$  if either  $e_i$  is degenerate at  $v$  or  $e_i$  is a boundary edge, or  
 $D_{e_{i-1}}^\alpha D_{e_i}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$  if both  $e_i$  and  $e_{i-1}$  are nondegenerate at  $v$ , or  
 $D_{e_{i-2}}^\alpha D_{e_{i-1}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$  if  $e_i$  is nondegenerate at  $v$ ,  $e_{i-1}$  is degenerate at  $v$ , and  $e_{i-2}$  is again nondegenerate at  $v$ , or  
 $D_{e_{i-3}}^\alpha D_{e_{i-2}}^\beta f(v)$  for all  $(\alpha, \beta) \in A_2$  if  $e_i$  is nondegenerate at  $v$ , both  $e_{i-1}$  and  $e_{i-2}$  are degenerate at  $v$ , and  $e_{i-3}$  is nondegenerate at  $v$ .

Furthermore, denote by

$$\mathcal{N}_{T,j}(v) \subset \mathcal{N}_T(v), \quad j = 1, 2, 3, 4,$$

the set of functionals corresponding to the nodal values listed in (vt1), (vt2), (vt3) and (vt4) respectively.

We also define

$$\tilde{\mathcal{N}}_T(v) \subset \mathcal{N}_T(v)$$

as follows: if each of two edges  $e_i$  and  $e_{i+1}$  is either degenerate or lies on the boundary, then  $\tilde{\mathcal{N}}_T(v) := \emptyset$ , if  $e_{i+1}$  is an interior nondegenerate at  $v$  edge, but  $e_i$  is not, then  $\tilde{\mathcal{N}}_T(v) := \mathcal{N}_{T,2}(v) \cup \mathcal{N}_{T,3}(v)$ , if, conversely,  $e_i$  is an interior nondegenerate at  $v$  edge, but  $e_{i+1}$  is not, then  $\tilde{\mathcal{N}}_T(v) := \mathcal{N}_{T,2}(v) \cup \mathcal{N}_{T,4}(v)$ , and, finally, if both  $e_i$  and  $e_{i+1}$  are interior nondegenerate at  $v$  edges, then  $\tilde{\mathcal{N}}_T(v) := \mathcal{N}_{T,2}(v) \cup \mathcal{N}_{T,3}(v) \cup \mathcal{N}_{T,4}(v)$ .

For every triangle  $T \in \Delta$  with vertices  $v_1, v_2, v_3$  and edges  $e_1, e_2, e_3$ , let

$$\begin{aligned} \mathcal{N}_T &:= \bigcup_{i=1}^3 \mathcal{N}_T(v_i) \cup \bigcup_{i=1}^3 \mathcal{N}(e_i) \cup \mathcal{N}(T), \\ \tilde{\mathcal{N}}_T &:= \bigcup_{i=1}^3 \tilde{\mathcal{N}}_T(v_i). \end{aligned}$$

Finally, we need a set of nodal functionals  $\mathcal{N}_T^*$  of finite-element type. Let  $T \in \Delta$ , let  $v$  be a vertex of  $T$ , and let the edges  $e_1, e_2$  of  $T$  be attached to  $v$ . Then  $\mathcal{N}_T^*(v)$  is defined to be the set of nodal functionals corresponding to the nodal values

$$D_{e_1}^\alpha D_{e_2}^\beta f(v), \text{ for all } (\alpha, \beta) \in A_1 \cup A_2 \cup A_3 \cup \tilde{A}_3,$$

where

$$\tilde{A}_3 := \{(\alpha, \beta) \in \mathbb{Z}^2 : (\beta, \alpha) \in A_3\}.$$

Furthermore, for every edge  $e$  of  $T$  we define  $\mathcal{N}_T^*(e)$  to be the set of nodal functionals assigning to every function  $f \in C^r(\Omega)$  the following nodal values:

$$D_{\tau'}^\mu f(z_e^{\mu,1}), \dots, D_{\tau'}^\mu f(z_e^{\mu, \kappa_\mu}) \text{ for all } \mu = 0, \dots, r,$$

where  $z_e^{\mu,i}$  and  $\kappa_\mu$  are defined in (2.5) and (2.6) respectively, and  $\tau'$  is the unit vector in the direction from the middle point of  $e$  to the vertex of  $T$  opposite to  $e$ .

For every triangle  $T \in \Delta$  with vertices  $v_1, v_2, v_3$  and edges  $e_1, e_2, e_3$ , we set

$$\mathcal{N}_T^* := \bigcup_{i=1}^3 \mathcal{N}_T^*(v_i) \cup \bigcup_{i=1}^3 \mathcal{N}_T^*(e_i) \cup \mathcal{N}(T).$$

**Lemma 3.7** *We have*

$$(3.14) \quad \text{card } \mathcal{N}_T = \text{card } \mathcal{N}_T^* = \binom{q+2}{2}.$$

Moreover,  $\mathcal{N}_T^*$  is  $\Pi_q$ -unisolvent, i.e., for any real data  $a_\nu$ ,  $\nu \in \mathcal{N}_T^*$ , there exists a unique polynomial  $p \in \Pi_q$  such that  $\nu p = a_\nu$  for all  $\nu \in \mathcal{N}_T^*$ .

**Proof.** Obviously,

$$\text{card } \mathcal{N}_T^* = 3 \text{ card } A_1 + 3 \text{ card } A_2 + 6 \text{ card } A_3 + 3 \text{ card } \mathcal{N}(e) + \text{card } \mathcal{N}(T).$$

By (2.9) and some elementary computation, we obtain  $\text{card } \mathcal{N}_T^* = \binom{q+2}{2}$ . Furthermore,

$$\text{card } \mathcal{N}_T(v) = \sum_{j=1}^4 \text{card } \mathcal{N}_{T,j}(v) = \text{card } A_1 + 3 \text{ card } A_2 = \text{card } \mathcal{N}_T^*(v).$$

Hence,  $\text{card } \mathcal{N}_T = \text{card } \mathcal{N}_T^*$ , which proves (3.14). Particularly,  $\text{card } \mathcal{N}_T^* = \dim \Pi_q$ . Because of this, the second statement of the lemma will follow if we show that the

only polynomial satisfying  $\nu p = 0$  for all  $\nu \in \mathcal{N}_T^*$  is the zero function. Following the lines of the proof of Lemma 3.5, with  $f \equiv 0$  and  $s = p$ , we get

$$\|D_{e_i}^\gamma D_{e_i^+}^\mu p\|_{C(e_i)} = 0, \quad i = 1, 2, 3,$$

for all  $\mu = 0, \dots, r$  and  $\gamma = 0, \dots, q - \mu$ , and every edge  $e_i$  of  $T$ . Therefore,

$$p = (l_1 l_2 l_3)^{r+1} \tilde{p},$$

where  $l_1, l_2$  and  $l_3$  are linear polynomials such that  $e_i \subset \{(x, y) : l_i(x, y) = 0\}$ , and  $\tilde{p}$  is a polynomial in  $\Pi_{q-3r-3}$ . Then  $\nu \tilde{p} = 0$ , for all  $\nu \in \mathcal{N}(T)$ . Since  $\mathcal{N}(T)$  is  $\Pi_{q-3r-3}$ -unisolvent, we have  $\tilde{p} = 0$ , and hence,  $p = 0$ . ■

We also need some local geometric characteristics of the triangulation.

Let  $e$  be any interior edge of the triangulation  $\Delta$ , and let  $v$  and  $v'$  be its vertices. Denote by  $e_1$  and  $e_2$  the adjacent edges of  $e$  at  $v$ , and by  $\theta_i$  the angle between  $e$  and  $e_i$ ,  $i = 1, 2$ . We set

$$\theta_{e,v} := \min\{\theta_1, \theta_2\}, \quad \tilde{\theta}_{e,v} := |\pi - \theta_1 - \theta_2|.$$

If  $e$  is a boundary edge, then  $\theta_{e,v}$  denotes the angle between  $e$  and its unique adjacent edge at  $v$ . Furthermore,

$$\theta_e := \min\{\theta_{e,v}, \theta_{e,v'}\},$$

and for an interior edge  $e$ ,

$$\tilde{\theta}_e := \begin{cases} \tilde{\theta}_{e,v}, & \text{if } e \text{ is degenerate at } v', \\ \min\{\tilde{\theta}_{e,v}, \tilde{\theta}_{e,v'}\}, & \text{if } e \text{ is nondegenerate at both } v \text{ and } v'. \end{cases}$$

(We note that no edge can be degenerate at both endpoints simultaneously.)

For every triangle  $T \in \Delta$  we denote by

$$\theta_T \quad \text{and} \quad \tilde{\theta}_T$$

the minimum of  $\theta_e$  over all edges of  $T$ , and the minimum of  $\tilde{\theta}_e$  over all edges of  $T$  lying in the interior of  $\Omega$ , respectively. Thus,  $\theta_T$  denotes the smallest angle around  $T$ , whereas  $\tilde{\theta}_T$  measures the “near-degeneracy” of the edges of  $T$ . Certainly,

$$\theta_T \geq \theta_\Delta.$$

The following key lemma shows that the nodal functionals in  $\mathcal{N}_T^*$  can be estimated in terms of those in  $\mathcal{N}_T$ . Moreover, only the contribution of  $\tilde{\mathcal{N}}_T$  to this estimation is influenced by  $\tilde{\theta}_T$ .

**Lemma 3.8** *Let  $T \in \Delta$ ,  $s \in S_q^{r,\rho}(\Delta)$  and  $f \in C^m(\Omega)$  ( $m \in \{2r, \dots, q+1\}$ ). Then for any  $\nu^* \in \mathcal{N}_T^*$*

$$(3.15) \quad \begin{aligned} |\nu^*(f-s)| &\leq Kh_T^{-d(\nu^*)} \left( h_T^m \max_{0 \leq m' \leq m} \|D_x^{m'} D_y^{m-m'} f\|_{C(\partial T)} \right. \\ &\left. + \max_{\nu \in \mathcal{N}_T \setminus \tilde{\mathcal{N}}_T} h_T^{d(\nu)} |\nu(f-s)| + \sin^{-r} \tilde{\theta}_T \max_{\nu \in \tilde{\mathcal{N}}_T} h_T^{d(\nu)} |\nu(f-s)| \right), \end{aligned}$$

where  $h_T$  is the diameter of  $T$ , and  $K$  depends only on  $r$ ,  $q$  and  $\theta_T$ .

**Proof.** Since  $\mathcal{N}(T) \subset \mathcal{N}_T^* \cap \tilde{\mathcal{N}}_T$ , we do not need to estimate  $|\nu^*(f-s)|$  for  $\nu^* \in \mathcal{N}(T)$ . Moreover, by symmetry, it is enough to consider  $\mathcal{N}_T^*(v)$  for a vertex  $v$  of  $T$ , and  $\mathcal{N}_T^*(e)$  for an edge  $e$  of  $T$ .

Let  $T = T_v^i$  for some  $i \in \{1, \dots, n(v)\}$ . Then  $\mathcal{N}_T^*(v)$  corresponds to the nodal values

$$\nu^* g = D_{e_i}^\alpha D_{e_{i+1}}^\beta g(v), \text{ for all } (\alpha, \beta) \in A_1 \cup A_2 \cup A_3 \cup \tilde{A}_3.$$

We consider three cases.

**Case 1:**  $(\alpha, \beta) \in A_1$ .

Then

$$\begin{aligned} h_T^{\alpha+\beta} |D_{e_i}^\alpha D_{e_{i+1}}^\beta (f-s)(v)| &\leq 2^{\alpha+\beta} h_T^{\alpha+\beta} \max_{\alpha' \geq 0, \beta' \geq 0, \alpha'+\beta'=\alpha+\beta} |D_x^{\alpha'} D_y^{\beta'} (f-s)(v)| \\ &\leq 2^\rho \max_{(\alpha', \beta') \in A_1} h_T^{\alpha'+\beta'} |D_x^{\alpha'} D_y^{\beta'} (f-s)(v)|. \end{aligned}$$

Therefore,

$$(3.16) \quad h_T^{\alpha+\beta} |D_{e_i}^\alpha D_{e_{i+1}}^\beta (f-s)(v)| \leq 2^\rho \max_{\nu \in \mathcal{N}_{T,1}(v)} h_T^{d(\nu)} |\nu(f-s)|, \quad (\alpha, \beta) \in A_1,$$

which proves (3.15).

**Case 2:**  $(\alpha, \beta) \in A_2$ .

If  $e_i$  is nondegenerate at  $v$ , then  $\nu^* \in \mathcal{N}_{T,2}(v)$  and (3.15) trivially holds. If  $e_i$  is degenerate at  $v$ , but  $e_{i-1}$  is nondegenerate at  $v$ , then, by (3.4),

$$h_T^{\alpha+\beta} |D_{e_i}^\alpha D_{e_{i+1}}^\beta (f-s)(v)| = h_T^{\alpha+\beta} |D_{e_i}^\alpha D_{e_{i-1}}^\beta (f-s)(v)| \leq \max_{\nu \in \mathcal{N}_{T,2}(v)} h_T^{d(\nu)} |\nu(f-s)|.$$

Similarly, if both  $e_i$  and  $e_{i-1}$  are degenerate at  $v$ , but  $e_{i-2}$  is nondegenerate at  $v$ , then a repeated application of (3.4) shows that

$$h_T^{\alpha+\beta} |D_{e_i}^\alpha D_{e_{i+1}}^\beta (f-s)(v)| = h_T^{\alpha+\beta} |D_{e_{i-2}}^\alpha D_{e_{i-1}}^\beta (f-s)(v)| \leq \max_{\nu \in \mathcal{N}_{T,2}(v)} h_T^{d(\nu)} |\nu(f-s)|.$$

Finally, if  $v$  is singular, then in the same manner we can see that

$$(3.17) \quad h_T^{\alpha+\beta} |D_{e_i}^\alpha D_{e_{i+1}}^\beta (f-s)(v)| \leq \max_{\nu \in \mathcal{N}_{T,2}(v)} h_T^{d(\nu)} |\nu(f-s)|, \quad (\alpha, \beta) \in A_2,$$

which, hence, holds in either case and confirms (3.15).

**Case 3:**  $(\alpha, \beta) \in A_3 \cup \tilde{A}_3$ .

By symmetry, assume without loss of generality that  $(\alpha, \beta) \in A_3$ .

If either  $e_i$  is degenerate at  $v$  or  $e_i$  is a boundary edge, then  $\nu^* \in \mathcal{N}_{T,4}(v)$  and (3.15) trivially holds. If, otherwise,  $e_i$  is a nondegenerate at  $v$  interior edge, then analysis similar to that in Case 2 shows that

$$(3.18) \quad h_T^{\alpha+\beta} |D_{e_i}^\alpha D_{e_{i-1}}^\beta (f-s)(v)| \leq \max_{\nu \in \mathcal{N}_{T,4}(v)} h_T^{d(\nu)} |\nu(f-s)|, \quad (\alpha, \beta) \in A_2.$$

Let us denote by  $v_{i-1}$ ,  $v_i$  and  $v_{i+1}$  the vertices of  $e_{i-1}$ ,  $e_i$  and  $e_{i+1}$  different from  $v$ , respectively, by  $e'_{i+1}$  the edge between  $v_i$  and  $v_{i+1}$ , and by  $e'_{i-1}$  the edge between  $v_i$  and  $v_{i-1}$ . The same argumentation as in the above shows that

$$(3.19) \quad \begin{aligned} h_T^{\alpha+\beta} |D_{e_i}^\alpha D_{e'_{i+1}}^\beta (f-s)(v_i)| &\leq 2^\rho \max_{\nu \in \mathcal{N}_{T,1}(v_i)} h_T^{d(\nu)} |\nu(f-s)|, \quad (\alpha, \beta) \in A_1, \\ h_T^{\alpha+\beta} |D_{e_i}^\alpha D_{e'_{i-1}}^\beta (f-s)(v_i)| &\leq \max_{\nu \in \mathcal{N}_{T,2}(v_i)} h_T^{d(\nu)} |\nu(f-s)|, \quad (\alpha, \beta) \in A_2. \end{aligned}$$

If now  $e_i$  is nondegenerate at  $v_i$ , then by the definition of  $\mathcal{N}_T$ ,

$$(3.20) \quad \mathcal{N}_{T,3}(v_i) = \{\nu g = D_{e_i}^\alpha D_{e'_{i-1}}^\beta g(v_i) : (\alpha, \beta) \in A_2\}.$$

In view of (3.16)–(3.20) and Lemma 3.6, 2), with  $T_1 = T$  and  $T_2 = T_v^{i-1}$ , we have for every  $\mu = 0, \dots, r$  and  $\gamma = 0, \dots, m - \mu$ ,

$$(3.21) \quad \begin{aligned} h_T^{\gamma+\mu} \|D_{e_i}^\gamma D_{e_i^\perp}^\mu (f-s)\|_{C(e_i)} &\leq K_1 \left( h_T^m \max_{0 \leq \mu' \leq \mu} \|D_{e_i}^{m-\mu'} D_{e_i^\perp}^{\mu'} f\|_{C(e_i)} \right. \\ &+ \max_{\nu \in \mathcal{N}_T(e_i)} h_T^{d(\nu)} |\nu(f-s)| + \max_{\nu \in \mathcal{N}_{T,1}(v) \cup \mathcal{N}_{T,1}(v_i)} h_T^{d(\nu)} |\nu(f-s)| \\ &\left. + \sin^{-r} \tilde{\theta}_T \max_{\nu \in \mathcal{N}_{T,2}(v) \cup \mathcal{N}_{T,4}(v) \cup \mathcal{N}_{T,2}(v_i) \cup \mathcal{N}_{T,3}(v_i)} h_T^{d(\nu)} |\nu(f-s)| \right), \end{aligned}$$



where  $K_1$  depends only on  $q$  and  $\theta_T$ . Since

$$(3.22) \quad |\nu^*(f - s)| = |D_{e_i}^\alpha D_{e_{i+1}}^\beta (f - s)(v)| \leq 2^\beta \max_{0 \leq \mu \leq \beta} |D_{e_i}^{\alpha+\beta-\mu} D_{e_i^\perp}^\mu (f - s)(v)|,$$

(3.15) follows from (3.21).

If  $e_i$  is degenerate at  $v_i$ , then

$$(3.23) \quad \mathcal{N}_{T,3}(v_i) = \{\nu g = D_{e_i}^\alpha D_{e_{i-1}}^\beta g(v_i) : (\alpha, \beta) \in A_3\}.$$

By (3.4),

$$(3.24) \quad |D_{e_i}^\alpha D_{e_{i+1}}^\beta g(v_i)| = |D_{e_i}^\alpha D_{e_{i-1}}^\beta g(v_i)|, \quad (\alpha, \beta) \in A_3.$$

In view of (3.16)–(3.19), (3.23), (3.24) and Lemma 3.6, 1), with  $T_1 = T$  and  $T_2 = T_v^{i-1}$ , we have for every  $\mu = 0, \dots, r$  and  $\gamma = 0, \dots, m - \mu$ ,

$$(3.25) \quad \begin{aligned} & h_T^{\gamma+\mu} \|D_{e_i}^\gamma D_{e_i^\perp}^\mu (f - s)\|_{C(e_i)} \leq K_2 \left( h_T^m \max_{0 \leq \mu' \leq \mu} \|D_{e_i}^{m-\mu'} D_{e_i^\perp}^{\mu'} f\|_{C(e_i)} \right. \\ & + \max_{\nu \in \mathcal{N}_T(e_i)} h_T^{d(\nu)} |\nu(f - s)| + \max_{\nu \in \mathcal{N}_{T,1}(v) \cup \mathcal{N}_{T,1}(v_i)} h_T^{d(\nu)} |\nu(f - s)| \\ & \left. + \max_{\nu \in \mathcal{N}_{T,2}(v) \cup \mathcal{N}_{T,3}(v_i)} h_T^{d(\nu)} |\nu(f - s)| + \sin^{-r} \tilde{\theta}_T \max_{\nu \in \mathcal{N}_{T,2}(v) \cup \mathcal{N}_{T,4}(v)} h_T^{d(\nu)} |\nu(f - s)| \right), \end{aligned}$$

with  $K_2$  being dependent only on  $q$  and  $\theta_T$ . Therefore, (3.15) follows from (3.22).

Finally, let  $e$  be one of the edges of  $T$ , say  $e = e_i$ . Then for any  $\nu^* \in \mathcal{N}_T^*$ ,

$$(3.26) \quad |\nu^*(f - s)| = |D_{\tau'}^\mu (f - s)(z_{e_i}^{\mu,j})| \leq 2^\mu \max_{0 \leq \mu' \leq \mu} |D_{e_i}^{\mu-\mu'} D_{e_i^\perp}^{\mu'} (f - s)(z_{e_i}^{\mu,j})|,$$

and (3.15) follows from (3.21) or (3.25) if  $e_i$  is an interior edge of  $\Omega$ . If, otherwise,  $e_i$  is a boundary edge, then, similar to the above, Lemma 3.5 implies that

$$(3.27) \quad \begin{aligned} & h_T^{\gamma+\mu'} \|D_{e_i}^\gamma D_{e_i^\perp}^{\mu'} (f - s)\|_{C(e_i)} \leq K_3 \left( h_T^m \max_{0 \leq \mu'' \leq \mu'} \|D_{e_i}^{m-\mu''} D_{e_i^\perp}^{\mu''} f\|_{C(e_i)} \right. \\ & + \max_{\nu \in \mathcal{N}_T(e_i)} h_T^{d(\nu)} |\nu(f - s)| + \max_{\nu \in \mathcal{N}_{T,1}(v) \cup \mathcal{N}_{T,1}(v_i)} h_T^{d(\nu)} |\nu(f - s)| \\ & \left. + \max_{\nu \in \mathcal{N}_{T,2}(v) \cup \mathcal{N}_{T,4}(v) \cup \mathcal{N}_{T,2}(v_i) \cup \mathcal{N}_{T,3}(v_i)} h_T^{d(\nu)} |\nu(f - s)| \right), \end{aligned}$$

with  $K_3$  being dependent only on  $q$  and  $\theta_T$ , which, in view of (3.26), implies (3.15).  $\blacksquare$

In the following lemma we use standard finite-element techniques to get an estimation of  $\|s|_T\|_{C(T)}$  in terms of the nodal functionals  $\nu \in \mathcal{N}_T^*$ .

**Lemma 3.9** *If  $s \in S_q^{r,\rho}(\Delta)$  and  $T \in \Delta$ , then*

$$(3.28) \quad \|s|_T\|_{C(T)} \leq K \max_{\nu \in \mathcal{N}_T^*} h_T^{d(\nu)} |\nu s|,$$

where  $h_T$  is the diameter of  $T$ , and  $K$  depends only on  $q$ .

**Proof.** Let  $\hat{T}$  be a fixed triangle in the plane, say, the triangle with vertices  $\hat{v}_1 = (-\frac{1}{2}, 0)$ ,  $\hat{v}_2 = (\frac{1}{2}, 0)$ ,  $\hat{v}_3 = (0, \frac{\sqrt{3}}{2})$ . Although  $\hat{T}$  may be not in  $\Delta$ , it is easy to see that the set of nodal functionals  $\mathcal{N}_{\hat{T}}^*$  is well-defined for  $\hat{T}$ .

For every  $T \in \Delta$ , let  $B_T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be an affine mapping such that  $B_T(\hat{T}) = T$ . Then

$$B_T z = A_T z + b_T, \quad z \in \mathbb{R}^2,$$

where  $A_T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is an invertible linear mapping and  $b_T \in \mathbb{R}^2$ . Since  $\hat{T}$  contains a disk of radius  $\frac{\sqrt{3}}{6}$ ,

$$(3.29) \quad \|A_T\| \leq 2\sqrt{3} h_T.$$

For every  $\hat{\nu} \in \mathcal{N}_{\hat{T}}^*$ , say of the form

$$\hat{\nu} g = D_{\hat{\tau}_1}^\alpha D_{\hat{\tau}_2}^\beta g(\hat{z}_0),$$

let us define  $\nu$  by

$$\nu g := D_{\tau_1}^\alpha D_{\tau_2}^\beta g(z_0),$$

where

$$\tau_i = \|A_T \hat{\tau}_i\|^{-1} A_T \hat{\tau}_i, \quad i = 1, 2, \quad z_0 = B_T \hat{z}_0.$$

Then it is easy to check that

$$(3.30) \quad \hat{\nu} \in \mathcal{N}_{\hat{T}}^* \iff \nu \in \mathcal{N}_T^*.$$

Moreover, a standard computation shows that

$$(3.31) \quad D_{\hat{\tau}_1}^\alpha D_{\hat{\tau}_2}^\beta g(B_T \hat{z}_0) = \|A_T \hat{\tau}_1\|^\alpha \|A_T \hat{\tau}_2\|^\beta D_{\tau_1}^\alpha D_{\tau_2}^\beta g(z_0).$$

By Lemma 3.7,  $\mathcal{N}_T^*$  is  $\Pi_q$ -unisolvent. Therefore, for every  $\nu \in \mathcal{N}_T^*$  there exists a unique fundamental polynomial  $p_\nu \in \Pi_q$  such that

$$\nu^* p_\nu = \begin{cases} 1, & \text{if } \nu^* = \nu, \\ 0, & \text{if } \nu^* \in \mathcal{N}_T^* \setminus \{\nu\}. \end{cases}$$

Similarly, for every  $\hat{\nu} \in \mathcal{N}_{\hat{T}}^*$  there exists a unique fundamental polynomial  $\hat{p}_{\hat{\nu}} \in \Pi_q$  such that

$$\nu^* \hat{p}_{\hat{\nu}} = \begin{cases} 1, & \text{if } \nu^* = \hat{\nu}, \\ 0, & \text{if } \nu^* \in \mathcal{N}_{\hat{T}}^* \setminus \{\hat{\nu}\}. \end{cases}$$

It follows from (3.31) that

$$(3.32) \quad p_\nu(B_T z) \equiv \|A_T \hat{\tau}_1\|^\alpha \|A_T \hat{\tau}_2\|^\beta \hat{p}_{\hat{\nu}}(z).$$

We are now ready to prove (3.28). Since  $s|_T$  is a polynomial in  $\Pi_q$ , we have

$$s|_T = \sum_{\nu \in \mathcal{N}_T^*} (\nu s) p_\nu.$$

Therefore, by (3.32) and (3.29),

$$\begin{aligned} \|s|_T\|_{C(T)} &\leq \sum_{\nu \in \mathcal{N}_T^*} |\nu s| \cdot \|p_\nu\|_{C(T)} = \sum_{\nu \in \mathcal{N}_T^*} |\nu s| \cdot \|p_\nu \circ B_T\|_{C(\hat{T})} \\ &\leq \sum_{\nu \in \mathcal{N}_T^*} |\nu s| \cdot \|A_T\|^{d(\nu)} \|\hat{p}_{\hat{\nu}}\|_{C(\hat{T})} \\ &\leq \left( \sum_{\hat{\nu} \in \mathcal{N}_{\hat{T}}^*} (2\sqrt{3})^{2r} \|\hat{p}_{\hat{\nu}}\|_{C(\hat{T})} \right) \max_{\nu \in \mathcal{N}_T^*} h_T^{d(\nu)} |\nu s|, \end{aligned}$$

and (3.28) follows. ■

**Remark 3.10** It is not difficult to see that the triples  $(T, \Pi_q, \mathcal{N}_T^*)$ ,  $T \in \Delta$ , form an affine family of finite elements in the sense of [9, p. 87], and  $(\hat{T}, \Pi_q, \mathcal{N}_{\hat{T}}^*)$  plays the role of the reference finite element for this family. Particularly, (3.30) shows that  $(T, \Pi_q, \mathcal{N}_T^*)$  is affine-equivalent to  $(\hat{T}, \Pi_q, \mathcal{N}_{\hat{T}}^*)$ , with  $B_T$  being the corresponding affine mapping.

**Proof of Theorem 3.1.** It follows from Lemma 3.8 and Lemma 3.9 that the only spline  $s \in S_q^{r,\rho}(\Delta)$  that satisfies  $\nu s = 0$  for all  $\nu \in \mathcal{N}$ , is the zero function. In view of (2.8), a standard linear algebra argument shows that for any real data  $a_\nu$ ,  $\nu \in \mathcal{N}$ , there exists a unique spline  $s \in S_q^{r,\rho}(\Delta)$  such that  $\nu s = a_\nu$  for all  $\nu \in \mathcal{N}$ . Particularly, for every function  $f \in C^{2r}(\Omega)$  the Hermite type interpolation problem

$$\nu s = \nu f \quad \text{for all } \nu \in \mathcal{N}$$

has a unique solution  $s_f \in S_q^{r,\rho}(\Delta)$ , which proves the first statement of the theorem.

Let us fix a function  $f \in C^m(\Omega)$  and a triangle  $T \in \Delta$ . Without loss of generality assume that  $(0, 0) \in T$ . Then for the Taylor polynomial

$$\tilde{p}(x, y) := \sum_{j=0}^{m-1} \sum_{j'=0}^j \frac{D_x^{j'} D_y^{j-j'} f(0,0)}{j'!(j-j')!} x^{j'} y^{j-j'}$$

we have

$$(3.33) \quad \|D_x^\alpha D_y^\beta (f - \tilde{p})\|_{C(T)} \leq \frac{2^{m-\alpha-\beta}}{(m+1-\alpha-\beta)!} h_T^{m-\alpha-\beta} \max_{0 \leq m' \leq m} \|D_x^{m'} D_y^{m-m'} f\|_{C(T)}$$

for all  $\alpha, \beta \geq 0$ ,  $\alpha + \beta \leq m$ . As a consequence, for every  $\nu \in \mathcal{N}_T^*$ ,

$$(3.34) \quad |\nu(f - \tilde{p})| \leq K_1 h_T^{m-d(\nu)} \max_{0 \leq m' \leq m} \|D_x^{m'} D_y^{m-m'} f\|_{C(T)},$$

where  $K_1$  is a constant depending only on  $q$ .

We have

$$(3.35) \quad \|D_x^\alpha D_y^\beta (f - s_f)\|_{L_\infty(T)} \leq \|D_x^\alpha D_y^\beta (f - \tilde{p})\|_{C(T)} + \|D_x^\alpha D_y^\beta (\tilde{p} - s_f)\|_{L_\infty(T)}.$$

By the bivariate Markov inequality (see, for example, [10]),

$$(3.36) \quad \|D_x^\alpha D_y^\beta p\|_{C(T)} \leq K_2 (h_T \sin \theta)^{-\alpha-\beta} \|p\|_{C(T)} \quad \text{for all } p \in \Pi_q,$$

where  $h_T$  and  $\theta$  are the diameter and the smallest angle of  $T$ , respectively, and  $K_2$  depends only on  $q$ .

Since  $\tilde{p} - s_f \in S_q^{r,\rho}(\Delta)$  and  $(\tilde{p} - s_f)|_T \in \Pi_q$ , it follows from (3.36) and Lemma 3.9 that

$$(3.37) \quad \begin{aligned} \|D_x^\alpha D_y^\beta (\tilde{p} - s_f)\|_{L_\infty(T)} &\leq K_2 (h_T \sin \theta)^{-\alpha-\beta} \|(\tilde{p} - s_f)|_T\|_{C(T)} \\ &\leq K_3 h_T^{-\alpha-\beta} \max_{\nu^* \in \mathcal{N}_T^*} h_T^{d(\nu^*)} |\nu^*(\tilde{p} - s_f)|, \end{aligned}$$

where  $K_3$  depends only on  $q$  and  $\theta_T$ . By (3.34) and Lemma 3.8, since  $\nu(f - s_f) = 0$  for all  $\nu \in \mathcal{N}_T$ , we have for every  $\nu^* \in \mathcal{N}_T^*$

$$(3.38) \quad \begin{aligned} h_T^{d(\nu^*)} |\nu^*(\tilde{p} - s_f)| &\leq h_T^{d(\nu^*)} |\nu^*(\tilde{p} - f)| + h_T^{d(\nu^*)} |\nu^*(f - s_f)| \\ &\leq K_4 h_T^m \max_{0 \leq m' \leq m} \|D_x^{m'} D_y^{m-m'} f\|_{C(T)}, \end{aligned}$$

where  $K_4$  depends only on  $r, q$  and  $\theta_T$ .

Since  $\theta_T \geq \theta_\Delta$ , now (3.35), (3.33), (3.37) and (3.38) imply (3.2). ■

**Remark 3.11** It is easy to see from the above proof that Theorem 3.1 in fact holds with  $\theta_T$  in place of  $\theta_\Delta$ .

## 4 A Basis for $S_q^{r,\rho}(\Delta)$

Let  $\mathcal{N} = \{\nu_i\}_{i=1}^n$ , where  $n = \dim S_q^{r,\rho}(\Delta)$  in view of (2.8). For every  $f \in C^{2r}(\Omega)$ , it follows from Theorem 3.1 that the interpolating spline  $s_f \in S_q^{r,\rho}(\Delta)$  satisfying (3.1) can be represented as

$$(4.1) \quad s_f = \sum_{i=1}^n (\nu_i f) s_i,$$

where the *fundamental functions*  $s_i \in S_q^{r,\rho}(\Delta)$ ,  $i = 1, \dots, n$ , are uniquely determined by the conditions

$$(4.2) \quad \nu_i s_j = \delta_{ij}, \quad i, j = 1, \dots, n.$$

Therefore,  $\{s_1, \dots, s_n\}$  is a basis for  $S_q^{r,\rho}(\Delta)$ . The following theorem establishes some useful properties of this basis.

**Theorem 4.1** *The fundamental functions  $s_1, \dots, s_n$  form a basis for  $S_q^{r,\rho}(\Delta)$  such that*

1)  $\{s_1, \dots, s_n\}$  is locally linearly independent, i.e., for every open  $B \subset \Omega$  the subsystem  $\{s_i : B \cap \text{supp } s_i \neq \emptyset\}$  is linearly independent on  $B$ ,

2)  $\{s_1, \dots, s_n\}$  is least supported, i.e., for every basis  $\{b_1, \dots, b_n\}$  of  $S_q^{r,\rho}(\Delta)$  there exists a permutation  $\pi$  of  $\{1, \dots, n\}$  such that

$$\text{supp } s_i \subset \text{supp } b_{\pi(i)}, \quad \text{for all } i = 1, \dots, n,$$

3) for each  $i = 1, \dots, n$ ,  $\text{supp } s_i$  is either a triangle or the union of some triangles sharing one common vertex,

4)  $K_1 h_i^{d(\nu_i)} \leq \|s_i\|_{C(\Omega)} \leq K_2 K_3 h_i^{d(\nu_i)}$ , where

$$h_i := \max_{T \subset \text{supp } s_i} h_T,$$

$K_1, K_2$  depend only on  $r, q$  and  $\theta_\Delta$ , and

$$K_3 = \begin{cases} 1, & \text{if } \nu_i \in \mathcal{N} \setminus \tilde{\mathcal{N}}, \\ \sin^{-r} \tilde{\theta}_\Delta, & \text{if } \nu_i \in \tilde{\mathcal{N}}, \end{cases}$$

with

$$\tilde{\mathcal{N}} := \bigcup_{T \in \Delta} \tilde{\mathcal{N}}_T, \quad \tilde{\theta}_\Delta := \min_{T \in \Delta} \tilde{\theta}_T,$$

5) the corresponding normalized basis  $\{s_1^*, \dots, s_n^*\}$ , with  $s_i^* := h_i^{-d(\nu_i)} s_i$ , is stable in the sense that

$$(4.3) \quad K_4 \max_i |a_i| \leq \left\| \sum_{i=1}^n a_i s_i^* \right\|_{C(\Omega)} \leq \frac{K_5}{\sin^r \tilde{\theta}_\Delta} \max_i |a_i|,$$

where  $K_4$  and  $K_5$  depend only on  $r, q$  and  $\theta_\Delta$ .

**Proof.** 1) As shown in [11], a system of functions  $\{s_1, \dots, s_n\} \subset S_q^r(\Delta) \setminus \{0\}$  such that  $\Pi_q \subset \text{span} \{s_1, \dots, s_n\}$  is locally linearly independent if and only if

$$(4.4) \quad \text{card} \{i : T \subset \text{supp } s_i\} \leq \binom{q+2}{2} = \dim \Pi_q, \quad \text{for every } T \in \Delta.$$

Since  $\Pi_q \subset S_q^{r,\rho}(\Delta)$ , we only have to check (4.4). Fix a triangle  $T$  in  $\Delta$ . By applying consecutively Lemma 3.9 and Lemma 3.8 (the latter with  $f \equiv 0$ ), we get

$$(4.5) \quad \|s_{i|_T}\|_{C(T)} \leq K_6 \left( \max_{\nu \in \mathcal{N}_T \setminus \tilde{\mathcal{N}}_T} h_T^{d(\nu)} |\nu s_i| + \sin^{-r} \tilde{\theta}_T \max_{\nu \in \tilde{\mathcal{N}}_T} h_T^{d(\nu)} |\nu s_i| \right),$$

where  $K_6$  depends only on  $r, q$  and  $\theta_T$ . Therefore,  $s_{i|_T} \equiv 0$  if  $\nu_i \notin \mathcal{N}_T$ , so that (4.4) follows from (3.14).

2) Least supportedness of  $\{s_1, \dots, s_n\}$  follows from its local linear independence in view of [5, Theorem 3.4].

3) As we have seen,  $T \subset \text{supp } s_i$  implies  $\nu_i \in \mathcal{N}_T$ . Therefore, it suffices to show that for each fixed  $\nu \in \mathcal{N}$  the set

$$\mathcal{T}_\nu := \{T \in \Delta : \nu \in \mathcal{N}_T\}$$

consists either of a single triangle or of some triangles sharing one common vertex. Since

$$\mathcal{N} = \bigcup_v \mathcal{N}(v) \cup \bigcup_e \mathcal{N}(e) \cup \bigcup_T \mathcal{N}(T),$$

we consider several cases. First, if  $\nu \in \mathcal{N}(T)$  for some  $T \in \Delta$ , then obviously  $\mathcal{T}_\nu = \{T\}$ . If  $\nu \in \mathcal{N}(e)$  for some interior edge  $e$  of  $\Delta$ , then  $\mathcal{T}_\nu = \{T_1, T_2\}$ , where  $T_1$  and  $T_2$  are the two triangles sharing the edge  $e$ . (If  $e$  is a boundary edge, then, of course, there is only one such triangle  $T$ , and  $\mathcal{T}_\nu = \{T\}$ .) Finally, assume that  $\nu \in \mathcal{N}(v)$  for some vertex  $v$  of  $\Delta$ . Then  $\nu \in \mathcal{N}_T$  implies  $\nu \in \mathcal{N}_T(v)$ . The latter is possible only for the triangles  $T$  that are attached to  $v$ . Hence,  $\mathcal{T}_\nu \subset \{T \in \Delta : v \in T\}$ .

4) By (4.5),

$$\|s_i\|_{C(\Omega)} = \max_{T \subset \text{supp } s_i} \|s_{i|_T}\|_{C(T)} \leq K_6 K_3 \max_{T \subset \text{supp } s_i} h_T^{d(\nu_i)} = K_6 K_3 h_i^{d(\nu_i)},$$

which gives the upper bound. Furthermore, by Markov inequality (3.36),

$$1 = |\nu_i s_i| \leq K_7 h_T^{-d(\nu_i)} \|s_{i|_T}\|_{C(T)}, \quad \text{for some } T \subset \text{supp } s_i,$$

where  $K_7$  depends only on  $q$  and  $\theta_\Delta$ . It is not difficult to check that

$$\max_{T \subset \text{supp } s_i} h_T \leq K_8 \min_{T \subset \text{supp } s_i} h_T$$

where  $K_8$  depends only on  $q$  and  $\theta_\Delta$  (see, e.g., [18, Lemma 3.2]). Therefore,

$$\|s_i\|_{C(\Omega)} \geq K_7^{-1} K_8^{-2r} h_i^{d(\nu_i)},$$

and the lower bound is also shown.

5) We fix  $\{a_i\}_{i=1}^n$  and set  $s = \sum_{i=1}^n a_i s_i^*$ . Let  $z \in T \in \Delta$ . By (4.4) and (4.5) we have

$$\left| \sum_{i=1}^n a_i s_i^*(z) \right| \leq \max_i |a_i| \sum_{i=1}^n |s_i^*(z)| \leq \max_i |a_i| \binom{q+2}{2} K_6 \sin^{-r} \tilde{\theta}_T,$$

which proves the upper bound for  $\|s\|_{C(\Omega)}$ . Moreover, let  $|a_j| = \max_i |a_i|$ . Then

$$\nu_j s = h_j^{-d(\nu_j)} a_j.$$

Therefore, in view of Markov inequality, we have for some  $T' \subset \text{supp } s_j$ ,

$$|a_j| = h_j^{d(\nu_j)} |\nu_j s| \leq K_7 h_j^{d(\nu_j)} h_{T'}^{-d(\nu_j)} \|s|_{T'}\|_{C(T')} \leq K_7 K_8^{2r} \|s\|_{C(\Omega)},$$

which completes the proof. ■

**Remark 4.2** A similar interpolation scheme can be done for the superspline space  $S_q^{r,\rho}(\Delta)$  with any  $\rho$  within the range

$$r + \left\lfloor \frac{r+1}{2} \right\rfloor \leq \rho \leq \min \left\{ 2r, \left\lfloor \frac{q-1}{2} \right\rfloor \right\}.$$

The only necessary change in the construction is that in the definition of  $\mathcal{N}(e)$  one should take

$$\kappa_\mu := \min \{ q - 2\rho - 1 + \mu, q - 3r - 1 - (r - \mu) \bmod 2 \}.$$

All results of Section 3 and Section 4 remain valid.

We conclude the paper with a discussion of the results of Section 4.

First of all, it immediately follows from Theorem 3.1 that the norm of the interpolation operator  $s_f : C^{2r}(\Omega) \rightarrow S_q^{r,\rho}(\Delta)$  is bounded by a constant which depends only on  $r, q$  and the smallest angle  $\theta_\Delta$  in  $\Delta$ . On the other hand, it is easy to see that some of the fundamental functions  $s_i$  can grow unboundedly if the triangulation contains near-degenerate edges. (In Theorem 4.1 we could only estimate  $\|s_i\|_{C(\Omega)}$  with a constant depending on  $\tilde{\theta}_\Delta$ .) This seems to be controversial at first glance. We will try to explain this phenomenon. Let  $v$  be a vertex of the triangulation  $\Delta$ . The nodal values  $\nu$  in the set  $\mathcal{N}(v)$  are linearly independent, as we have shown, if they are considered as linear functionals on the spline space  $S_q^{r,\rho}(\Delta)$ . Contrary to this, the nodal values  $\nu \in \mathcal{N}(v)$  corresponding to the partial derivatives of the same order  $k$ , with  $\rho < k \leq 2r$ , do stay in a linear relation as linear functionals on the space  $C^{2r}(\Omega)$ . (Recall that  $S_q^{r,\rho}(\Delta)$  is not a subspace of  $C^{2r}(\Omega)$ .) Indeed, there exist exactly  $k+1$  linearly independent partial derivatives  $D_{\tau_1}^\alpha D_{\tau_2}^\beta (f)(v)$ , with  $\alpha + \beta = k$ , for any



$k$ -times differentiable function  $f$ , and we certainly have in  $\mathcal{N}(v)$  more than  $k+1$  nodal values of this type. As a consequence, the coefficients  $\nu_i f$  in (4.1) satisfy some linear relations reflecting the fact that  $f$  is  $2r$ -times differentiable at each vertex. This leads to some cancellations in the sum and makes possible estimation (3.2).

Let us also remark that, according to Theorem 4.1, 2), our basis is best possible for the space  $S_q^{r,\rho}(\Delta)$  in regard to the size of the supports of the basis functions. As mentioned in Introduction, it shares this property with the basis for  $S_q^{r,\rho}(\Delta)$  constructed in [16]. The bases in [7, 18] fail to be least supported, but they have the advantage that the stability inequality (4.3) holds for them without  $\sin^r \tilde{\theta}_\Delta$  in the right hand side, *i.e.*, they enjoy stability even in the presence of near-degenerate edges.

Finally, we note that the property of local linear independence established for our basis in Theorem 4.1, 1), plays an important role in the theory of almost interpolation (see [11, 13, 14]).

## References

- [1] M. H. Adam, *Bivariate Spline-Interpolation auf Crosscut-Partitionen*, Dissertation, Mannheim, 1995.
- [2] C. de Boor and K. Höllig, Bivariate box splines and smooth pp functions on a three direction mesh, *J. Comput. Appl. Math.* **9** (1983), 13–28.
- [3] C. de Boor and K. Höllig, Approximation power of smooth bivariate pp functions, *Math. Z.* **197** (1988), 343–363.
- [4] C. de Boor and R.-Q. Jia, A sharp upper bound on the approximation order of smooth bivariate pp functions, *J. Approx. Theory* **72** (1993), 24–33.
- [5] J. M. Carnicer and J. M. Peña, Least supported bases and local linear independence, *Numer. Math.* **67** (1994), 289–301.
- [6] C. K. Chui and T. X. He, On location of sample points in  $C^1$  quadratic bivariate spline interpolation, in “Numerical Methods of Approximation Theory,” (L. Collatz, G. Meinardus and G. Nürnberger, Eds.), ISNM 81, Birkhäuser, Basel, 1987, 30–43.

- [7] C. K. Chui, D. Hong and R.-Q. Jia, Stability of optimal order approximation by bivariate splines over arbitrary triangulations, *Trans. Amer. Math. Soc.* **347** (1995), 3301–3318.
- [8] C. K. Chui and M. J. Lai, On bivariate super vertex splines, *Constr. Approx.* **6** (1990), 399–419.
- [9] P. G. Ciarlet, “The finite element method for elliptic problems,” North-Holland, Amsterdam, 1978.
- [10] C. Coatmélec, Approximation et interpolation des fonctions différentiables de plusieurs variables, *Ann. Sci. Ecole Norm. Sup. (3)* **83** (1966), 271–341.
- [11] O. Davydov, Locally linearly independent basis for  $C^1$  bivariate splines of degree  $q \geq 5$ , in M. Daehlen, T. Lyche, L. L. Schumaker (eds.), *Mathematical Methods for Curves and Surfaces II*, Vanderbilt University Press, 1998, 71–78.
- [12] O. Davydov, G. Nürnberger and F. Zeilfelder, Approximation order of bivariate spline interpolation for arbitrary smoothness, *J. Comput. Appl. Math.* **90** (1998), 117–134.
- [13] O. Davydov, M. Sommer and H. Strauss, Locally linearly independent systems and almost interpolation, in: G. Nürnberger, J. W. Schmidt, G. Walz (eds.), *Multivariate Approximation and Splines*, pp. 59–72, ISNM, Birkhäuser, Basel, 1997.
- [14] O. Davydov, M. Sommer and H. Strauss, On almost interpolation and locally linearly independent bases, *East J. Approx.* **5** (1999), 67–88.
- [15] D. Hong, Spaces of bivariate spline functions over triangulation, *Approx. Theory Appl.* **7** (1991), 56–75.
- [16] A. Kh. Ibrahim and L. L. Schumaker, Super spline spaces of smoothness  $r$  and degree  $d \geq 3r + 2$ , *Constr. Approx.* **7** (1991), 401–423.
- [17] F. Jeeawock-Zedek, Interpolation scheme by  $C^1$  cubic splines on a non uniform type-2 triangulation of a rectangular domain, *C.R. Acad. Sci. Ser. I Math.* **314** (1992), 413–418.
- [18] M. J. Lai and L. L. Schumaker, On the approximation power of bivariate splines, *Advances in Comp. Math.* **9** (1998), 251–279.

- [19] J. Morgan and R. Scott, A nodal basis for  $C^1$  piecewise polynomials of degree  $n \geq 5$ , *Math. Comp.* **29** (1975), 736–740.
- [20] G. Nürnberger, “Approximation by Spline Functions,” Springer-Verlag, Berlin, Heidelberg, New York, 1989.
- [21] G. Nürnberger, Approximation order of bivariate spline interpolation, *J. Approx. Theory* **78** (1996), 117–136.
- [22] G. Nürnberger, O. Davydov, G. Walz and F. Zeilfelder, Interpolation by bivariate splines on crosscut partitions, in “Multivariate Approximation and Splines,” (G. Nürnberger, J. W. Schmidt and G. Walz, Eds.), ISNM, Birkhäuser, 1997, 189–203.
- [23] G. Nürnberger and Th. Riessinger, Lagrange and Hermite interpolation by bivariate splines, *Numer. Func. Anal. Optim.* **13** (1992), 75–96.
- [24] G. Nürnberger and Th. Riessinger, Bivariate spline interpolation at grid points, *Numer. Math.* **71** (1995), 91–119.
- [25] G. Nürnberger and G. Walz, Error analysis in interpolation by bivariate  $C^1$ -splines, *IMA J. Numer. Anal.* **18** (1998), 485–507.
- [26] L. L. Schumaker, On super splines and finite elements, *SIAM J. Numer. Anal.* **26** (1989), 997–1005.
- [27] Z. Sha, On interpolation by  $S_2^1(\Delta_{m,n}^2)$ , *Approx. Theory Appl.* **1**, (1985), 71–82.
- [28] Z. Sha, On interpolation by  $S_3^1(\Delta_{m,n}^2)$ , *Approx. Theory Appl.* **1**, (1985), 1–18.
- [29] G. Strang and G. J. Fix, “An analysis of the finite element method,” Prentice-Hall, Englewood Cliffs, N. J., 1973.
- [30] A. Ženíšek, Interpolation polynomials on the triangle, *Numer. Math.* **15** (1970), 283–296.

Oleg Davydov  
Justus-Liebig-Universität Giessen, Mathematisches Institut  
35392 Giessen, Germany  
davydov@math.uni-dortmund.de

Günther Nürnberger and Frank Zeilfelder  
Universität Mannheim, Fakultät für Mathematik und Informatik  
68131 Mannheim, Germany  
nuern@euklid.math.uni-mannheim.de  
zeilfeld@fourier.math.uni-mannheim.de