



**Reilly, John J. and Janssen, Xanne and Cliff, Dylan P. and Okely, Anthony D. (2015) Appropriateness of the definition of 'sedentary' in young children : whole-room calorimetry study. Journal of Science and Medicine in Sport. ISSN 1878-1861 ,  
<http://dx.doi.org/10.1016/j.jsams.2014.07.013>**

This version is available at <https://strathprints.strath.ac.uk/52799/>

**Strathprints** is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<https://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: [strathprints@strath.ac.uk](mailto:strathprints@strath.ac.uk)

1 **Appropriateness of the Definition of ‘Sedentary’ in Young Children: Whole-Room Calorimetry**

2 **Study**

3 John J Reilly<sup>1\*</sup>, Xanne Janssen<sup>1,2</sup>, Dylan P Cliff<sup>2</sup> and Anthony D Okely<sup>2</sup>.

4 1. School of Psychological Sciences and Health, University of Strathclyde, Glasgow, Scotland,  
5 G1 1XQ.

6 2. Interdisciplinary Educational Research Institute, Faculty of Social Sciences, University of  
7 Wollongong, Wollongong, New South Wales, Australia, 2522.

8

9 \*Author for correspondence and guarantor:

10 John J. Reilly

11 Physical Activity for Health Group, School of Psychological Sciences and Health, University of  
12 Strathclyde

13 Glasgow, Scotland, G1 1XQ.

14 Email: john.j.reilly@strath.ac.uk; Phone: 0141 548 4235

15

16

17 Word Count: 2266

18 Abstract Word Count: 163

19

20

21 Running head: Energy expenditure of sedentary activities

22 Conflicts of interest: none

23

24

25

26

27

28

29 **Abstract**

30 **Objective:** The present study aimed to measure the energy cost of three common sedentary activities  
31 in young children to test whether energy expended was consistent with the recent consensus definition  
32 of ‘sedentary’ as ‘any behaviour conducted in a sitting or reclining posture and with an energy cost  
33  $\leq 1.5$  metabolic equivalents (METs)’ (Sedentary Behavior Research Network, 2012).

34 **Methods:** Whole-room calorimetry measures of television viewing, sitting at a table drawing and  
35 reading, and sitting on the floor playing with toys were made in 40 young children (mean age 5.3  
36 years, SD 1.0).

37 **Results:** The energy cost of each sedentary activity was consistent with the recent consensus  
38 definition of sedentary: 1.17 METs (95% CI 1.07-1.27) for TV viewing; 1.38 METs (95% CI 1.30-  
39 1.46) for sitting at a table; and 1.35 METs (95% CI 1.28-1.43) for floor-based play.

40 **Conclusions:** Common sedentary activities in young children have energy costs which are consistent  
41 with the recent consensus definition of ‘sedentary’, and the present study is supportive of this  
42 definition.

43

44 **Keywords:** calorimetry; obesity; children; preschool; sedentary behaviour; measurement.

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60 **Introduction**

61

62 Epidemiological studies have recently established that time spent in sedentary behaviour (sitting)  
63 influences several major health outcomes in adults<sup>1,2</sup>. There may also be measurable health effects of  
64 sitting behaviour during childhood and adolescence<sup>3,4</sup>, and sitting behaviour during childhood and  
65 adolescence may influence adult sitting behaviour<sup>5,6</sup>. An international consensus has been reached  
66 recently on the definition of ‘sedentary’ as ‘any behaviour conducted in a sitting or reclining posture  
67 and with an energy cost  $\leq 1.5$  metabolic equivalents (METs)’.<sup>7</sup>

68

69 Newton et al<sup>8</sup> recently demonstrated, in a whole-room calorimetry (WRC) study of 25 overweight and  
70 obese African-American adults, that the energy cost of common sedentary activities performed when  
71 sitting upright was clustered tightly around 1.0 MET, suggesting that a definition of sedentary might  
72 usefully incorporate an energy expenditure threshold below 1.5 MET. However, Newton et al<sup>8</sup>  
73 expressed a concern over the generalisability of their findings, noting the need to extend research of  
74 this kind to other populations. Evidence on the energy cost of sedentary behaviours in children has  
75 focused largely on older children and adolescents, notably the studies by Harrell et al in 8-18y olds<sup>9</sup>,  
76 Ridley and Olds in 6-18y olds<sup>10</sup>, Puyau et al in 6-16y olds<sup>11</sup>, and Evenson et al (mean age 7.3y)<sup>12</sup>. The  
77 energy cost of 5 common sedentary activities reported by Harrell et al was consistent with the  
78 consensus definition, with the 95% confidence intervals not exceeding 1.5 METs<sup>9</sup>. The mean energy  
79 cost of four common sedentary activities was reported by Ridley and Olds, and this approached 1.5  
80 METs for only one of the four<sup>10</sup>. Puyau et al found that the range of energy expenditure exceeded 1.5  
81 METs for one of the two sedentary activities studied<sup>11</sup>. Evenson et al reported the mean energy cost of  
82 two sedentary activities as well below 1.5 METs<sup>12</sup>.

83

84 In pre-school-aged children (3-5 years), evidence on the energy cost of common sedentary behaviours  
85 is scarce. We have been able to find only a single study, in which Adolph et al<sup>13</sup> used WRC to  
86 measure the energy cost of two sedentary activities: reclining watching TV; sitting and colouring. The  
87 range of energy cost of watching TV did not exceed 1.5 METs, but the range of energy costs for

88 sitting and colouring did (mean 1.4, SD 0.2). The primary aim of the present study was therefore to  
89 test whether the energy cost of common sedentary activities was consistent with a '1.5 MET  
90 threshold'<sup>7</sup> definition of sedentary behaviour in a sample of young children.

91

## 92 **Methods**

93 The present study was based on a sample of forty healthy 4- to 6-year-old children and was part of a  
94 larger study that aimed to validate various objective methods of estimation of free-living energy  
95 expenditure and physical activity in young children. Children were recruited from childcare centers  
96 (pre-schools, long-day and family-day care) in the Illawarra region of New South Wales, Australia.  
97 Exclusion criteria included the child having a disease known to influence their energy balance (e.g.  
98 hypothyroidism), a physical disability, or claustrophobia. No children were excluded on these  
99 grounds. The study was approved by the University of Wollongong/ SESIAHS Health and Medical  
100 Human Research Ethics Committee and all participating parents provided informed written consent  
101 and their children assented to participation.

102

103 Whole-room calorimetry provides a criterion measure of physical activity, energy expenditure, by  
104 measurement of oxygen consumption and carbon dioxide production ('calorimetry') while study  
105 participants are confined within a room (the calorimeter). Moreover, WRC avoids the need for face  
106 masks for collection of expired air which can be problematic in young children, and the avoidance of  
107 facemask- based collection systems combined with the amount of space within the WRC allows  
108 young children to behave in a fairly natural way. In the present study all children had a familiarisation  
109 visit to the WRC before the measurement. On the morning of measurement, parents were asked to  
110 give their children a standardised breakfast provided by the researchers (170 kcal) at 07.00 h and only  
111 give them sips of water thereafter<sup>14</sup>. Children and their parents arrived at the laboratory at  
112 approximately 08.15 h before entering the WRC at around 08.30 h. For the present study of sedentary  
113 behavior children spent ~70 minutes in the WRC, but this was nested within a more extended protocol  
114 of ~150 minutes which included activities of light and moderate-vigorous intensity which are  
115 described elsewhere<sup>14</sup>. We have established that giving a small standardised breakfast has a negligible

116 impact on subsequent measures of energy expenditure within the WRC<sup>14</sup>, and no decline in energy  
117 expenditure associated with declining diet-induced thermogenesis was detectable<sup>14</sup>.

118

119 Children's height and weight were measured using standardised procedures. Height was measured to  
120 the nearest 0.1 cm using a portable stadiometer (PE87, Mentone Educational Centre, Victoria,  
121 Australia) and weight was measured to the nearest 0.1 kg using a calibrated electronic scale (Tanita  
122 BC-418A, Tanita Corporation of America, Illinois, USA). Children then entered the WRC and were  
123 asked to follow a protocol which consisted of: sitting watching TV/DVD, sitting at a table while  
124 talking on the phone, reading, colouring, drawing, and sitting playing with toys on the floor.

125 According to the compendium of energy expenditure for children playing with toys was classified as a  
126 light physical activity<sup>15</sup>. However, in the current study it was completed while staying in a seated  
127 position and therefore it was included as a possible sedentary behaviour. The duration and order of the  
128 activities was pre-set and the same for each child (Table 1). Children were requested to complete one  
129 activity before moving on to the next. Children were not requested or instructed to sit still, but simply  
130 to complete the activity while in a seated position as they would do in a free-living situation. A degree  
131 of variation in the ways each behaviour in the protocol were carried out is inevitable, but energy  
132 expenditure data were only included in the present study if direct observation confirmed that they  
133 were sitting and carrying out the behaviour required at each stage of the protocol. All children were  
134 guided through the protocol by a research assistant who observed through a window and  
135 communicated via an intercom. The research assistant was able to encourage compliance with the  
136 protocol, and compliance was confirmed independently by filming and direct observation<sup>16,17</sup>.  
137 'Compliance' as defined by the filmed record represented periods when the child was following the  
138 protocol (i.e. completing the activity while in a seated position).

139

140 Oxygen consumption ( $\text{VO}_2$ ) and carbon dioxide production ( $\text{VCO}_2$ ) were measured continuously  
141 (paramagnetic  $\text{O}_2$  and infrared  $\text{CO}_2$  analyzers, Sable System Inc, Las Vegas USA) and corrected to  
142 standard temperature, pressure and humidity in the room calorimeter (3m x2.1m x2.1m) at the  
143 University of Wollongong. Technical procedures are described in more detail elsewhere, along with

144 full details of the protocol which children followed in the present study<sup>14</sup>. Chamber air was sampled  
145 every two minutes and rates of O<sub>2</sub> consumption and CO<sub>2</sub> production were calculated from in- and  
146 outflow as described in the literature<sup>18</sup>. Rates of O<sub>2</sub> consumption and CO<sub>2</sub> production were then  
147 averaged over 10 min to produce stable measures of EE,<sup>14,16</sup> and rates of energy expenditure were  
148 calculated using the Weir equation<sup>19</sup>.

149

150 Individualised MET values were calculated by dividing measured energy expenditure for each child  
151 by their predicted BMR. BMR was calculated using the following equations developed by Schofield  
152 et al.<sup>20</sup> in children aged 3- to 10-years:

153

$$154 \text{ BMR}_{\text{boys}} = 0.082 * \text{weight (kg)} + 0.545 * \text{height (m)} + 1.736$$

$$155 \text{ BMR}_{\text{girls}} = 0.071 * \text{weight (kg)} + 0.677 * \text{height (m)} + 1.553$$

156

157 The BMR was calculated as MJ/day and converted to kcal/kg/min. The Schofield equation was used  
158 because of the practical difficulties associated with obtaining BMR measures in children of this age<sup>21-</sup>  
159 <sup>23</sup>, and in one study it had no bias relative to measured BMR in a small sample of young children<sup>21</sup>.

160

161 Participants were filmed during the protocol. Video footage was coded using Vitessa 0.1 (Version 0.1,  
162 University of Leuven, Belgium) which generated a time stamp every time a change in posture or  
163 intensity was coded by the observer<sup>24</sup>. Every second following a given time stamp was coded as being  
164 at the same posture as that occurring at the point of the time stamp itself. Each second was coded in  
165 this way until a change in posture was indicated by the appearance of the next time stamp. This  
166 resulted in second-by-second coding. Children's postures were classified as sit/lie (i.e. sedentary) or  
167 non-sedentary. Postures were classified as sit/lie whenever the child's bottom touched the ground, a  
168 chair, or their legs (e.g. kneeling on both knees with their bottom touching the legs or heels)<sup>25,26</sup>. The  
169 postural coding meant that, as far as possible, only sedentary behaviours were included in the energy  
170 expenditure data.

171

172 Average energy expenditure values collected during watching television, sitting at a table, and playing  
173 with toys on the floor were used for analyses. Ten-minute data points were defined as valid if during  
174 these 10 minutes children participated consistently in sedentary activity (i.e. watching television,  
175 sitting at a table, playing with toys on the floor) as confirmed by direct observation<sup>14,16,17</sup>. Repeated-  
176 measures ANOVA with Bonferroni adjustment was used to compare energy expenditure between  
177 each of the sedentary activities and predicted BMR using the Schofield equation. In addition, to  
178 compare the effect of using predicted BMR and measured REE on energy expenditure values during  
179 sitting at a table and playing with toys, paired sample t-tests were used.

180

## 181 **Results**

182

183 Of the 40 children who participated in the study, two had missing data due to calorimeter malfunction.  
184 For the remaining 38 children, 34 (92.1%), 28 (73.7%), and 35 (92.1%) had at least one 10-min block  
185 of watching television, sitting at a table, and playing with toys on the floor, respectively. Missing data  
186 were due to children breaking up the sedentary activity by moving to a non-sit/lie position.

187 Descriptive characteristics for the study sample are presented in Table 2.

188

189 Energy expenditure data are shown in Table 3. Energy expenditure values for watching TV, sitting at  
190 a table and playing with toys were 0.037 kcal/kg/min ( $\pm 0.010$ ), 0.044 kcal/kg/min ( $\pm 0.008$ ) and  
191 0.043 kcal/kg/min ( $\pm 0.009$ ), respectively ( $P < 0.05$ ). Post-hoc analyses showed that measured energy  
192 expenditure while sitting at a table and playing with toys on the floor was significantly higher than  
193 while watching television ( $P < 0.05$ ).

194

195 For each of the three sedentary activities, average values were consistently  $< 1.5$  METs when  
196 predicted BMR was used to define 1 MET. The majority of the data points which exceeded 1.5 times  
197 predicted BMR were observed while sitting at a table ( $n = 8$  out of 28) and playing with toys on the  
198 floor ( $n = 11$  out of 35). The 95% confidence intervals for the three sedentary activities, expressed as



199 multiples of predicted BMR were as follows: 1.07 – 1.27; 1.30-1.46; 1.28 – 1.43 for watching TV,  
200 sitting at a table and playing with toys on the floor, respectively.

201

202

## 203 **DISCUSSION**

204 Consistency with the current ‘1.5 MET’ definition of sedentary was high in the present study: the  
205 95% confidence intervals for the energy cost of all three sedentary activities did not exceed 1.5 MET.

206 The behaviours included in the present study were common early childhood activities which were  
207 sedentary on postural grounds, and on energy expenditure grounds, suggesting a high degree of  
208 concordance between the postural and energy expenditure components of the current consensus  
209 definition of ‘sedentary’<sup>7</sup>.

210

211 A few studies in older children and adolescents<sup>7-10,27</sup> suggested that some sedentary activities had  
212 mean energy costs which exceed 1.5 METs, or a range which greatly exceeded 1.5 METs, but  
213 generally the evidence from older children and adolescents is consistent with the current consensus  
214 definition of ‘sedentary’<sup>9-12</sup>. The study of pre-school children (mean age 4.5y) by Adolph et al, which  
215 did not set out to test the appropriateness of the sedentary behaviour definition, found that the energy  
216 cost of sitting and colouring averaged 1.4 METs, with an SD of 0.2<sup>13</sup>. The 1.5 MET threshold to  
217 define sedentary behaviour is accepted as a mean, and a degree of inconsistency with it in certain  
218 populations, and for certain activities could presumably be tolerated.

219

220 The present study had a number of strengths. The use of a criterion method (energy expenditure) for  
221 validation of physical activity measures, the fairly natural setting, the combination of WRC with  
222 direct observation to confirm that activities within the WRC were occurring as instructed<sup>14,16,17</sup>, and  
223 the inclusion of three common sedentary activities, were notable strengths. One weakness was our  
224 inability to obtain a measured BMR measure which led to the need to use predicted BMR values, a  
225 problem common to almost all studies of young children<sup>22,22,28</sup>. However, study conclusions did not  
226 differ whether predicted values BMR were used, or measured resting metabolic rate (reclining in a

227 beanbag watching TV) values were used. The WRC measures were made after an overnight fast  
228 followed by a small standard breakfast. We have shown that providing a small standardised breakfast  
229 has no marked impact on energy expenditure >90 minutes later within the WRC, and so the  
230 measurement conditions approximate measurement in the post-absorptive state<sup>14</sup>. Many free-living  
231 sedentary behaviours will be carried out in a post-prandial state, and so will have energy costs slightly  
232 higher than those measured in the present study. Sit- to -stand postural transitions are very common  
233 among pre-school children during sedentary activities<sup>25,26,29</sup> and the exclusion of these from the  
234 measures made in the WRC (using direct observation) in the present study means that the estimates of  
235 the energy cost of these behaviours in the present study may be conservative. The present study could  
236 not include all forms of sedentary behaviour which young children experience<sup>30</sup>, and did not include  
237 screen-based gaming (some relatively new screen-based gaming devices appear to be particularly  
238 popular with young children)<sup>31</sup>. Finally, the dearth of evidence on the energy cost of common  
239 sedentary behaviours, with contemporary definitions of sedentary<sup>7,32</sup>, in a range of contemporary  
240 populations, means that generalisability of the present study should not be assumed.

241

## 242 **Conclusions**

243 In conclusion, the present study suggests that common ‘postural sedentary’ activities in young  
244 children have an energy cost which would mean that they can be considered as ‘sedentary’ according  
245 to the current consensus definition which incorporates energy expenditure. The present study is  
246 therefore supportive of the use of the current definition of ‘sedentary’ in young children<sup>7</sup>.

247

## 248 **Practical Implications**

- 249 • Defining sedentary behaviour in young children is important to evaluating interventions,  
250 understanding prevalence and trends, and assessing the health impact of sedentariness.
- 251 • The current consensus definition of sedentary behaviour is robust in young children and can  
252 be used with greater confidence as a result.

- 253       • Common sedentary behaviours in young children have energy costs between 1.2-1.4 METs,  
254           and energy expenditure during sedentary time can be estimated using these values.

255

256   **Acknowledgements**

257   We thank Harry Battam for his technical support and Melinda Smith for her assistance with  
258   recruitment and for leading the participants through the activity protocol. We also thank the children  
259   and their parents for their participation. This study was supported by the National Heart Foundation of  
260   Australia (project grant GIA09S4441; plus personal awards PH 11S 6025; CR 11S 6099).

261   **Conflicts of Interest: None Declared.**

262 **References**

- 263 1. Thorp AA, Owen N, Neuhaus M et al. Sedentary Behaviors and Subsequent Health Outcomes in  
264 Adults: A Systematic Review of Longitudinal Studies, 1996-2011. *Am J Prev Med* 2011; 41:207-215.
- 265 2. Tremblay MS, Colley RC, Saunders TJ et al . Physiological and health implications of a sedentary  
266 lifestyle. *Appl Physiol Nutr Metab* 2010; 35:725-740.
- 267 3. Tremblay MS, Le Blanc AG, Kho ME, Saunders TJ, Larouche R, Colley RC, Goldfield G, Connor-  
268 Gorber S. Systematic review of sedentary behaviour and health indicators in school-age children and  
269 youth. *Int J Behav Nutr Phys Act* 2011; 8:98.
- 270 4. Le Blanc AG, Spence JC, Carson V, Connor-Gorber S, Dillman C, Janssen I, Kho ME, Stearns JA,  
271 Timmons BW, Tremblay MS. Systematic review of sedentary behaviours and health indicators in the  
272 early years. *Appl Physiol Nutr Metab* 2012; 37:753-772.
- 273 5. Verloigne M, van Lippenvelde W, Maes L et al. Levels of physical activity and sedentary time  
274 among 10-12y old boys and girls across 5 European countries using accelerometers. *Int J Behav Nutr*  
275 *Phys Act* 2012; 9:34.
- 276 6. Mitchell JA, Pate RR, Dowda M et al . Prospective study of sedentary behavior in a large cohort of  
277 youth. *Med Sci Sports Exerc* 2012; 44:1081-7.
- 278 7. Sedentary Behaviour Research Network. Standardised use of the terms" sedentary" and" sedentary  
279 behaviours": letter to the editor. *Appl Physiol Nutr Metab* 2012; 37:540-542.
- 280 24.
- 281 8. Newton Jr RL, Han H, Zderic T et al. The Energy Expenditure of Sedentary Behavior: A Whole  
282 Room Calorimeter Study. *PLoS ONE* 2013; 8:e63171.
- 283 9. Harrell JS, McMurray RG, Baggett CD et al. Energy costs of physical activities in children and  
284 adolescents. *Med Sci Sports Exerc* 2005; 37:329-336.
- 285 10. Ridley K, and Olds TS. Assigning energy costs to activities in children: a review and synthesis.  
286 *Med Sci Sports Exerc* 2008; 40:1439-1446.
- 287 11. Puyau MR, Adolph AL, Vohra FA et al. Validation and calibration of physical activity monitors  
288 in children. *Obes Res* 2002; 10:150-157.

289 12. EvensonKR, Catellier DJ, Gill K et al Calibration of two objective measures of physical activity  
290 for children. *J Sports Sci* 2008; 26:1557-1568.

291 13. Adolph AL, Puyau MR, Vohra FA et al. Validation of uniaxial and triaxial accelerometers for the  
292 assessment of physical activity in preschool children. *J Phys Act Health* 2012; 9:944-953.

293 14. Janssen X, Cliff DP, Okely AD et al. Practical utility and reliability of whole-room calorimetry  
294 in young children. *Br J Nutr* 2013; 109:1917-1922.

295 15. Ridley K, Ainsworth BE, Olds TS. Development of a compendium of energy expenditures for  
296 youth. *Int J Behav Nutr Phys Act* 2008; 5:45.

297 16. Janssen X, Cliff D, Reilly JJ et al. Validation of activPAL™ Defined Sedentary Time and Breaks  
298 in Sedentary Time in 4-6 Year Olds. *Pediatr Exerc Sci* 2014; 26:110-117.

299 17. Janssen X, Cliff DP, Reilly JJ et al. Predictive validity and classification accuracy of ActiGraph  
300 energy expenditure equations and cut-points in young children. *PLoS ONE* 2013; 8:e79124.

301 18. Schoffelen PFM, Westerterp KR, Saris WHM et al. A dual-respiration chamber system with  
302 automated calibration. *J Appl Physiol* 1997; 83:2064-2072.

303 19. Weir JB. New methods for calculating metabolic rate with special reference to protein  
304 metabolism. *J Physiol* 1949; 109:1-9.

305 20. Schofield WN. Predicting basal metabolic rate, new standards and review of previous work. *Hum*  
306 *Nutr Clin Nutr* 1985; 39:5-41.

307 21. Reilly JJ, Jackson DM, Montgomery C et al. Total energy expenditure and physical activity in  
308 young Scottish children: mixed longitudinal study. *The Lancet* 2004; 363:211-212.

309 22. Jackson DM, Pace L, and Speakman JR. The Measurement of Resting Metabolic Rate in  
310 Preschool Children. *Obesity* 2007; 15:1930-1932.

311 23. Heil DP. Predicting Activity Energy Expenditure Using the Actical® Activity Monitor. *Res Q*  
312 *Exerc Sport* 2006; 77:64-80.

313 24. Van Puyenbroeck J, Maes B, and Laeremans P. Vitessa (Version 0.1). University of Leuven,  
314 Belgium., 2005, p. Computer software; retrieved from <http://ppw.kuleuven.be/ortho/vitessa>.accessed  
315 19th November 2013.

316 25. Davies G, Reilly JJ, and Paton JY. Objective measurement of posture and posture transitions in  
317 the pre-school child. *Physiol Meas* 2012; 33:1913-1921.

318 26. Davies G, Reilly JJ, McGowan A et al..Validity, practical utility, and reliability of the activPAL in  
319 pre-school children. *Med Sci Sports Exerc* 2012; 44:761-768.

320 27. Trost SG, Loprinzi PD, Moore R et al. Comparison of accelerometer cut points for predicting  
321 activity intensity in youth. *Med Sci Sports Exerc* 2011; 43:1360-1368.

322 28. Rodriguez G, Moreno LA, Sarria A, Fleta J, Bueno M. Resting energy expenditure in children and  
323 adolescents: agreement between calorimetry and prediction equations. *Clin Nutr* 2002; 21:255-260.

324 29. Alghaeed Z, Reilly JJ, Chastin SF et al. The Influence of minimum sitting period of the  
325 ActivPAL™ on the measurement of breaks in sitting in young children. *PLoS ONE* 2013; 8:e71854.

326 30. Reilly JJ. Physical activity, sedentary behaviour, and energy balance in the pre-school child. *Proc*  
327 *Nutr Soc* 2008; 67:317-325.

328 31. Christakis DA. Interactive media use under the age of 2 years. *JAMA Pediatr* 2014; 168:399-400.

329 32. Pate RR, O'Neill JR, Lobelo F. The evolving definition of 'sedentary'. *Exerc Sport Sci Rev* 2008;  
330 36:173-178.

331

332 **Table 1.** Whole room calorimetry protocol

Activity	Time (min)
Behaviour	
Watching TV–sitting in a beanbag	30
Talking on telephone with parents – sitting	2
Reading books with a cassette/CD – sitting	5
Drawing/colouring in – sitting	10
Playing with toys, blocks (Lego), dolls, puzzles, games – sitting on floor	20

333

334

335

336

337

338

339

340

341

342

343

344 **Table 2.** Characteristics of study participants, mean (SD).

	Total sample (n=40)	Boys (n=22)	Girls (n=18)
Age (years)	5.3 (1.0)	5.2 (1.0)	5.3 (1.1)
Height (cm)	112.7 (8.1)	114.3 (6.2)	110.9 ( 9.7)
Weight (kg)	20.6 (3.7)	21.5 (2.4)	19.4 (4.6)
BMI (kg/m <sup>2</sup> )	16.1 (1.5)	16.5 (1.3)	15.5 (1.6)
BMI z-score	0.34 (1.07)	0.66 (0.88)	-0.05 (1.18)

345

346



347 **Table 3.** Energy expenditure values for each sedentary activity, mean (SD).

	O <sub>2</sub> ml/kg/min	kcal/kg/min	METs <sup>a</sup>
Predicted BMR <sup>b</sup>		0.032 (0.003)	1.00
Watching TV	7.64 (2.47)	0.037 (0.010) <sup>†</sup>	1.17 (0.30) <sup>†</sup>
Sitting at a table	9.25 (1.82) <sup>*</sup>	0.044 (0.007) <sup>*†</sup>	1.38 (0.22) <sup>*†</sup>
Playing with toys	8.83 (2.10)	0.043 (0.009) <sup>*†</sup>	1.35 (0.23) <sup>*†</sup>

348 <sup>a</sup> METs defined as multiples of predicted basal metabolic rate using the Schofield equation<sup>19</sup>;

349 <sup>b</sup> predicted basal metabolic rate using the Schofield equation; \* p<0.05 compared to watching  
 350 television; † p<0.05 compared to predicted BMR

351

352