

1 **Lifetime exposure to ambient ultraviolet radiation and the risk for cataract**  
2 **extraction and age-related macular degeneration: the Alienor Study**

3

4 Cécile Delcourt<sup>1,2</sup>, Audrey Cougnard-Grégoire<sup>1,2</sup>, Mathieu Boniol<sup>3</sup>, Isabelle Carrière<sup>4,5</sup>, Jean-François  
5 Doré<sup>6</sup>, Marie-Noëlle Delyfer<sup>1,2,7</sup>, Marie-Bénédicte Rougier<sup>7</sup>, Mélanie Le Goff<sup>1,2</sup>, Jean-François  
6 Dartigues<sup>1,2</sup>, Pascale Barberger-Gateau<sup>1,2</sup>, Jean-François Korobelnik<sup>1,2,7</sup>.

7

8

9 1 : Univ. Bordeaux, F-33000 Bordeaux, France ;

10 2 : INSERM, ISPED, Centre INSERM U897-Epidemiologie-Biostatistique, F-33000 Bordeaux, France

11 3 : International Prevention Research Institute, Lyon, France

12 4 : Inserm, U1061, Montpellier, F-34093 France

13 5 : University Montpellier I, Montpellier, F-34000 France

14 6 : Centre de Recherche en Cancérologie de Lyon, UMR Inserm U1052 - CNRS U5286, Lyon, F-  
15 69008, France

16 7 : CHU de Bordeaux, Service d'Ophtalmologie, Bordeaux, F-33000, France

17

18

19 Corresponding author/reprint request : Cécile Delcourt, Inserm U897, ISPED, Université Bordeaux  
20 Segalen, 146 rue Léo Saignat, 33076 Bordeaux Cedex. Tel : +33 5 47 30 42 04; Fax : +33 5 57 57 14  
21 86 ; email : [cecile.delcourt@isped.u-bordeaux2.fr](mailto:cecile.delcourt@isped.u-bordeaux2.fr)

22

23 Financial support: This study received financial support from Laboratoires Théa (Clermont-Ferrand,  
24 France) and Fondation Voir et Entendre (Paris, France). Laboratoires Théa participated in the design  
25 of the study, but no sponsor participated in the collection, management, statistical analysis and  
26 interpretation of the data, nor in the preparation, review or approval of the present manuscript.  
27 Eurosun study received a funding from the European Union's Public Health Executive Agency under  
28 Grant Agreement No. 2006320 (EUROSUN project).

29

30 **ABSTRACT**

31 Purpose: While exposure to ultraviolet radiation is a recognized risk factor for cataract, its association  
32 is more controversial with age-related macular degeneration (AMD). We report the associations of  
33 lifetime exposure to ambient ultraviolet radiation (UVR) with cataract extraction and AMD.

34 Methods: The Alienor Study is a population-based study of 963 residents of Bordeaux (France), aged  
35 73 years or more. Lifetime exposure to ambient UVR was estimated from residential history and  
36 Eurosun satellite-based estimations of ground UVR. It was divided in 3 groups (lower quartile,  
37 intermediate quartiles, upper quartile), using the intermediate quartiles as the reference. Early and late  
38 AMD were classified from retinal colour photographs. Cataract extraction was defined as absence of  
39 the natural lens at slit lamp.

40 Results: After multivariate adjustment, subjects in the upper quartile of lifetime ambient UVR exposure  
41 were at increased risk for cataract extraction (OR=1.53, 95 % confidence interval (CI): 1.04-2.26,  
42 p=0.03) and for early AMD (OR= 1.59, 95 % CI: 1.04-2.44, p=0.03), by comparison with subjects in the  
43 intermediate quartiles. Subjects in the lower quartile of UVR exposure were also at increased risk for  
44 early AMD (OR=1.69, 95 % CI: 1.06-2.69, p=0.03), by comparison with those with medium exposure.  
45 Associations of late AMD with UVR exposure was not statistically significant.

46 Conclusions: This study further confirms the increased risk for cataract extraction in subjects exposed  
47 to high ambient UVR. Moreover, it suggests that risk for early AMD is increased in subjects exposed to  
48 high UVR, but also to low UVR, by comparison with medium exposures.

49

50 Keywords: macular degeneration, cataract, light exposure, ultraviolet radiation, risk factors,  
51 epidemiology

52

53

54 Cataract and age-related macular degeneration (AMD) are leading causes of blindness and visual  
55 impairment worldwide <sup>1</sup>. Both of these diseases are multifactorial, involving non-modifiable (e.g. age,  
56 gender, genetics) and modifiable factors <sup>2,3</sup>. The control of these modifiable factors may represent a  
57 preventive strategy for decreasing the incidence of these diseases and of the related visual  
58 impairment. Among other modifiable factors (in particular smoking and nutrition) <sup>2,3</sup>, the potential role  
59 of sunlight exposure in the aetiology of these diseases has been investigated <sup>4 5</sup>. The absorption of  
60 solar radiations by biological tissues results in photochemical reactions and the formation of reactive  
61 oxygen species (including singlet oxygen) which may damage all cellular components (lipids, proteins,  
62 DNA)<sup>6</sup>. The part of solar radiations that interacts with the eye is known as “optical radiations” and  
63 includes wavelengths from ultraviolet (100-400 nm), visible light (400-760 nm) to infrared (760-10,000  
64 nm). Ultraviolet radiations represent the most energetic part of optical radiations, and are thus  
65 responsible for a large part of photochemical damage.

66 The crystalline lens is particularly exposed to phototoxic damage, because it absorbs most of  
67 ultraviolet radiations (UVR), together with the cornea. This has been confirmed by epidemiological  
68 studies, which have shown consistent associations of cataract (resulting from lens opacification) with  
69 sunlight exposure and, in particular, UVR exposure <sup>7-13</sup>. , Ultraviolet exposure has been consistently  
70 associated with the risk for cataract in numerous studies, performed in different continents with  
71 different methodologies, showing dose-dependent relationships, and specific associations with cortical  
72 cataracts. It is now a recognized risk factor for cataract <sup>2</sup>.

73 By contrast, epidemiological data regarding the associations of light exposure with the risk for AMD  
74 remain scarce and inconsistent. Several studies have suggested an increased risk for AMD in subjects  
75 highly exposed to sunlight <sup>14-17</sup>, but others showed no significant associations <sup>18-20</sup> and some even  
76 suggested a decreased risk for AMD in the most exposed subjects <sup>21, 22</sup>.

77 In the present study, we report the associations of lifetime exposure to ambient UVR with the risk for  
78 cataract extraction and AMD, in the framework of a population-based cross-sectional study of elderly  
79 subjects from the South of France.

80

## 81 **Subjects and Methods**

### 82 ***Study aims***

83 The Alienor (Antioxydants, Lipides Essentiels, Nutrition et maladies OculaiRes) Study is a population-  
84 based prospective study aiming at assessing the associations of age-related eye diseases (AMD,  
85 glaucoma, cataract, dry eye syndrome) with nutritional factors (in particular antioxidants, macular  
86 pigment and fatty acids).<sup>23</sup> It also takes into account other major determinants of eye diseases,  
87 including gene polymorphisms, lifestyle and vascular factors.

88

### 89 ***Study sample***

90 Subjects of the Alienor Study were recruited from an ongoing population-based study on the vascular  
91 risk factors for dementia, the Three City (3C) Study<sup>24</sup>. The 3C Study included 9,294 subjects aged 65  
92 years or more from three French Cities (Bordeaux, Dijon and Montpellier), among which 2,104 were  
93 recruited in Bordeaux. Subjects were contacted individually from the electoral rolls. They were initially  
94 recruited in 1999-2001 and followed-up about every two years since. The Alienor study consists in eye  
95 examinations, which are offered to all participants from the third follow-up (2006-2008) of the 3C  
96 cohort in Bordeaux. Among the 1,450 participants re-examined in 2006-2008, 963 (66.4 %)   
97 participated in the first eye examination of the Alienor Study.

98 This research followed the tenets of the Declaration of Helsinki. Participants gave written consent for  
99 the participation in the study. The design of this study has been approved by the Ethical Committee of  
100 Bordeaux (Comité de Protection des Personnes Sud-Ouest et Outre-Mer III) in May 2006.

101

### 102 ***Eye examination***

103 The eye examination took place in the Department of Ophthalmology of the University Hospital of  
104 Bordeaux. It included a recording of ophthalmological history, measures of visual acuity, refraction,  
105 two 45° non mydriatic color retinal photographs (one centred on the macula, the other centred on the  
106 optic disc), measures of intraocular pressure and central corneal thickness and tear film break-up time  
107 test. A self-completed questionnaire on risk factors specific to the eye (including residential history for

108 estimation of light exposure) and dry eye symptoms was filled at home and brought back on the day of  
109 the eye examination.

110 Retinal photographs were performed using a high-resolution digital non mydriatic retinograph (TRC  
111 NW6S, Topcon, Japan). Photographs were interpreted in duplicate by two specially trained  
112 technicians. Inconsistencies between the two interpretations were adjudicated by a retina specialist for  
113 classification of AMD and other retinal diseases, or by a glaucoma specialist for classification of  
114 glaucoma. All cases of late AMD, other retinal diseases and glaucoma were reviewed and confirmed  
115 by specialists. Graders had no access to light exposure variables, or any of the potential confounders.

116

### 117 ***Cataract extraction***

118 In each eye, cataract extraction was defined as the absence of the natural crystalline lens at slit lamp.

119

### 120 ***Classification of AMD***

121 Retinal photographs were interpreted according to the international classification<sup>25</sup> and to a  
122 modification of the grading scheme used in the Multi-Ethnic Study of Atherosclerosis for drusen size,  
123 location and area<sup>26</sup>. Eyes were classified according into one of the three exclusive groups: no AMD,  
124 early AMD, late AMD.

125 Late AMD was defined by the presence of neovascular AMD or geographic atrophy within the grid  
126 (3000 microns from the foveola). Neovascular AMD included serous or hemorrhagic detachment of the  
127 retinal pigment epithelium (RPE) or sensory retina, subretinal or sub-RPE hemorrhages and fibrous  
128 scar tissue. Geographic atrophy was defined as a discrete area of retinal depigmentation, 175 microns  
129 in diameter or larger, characterized by a sharp border and the presence of visible choroidal vessels.  
130 Five cases of late AMD had no gradable photographs and were classified using ophthalmological  
131 history of AMD and AMD therapy (in particular intravitreal antiangiogenic agents and photodynamic  
132 therapy), and confirmed by their treating ophthalmologist.

133 Early AMD was defined by the presence of soft distinct drusen and/or soft indistinct drusen and/or  
134 reticular drusen and/or pigmentary abnormalities. Soft distinct and indistinct drusen were larger than

135 125 microns in diameter and with uniform density and sharp edges or decreasing density from the  
136 center outwards and fuzzy edges, respectively. Pigmentary abnormalities were defined as areas of  
137 hyperpigmentation and/or hypopigmentation (without visibility of choroidal vessels).

138

### 139 ***Ambient UVR exposure***

140 For each participant, average annual ambient UVR exposure was estimated using the residential  
141 history, by weighting annual ambient UVR at each location by the time spent at that location.

142 Residential history from birth (locations, time spent at each location) was self-declared up to the first  
143 eye examination (2006-2008). In France, locations were divided in 101 geographical areas,  
144 corresponding to the 101 administrative « departments » (95 in metropolitan France and 6 overseas  
145 departments). Concerning foreign countries, ambient UVR was generally estimated for the capital of  
146 the country, except when the capital was very off-centered, in which case a more central location was  
147 chosen. Very large countries (United States, China, etc) were excluded from this analysis, since an  
148 estimation of solar radiation in a single geographic location is meaningless.

149 Average annual ambient solar radiation was assessed for the first 65 years of life, since almost all  
150 subjects (97.3 %) lived in Bordeaux area beyond this age.

151

### 152 ***Ultraviolet radiation***

153 In each location, UVR was extracted from Eurosun UV database ([www.eurosun-project.org](http://www.eurosun-project.org)). Briefly,  
154 UV irradiation levels were initially extracted from surface solar irradiance derived from the Meteosat  
155 satellite's images. From these irradiation levels, the UV component was computed by a model which  
156 exploits the algorithm set up by the Royal Institute of Meteorology, Belgium, published in the European  
157 Solar Radiation Atlas (spectral model of Joukoff ESRA). The algorithm converts the total irradiance (E)  
158 into its spectral distribution  $E(\lambda)$ , every 10nm, and gives estimates for total UV (280-400 nm), UVA  
159 (315-400 nm) and UVB (280-315 nm). The calculation of individual exposure assumes that the  
160 irradiation levels in different regions remained constant over the years. The information provided by  
161 Eurosun corresponds to the average of daily UV radiation over the period 1988-2007 for each location.

162 We conducted analysis of change in UV radiation over the period 1988-2007 in Eurosun maps and  
163 found only minor changes of less than 1% yearly change. This observations were in line with the  
164 results obtained from COST 726 which also reported a stability of UV irradiation in Europe for a period  
165 of 50 years of data up to 2002 (the project COST 726's report and details are available at [http://www-  
166 med-physik.vu-wien.ac.at/uv/cost726/cost726.htm](http://www-med-physik.vu-wien.ac.at/uv/cost726/cost726.htm)). So these analyses from Eurosun data and  
167 observation from COST 726 action are suggesting a stability of UV which let us assume no change  
168 over time. These estimates were available for European and North African countries, with a resolution  
169 of 5km. However, they are not available for other countries (Americas, Sub-saharian Africa, Asia,  
170 Oceania).

171

#### 172 *Estimation of missing ultraviolet radiation data using global solar radiation*

173 In each location, global ambient annual solar radiation (a measure of solar energy including all  
174 wavelengths) was estimated using astronomical equations and the statistics of sunshine hours, using  
175 the same methodology as in the POLA Study <sup>13</sup>. Overall, global ambient annual solar radiation  
176 estimates were available in 116 locations (101 French departments, 7 European countries, 3 North  
177 African countries, 5 other countries). In 105 locations (95 French departments, 7 European countries,  
178 3 North African countries), both UVR and global solar radiation (GSR) estimates were available.

179 Pearson's correlation coefficient between these two variables was 0.952. For the 11 areas with  
180 missing UVR but available solar radiation (6 French overseas departments and 5 countries), we thus  
181 estimated ambient UVR from global solar radiation, using linear regression modeling. The regression  
182 equation derived from the 105 locations with both UVR and GSR was:  $UVR = 5613.105 + 0.0729 \times$   
183  $GSR$  ( $r^2=0.91$ ). The same analyses were performed for UVA and UVB, leading to the following  
184 equations:  $UVA = 5467.398 + 0.0710 \times GSR$  ;  $UVB = 145.708 + 0.00189 \times GSR$ .

185 Finally, estimates of UVR exposure still could not be estimated for some countries. When, for a given  
186 subject, the number of years spent in such countries was less or equal to 3 years, these countries  
187 were eliminated from the calculations. For 71 subjects, average ambient UVR was therefore calculated  
188 on 62, 63 or 64 years instead of 65. In addition, 113 subjects were excluded from the analyses  
189 because they had spent more than 3 years in countries where UVR could not be estimated.

190

191 **Other variables**

192 The following potential confounders have been selected, based on literature results reporting  
193 significant associations of AMD and/or cataract and/or light exposure with age, sex, educational level,  
194 diabetes, hypertension, asthma, oral corticosteroid use, smoking, physical activity, body mass index  
195 (BMI), plasma glucose and lipids, dietary intake of energy, antioxidants, lutein and zeaxanthin and  
196 omega3 fatty acids<sup>2,3</sup>. We also included the two major genetic polymorphisms associated with AMD  
197 Complement Factor H (CFH) Y402H (rs1061170) and Age-Related Maculopathy Susceptibility 2  
198 (ARMS2) A69S (rs10490924) polymorphisms<sup>27-29</sup>. These genetic polymorphisms were also strongly  
199 associated with AMD in the Alienor study<sup>30,31</sup>. By comparison with subjects homozygous for the most  
200 frequent allele (TT for CFH Y402H and GG for ARMS2 A69S), subjects homozygous for the less  
201 frequent allele (CC for CFH Y402H, TT for ARMS2 A69S) exhibited 5- to 15- fold increased risk for  
202 AMD. Heterozygous subjects (TC for CFH Y402H and GT for ARM2 A69S) exhibit intermediate risks  
203 (2- to 5- fold increased risks). Indeed, all of these factors may represent confounders in the  
204 relationship of AMD with lifetime ambient UVR, since lifestyle (smoking, physical activity, diet), health  
205 conditions and genetic characteristics have been reported to vary according to geographical location,  
206 and thus with lifetime ambient UVR<sup>32-34</sup>.

207 Data were collected during a face-to-face interview using a standardized questionnaire administered  
208 by a trained psychologist or nurse. At baseline (1999-2000), general data included: demographic  
209 characteristics, educational level and smoking. BMI (kg/m<sup>2</sup>) was calculated as weight/height<sup>2</sup> using  
210 weight and height. Physical activity was assessed by 2 questions: "Do you practice sports?" (yes/no)  
211 and "Do you perspire when you practice sports?" (Never/sometimes/most of the time/always). A 3-  
212 level variable was computed to describe intensity of physical activity, as already published.<sup>35</sup> Plasma  
213 glucose and lipids were measured at the Biochemistry Laboratory of the University Hospital of Dijon  
214 from baseline fasting blood samples. Dietary intake of energy, antioxidants, lutein and zeaxanthin and  
215 omega3 fatty acids were estimated from a 24h recall performed through face-to-face interview by  
216 specifically trained dietitians in 2001-2002<sup>35,36</sup>. Genetic polymorphisms were determined by the Lille  
217 Génopôle, from the DNA samples collected at baseline (1999–2001).

218

219

## 220 **Statistical methods**

221 First, associations of ambient solar radiation with socio-demographic factors, lifestyle, biological and  
222 dietary parameters were performed using chi-square test or analysis of variance.

223 Associations of cataract extraction and AMD with ambient UVR were estimated using logistic  
224 Generalized Estimating Equations (GEE) models <sup>37</sup>, which allow taking into account the data from both  
225 eyes and their intra-individual correlations. Adjusted odds-ratios were estimated using cataract  
226 extraction or AMD as the dependent variable, and ambient UVR and potential confounders as the  
227 independent variables. Ambient UVR was divided in three groups (lower quartile, intermediate  
228 quartiles, upper quartile), the medium category being the reference. With regard to AMD, two models  
229 were performed (for early and late AMD, respectively), subjects without any AMD being the reference  
230 in both models. With regard to cataract, subjects without cataract extraction were the reference group.

231 With regard to cataract extraction, potential confounders retained in the analysis were known risk  
232 factors for cataract <sup>2</sup> (age, gender, smoking, diabetes, oral corticosteroid use and asthma) and factors  
233 significantly associated with ambient UVR (at  $p < 0.05$ ). Similarly, with regard to AMD, potential  
234 confounders retained in the analysis were well known risk factors, which are strongly associated with  
235 AMD in our cohort (age, gender, smoking <sup>30</sup>, CFH <sup>30</sup> and ARMS2 <sup>31</sup> polymorphisms, dietary intake of  
236 omega3 fatty acids <sup>36</sup>) and factors significantly associated with ambient UVR (at  $p < 0.05$ ). No  
237 collinearity was detected between the variables included in the final models. No major confounding  
238 was detected (variation by more than 10 % of the estimates of the odds-ratios when deleting one  
239 confounder from the models). All statistical analyses were performed using statistical software (SAS,  
240 version 9.1; SAS Institute Inc., Cary, NC).

241

## 242 **RESULTS**

243 Figure 1 presents the distribution of annual ambient UVR exposures (total UV, UVA and UVB),  
244 estimated from residential history. These variables showed important inter-individual variability,  
245 ranging respectively, from 32.39 kJ/cm<sup>2</sup> to 50.93 kJ/cm<sup>2</sup>, 31.55 kJ/cm<sup>2</sup> to 49.61 kJ/cm<sup>2</sup> and 0.81  
246 kJ/cm<sup>2</sup> to 1.32 kJ/cm<sup>2</sup>. The central peaks correspond to subjects who spent all their life in the

247 Bordeaux area. As shown in Table 1 for total UVR, these subjects constitute the middle group  
248 (intermediate quartiles), who were born and lived all their life in the South of France. By contrast, two  
249 thirds of subjects from the lower quartile of ambient solar radiation were born in Central or Northern  
250 France, where they spent about 23 years on average. Among subjects from the upper quartile, more  
251 than a quarter were born in Northern Africa and remained there for 32 years on average  
252 (corresponding to the independence of these countries in the 1960s) and 8.13 % were born in  
253 Southern Europe. Although the definition of the intermediate quartiles is narrow (39.649-40.173  
254 kJ/cm<sup>2</sup>), there is significant variability, with a mean of 38.375 kJ/cm<sup>2</sup> in the lower quartile, compared  
255 with 42.718 kJ/cm<sup>2</sup> in the upper quartile. Similar findings were found for UVA and UVB (data not  
256 shown).

257 Socio-demographic, lifestyle and biological characteristics of participants appear different according to  
258 their lifetime solar exposure. As shown in Table 2, compared with subjects from the middle quartiles,  
259 subjects in the higher quartile were more often males (42.58 % versus 31.74 %) and tended to have a  
260 higher frequency of the TC genotype of CFH Y402H (47.12 % versus 37.43 %) and lower frequency of  
261 TT genotype of CFH Y402H (45.03 % versus 49.21 %), while subjects from the lower quartiles had a  
262 higher educational level (48.80 % with University degree versus 26.97 %) and tended to have a higher  
263 frequency of the TC genotype of CFH Y402H (47.15 % versus 37.43 %) and lower frequency of TT  
264 genotype of CFH Y402H (40.93 % versus 49.21%). As shown in Table 3, both subjects from the upper  
265 and lower quartiles had more smoked during their life (18.36 % and 22.93 % having smoked more  
266 than 20 pack-years respectively), by comparison with intermediate exposures (12.50 %). Subjects in  
267 the upper quartile also had higher total energy intake (1835.7 Kcal/day versus 1678.2 Kcal/day).  
268 These variables (education, smoking, CFH Y402H genotypes and total energy intakes) were therefore  
269 considered as potential confounders in the analyses of association of solar radiation with cataract  
270 extraction and AMD.

271 Cataract extraction status was available in at least one eye of 958 subjects (99.5%). Of those, 837 had  
272 complete data for light exposure variables. In addition, 143 subjects had missing values in potential  
273 confounders, leaving 694 subjects in the fully adjusted models, representing 1388 eyes among which  
274 542 had undergone cataract surgery. Participants with missing data were not significantly different  
275 from those without missing data, with regard to age, gender, ambient UVR or genetic polymorphisms  
276 (data not shown). As shown in Table 4, after adjustment for confounders, by comparison with

277 participants exposed to medium ambient total UVR, participants exposed to high ambient total UVR  
278 were at increased risk for cataract extraction (OR=1.53, 95 % CI: 1.04-2.26, p=0.03), while participants  
279 with low exposure had similar risk as those with medium exposures (0.75 (0.51-1.12)). Similar results  
280 were observed for UVA and UVB. Among the potential confounders included in the multivariate  
281 models, those significantly associated with cataract surgery were age (OR=1.18 for 1-year increase,  
282 95 % CI: 1.13-1.22, p<0.0001), gender (OR=1.86 for females versus males, 95 % CI: 1.23-2.80,  
283 p=0.003), having smoked 20 pack-years or more (OR= 2.23, 95 % CI: 1.37-3.64, p=0.001), history of  
284 asthma (OR=1.99 1.15-3.42, p=0.01).

285 AMD status was available in at least one eye of 875 subjects (91 %). Of those, 769 had complete data  
286 for light exposure variables. In addition, 172 subjects had missing values in potential confounders,  
287 leaving 597 subjects in the fully adjusted models, representing 1154 eyes among which 238 with early  
288 AMD and 49 with late AMD. Participants with missing data were not significantly different from those  
289 without missing data, with regard to age, gender, ambient UVR or genetic polymorphisms (data not  
290 shown). Participants exposed to high ambient total UVR tended to be at increased risk for early AMD  
291 (OR=1.59, 95 % CI: 1.04-2.44, p=0.03), by comparison with participants with medium exposures  
292 (Table 5). In addition, participants exposed to low ambient solar radiation were increased risk for early  
293 AMD (OR=1.69, 95 %CI: 1.06-2.69, p=0.03), by comparison with medium exposures. By contrast, no  
294 statistically significant associations of late AMD with UVR exposures were found. Overall, results were  
295 very similar for total UV, UVA and UVB exposures. Among the potential confounders included in the  
296 multivariate models, those significantly associated with early AMD were age (OR=1.06 for 1-year  
297 increase, 95% CI: 1.02-1.10, p=0.008), gender (OR=1.95 for females versus males, 95% CI: 1.19-  
298 3.21, p=0.01), ARMS2 A69S GT genotype (OR=1.62, 95 % CI: 1.12-2.35, p=0.01), ARMS A69S TT  
299 genotype (OR=12.09, 95 % CI: 4.63-31.55, p<0.0001) and CFH Y402H CC genotype (OR=1.89, 95%  
300 CI: 1.13-3.16, p=0.01). The factors significantly associated with late AMD were age (OR=1.24, 95%  
301 CI: 1.14-1.35, p<0.0001), gender (OR=2.93, 95 % CI: 1.16-7.43, p=0.02), ARMS A69S TT genotype  
302 (OR=59.71, 95 % CI: 15.53-229.57, p<0.0001), CFH Y402H TC genotype (OR=3.49, 95% CI: 1.31-  
303 9.30, p=0.009), CFH Y402H CC genotype (OR=6.68, 95 % CI: 2.05-21.77, p=0.001), dietary omega3  
304 fatty acids intake (OR=0.63, 95% CI: 0.41-0.96, p=0.03) and secondary education (OR=7.46, 95 % CI:  
305 1.63-34.05, p=0.01). We detected no significant interactions of genetic polymorphisms with UVR  
306 exposure.

307

308 **DISCUSSION**

309 While confirming the well-known association of cataract with high ambient UVR,<sup>7-13</sup> the present study  
310 suggests a U-shaped association of early AMD with ambient UVR, with increased risk both in low- and  
311 high-UVR exposures. The effects of UVR on human health generally follows such a U-shaped (or J-  
312 shaped) relationship<sup>38</sup>. Indeed, on one hand, high exposures have well documented effects on skin  
313 diseases (skin cancers, sunburns and chronic sun damage...) and several eye diseases (acute  
314 photokeratitis and conjunctivitis, acute solar retinopathy, pterygium, cortical cataract...) <sup>38</sup>. On the other  
315 hand, cutaneous UVR exposure is necessary for endogenous production of vitamin D, so that low  
316 UVR exposures are associated with an increased risk for osteo-muscular diseases and possibly of  
317 other diseases related to vitamin D deficiency (in particular several cancers, cardiovascular disease,  
318 autoimmune diseases) <sup>38-40</sup>. Indeed, besides its effects on bone health, vitamin D has many cellular  
319 effects, including regulation of cellular differentiation, proliferation, apoptosis and angiogenesis <sup>40</sup>, and  
320 anti-inflammatory properties <sup>41</sup>. Many of these processes are implicated in the physiopathology of  
321 AMD and some studies have suggested that low vitamin D status may be associated with an  
322 increased risk for AMD <sup>42-45</sup>. Whether vitamin D deficiency may explain the higher risk for AMD  
323 observed with low sunlight exposure in the present study will need to be determined in future studies.

324

325 Only few epidemiological studies have addressed the potential link of AMD with UVR exposure, with  
326 inconsistent results. This is probably due to major methodological difficulties in the assessment of  
327 UVR exposure. Indeed, UVR exposure and its health effects result from distal factors (such as latitude,  
328 season, cloud cover, stratospheric ozone levels and lower atmospheric pollution), conditioning  
329 ambient UVR, in addition to proximal factors (in particular occupational and leisure exposures to  
330 sunlight, use of hats and sunglasses, skin pigmentation and sun sensitivity...) <sup>46</sup> conditioning individual  
331 exposure and response to exposure. Moreover, previous studies have shown that health effects of  
332 UVR exposure are cumulative over the lifetime. Thus, estimation of lifetime ocular UVR exposure  
333 requires detailed questionnaires on lifetime distal and proximal factors, combined with complex  
334 modeling of their effects on ocular UVR exposure. Since most epidemiological studies in the field of  
335 AMD, including the present study, addressed not only the effect of sunlight exposure, but also of many

336 other factors (smoking, cardiovascular risk factors, genetics, nutrition...), only partial information on  
337 distal and/or proximal factors was collected, limiting the validity of the estimates of UVR exposure, and  
338 the comparability of studies.

339 The Maryland Watermen Study included detailed questionnaires of ocular exposures, combined with  
340 field and laboratory data, in order to assess ocular UVR exposure. In this study, ocular UVR exposure  
341 was not significantly associated with prevalent AMD<sup>47</sup>. In the Beaver Dam Eye Study, ambient UV-B  
342 exposure, estimated from residential history, was not significantly associated with prevalent<sup>15</sup> or  
343 incident<sup>16, 17</sup> AMD, but subjects having spent more than 5 hours/day outside during summer in their  
344 youth were more likely to have prevalent<sup>15</sup> and incident AMD<sup>16, 17</sup>. In the POLA Study<sup>22</sup>, performed in  
345 the South of France, the risk for prevalent early AMD was higher in subjects exposed to low ambient  
346 solar radiation. In the Blue Mountains Eye Study, prevalent AMD was associated both with high and  
347 low sun sensitivity index, by comparison with medium sun sensitivity index<sup>48</sup>, although this was not  
348 confirmed in the prospective analysis of the same cohort<sup>20</sup>. Finally, an Australian case-control study,  
349 AMD was associated with poorer tanning ability and lower time of ocular sunlight exposure<sup>21</sup>.

350 In the present study, only estimates of ambient UVR exposure were available, based on residential  
351 history and satellite-based estimates of UVR, while no data were available on sun-related behaviors,  
352 skin pigmentation or sun sensitivity. This may have led to significant misclassification of exposures,  
353 since actual ocular UVR exposure may be quite different from ambient UVR exposure in those  
354 subjects with very low time spent outdoors. However, it seems most likely that misclassification was  
355 unrelated to ambient UVR (i.e. that there would be similar proportions of people with low/high outdoors  
356 activities in areas with different ambient UVR), and was thus unlikely to have biased the associations  
357 of eye diseases with UVR exposure. Moreover, the well known association of UVR exposure with  
358 cataract was confirmed in the present study, suggesting that our measure of UVR exposure had some  
359 validity.

360 While the lack of information on proximal factors is clearly a limitation, the use of satellite-based  
361 estimates of UVR is a strength. Indeed, most previous studies relied on formulas, estimating UVR  
362 mainly from latitude and altitude. While these are major determinants of UVR, weather conditions also  
363 contribute to actual ground UVR, in particular because of attenuation by cloud cover. UVR estimations  
364 in the present study rely on Meteosat satellite measurements of total solar irradiance by the earth's

365 surface by reflection, over thirty years.

366 Using detailed estimation of ocular lifetime exposure, the Maryland Watermen study and the EUREYE  
367 study have suggested that AMD risk may be associated with blue light exposure<sup>14,49</sup>, which is also  
368 supported by animal and laboratory studies<sup>6,50</sup>. In the present study, we could only estimate ambient  
369 UVR, since no data on blue light are currently available in the Eurosun project. It is however difficult to  
370 distinguish the effect of the different wavelengths, since they are naturally highly correlated. For  
371 instance, in the present study, global solar radiation (a measure of solar energy including all  
372 wavelengths) showed a correlation coefficient of 0.95 with UV radiation, measured in 105 geographical  
373 areas.

374 The present study has several limitations. One limitation of our study could come from the  
375 representativeness of the sample. The Alienor subsample tends to over-represent younger subjects  
376 and high socioeconomic status, among subjects participating to the 3C Study<sup>23</sup>. The individuals  
377 included in this study may therefore be healthier and have different lifestyles, particularly concerning  
378 sunlight exposure, than the general population. These differences may have affected the distribution  
379 of sunlight exposure or the prevalence of eye diseases. However, subjects participating in the 3C  
380 study and included in the Alienor study were not different from those who were not included for most  
381 parameters of interest in our study<sup>23</sup>. Furthermore, as described previously<sup>23</sup>, the age-gender-  
382 specific prevalence rates of AMD in the Alienor study were similar to that observed in other studies  
383 performed in Europe<sup>51,52</sup> and other industrialized countries<sup>53</sup>. Data collection was performed in the  
384 same way in all individuals regardless of their AMD stage and photograph graders had no access to  
385 sunlight exposure data. Therefore, we can assume that the error was not differential and was unlikely  
386 to have biased the estimation of any of the associations of AMD with ambient solar radiation.

387 In observational studies, confounding is always a concern. We therefore adjusted for potential  
388 confounders, including all known risk factors for cataract and AMD. However, we cannot totally  
389 exclude residual confounding. In the present study, it is in particular striking that not only socio-  
390 demographic and lifestyle variables varied with ambient UVR exposure, but also genetic background  
391 (mainly CFH Y402H). While there are major differences in distribution of CFH Y402H alleles among  
392 ethnic groups (with low frequency of the at-risk C allele in non-Europeans<sup>54,55</sup>), to our knowledge,  
393 there are no available data on its geographical distribution within the European continent (where most

394 of the Alienor participants were born). However, large variations of allele frequencies between  
395 European countries have been reported, in particular for Apolipoprotein E<sup>33</sup>. Since AMD has a strong  
396 genetic component, and there may be geographical variation in the distribution of gene  
397 polymorphisms, gene polymorphisms may be confounders in the relationship of AMD with any factor  
398 affected by geographical variation (including UVR exposure, but also lifestyle, diet...). This underlines  
399 the importance of taking into account both environmental and genetic confounders in studies of  
400 associations of eye diseases with sunlight exposure. None of the previously published studies took  
401 potential genetic confounders into account. In the present study, although we took into account the  
402 two major genetic risk factors for AMD (CFH Y402H and ARMS A69S), we cannot exclude that these  
403 groups of subjects differ for other genetic factors related to AMD (complement factors CF1, C2, C3  
404 and CFB, LIPC, CETP...<sup>3</sup>).

405 Since this study was cross-sectional, recall bias may have affected the results. However, only few  
406 prospective studies are available in this field. Our study included only a small number of cases of late  
407 AMD, which induced low statistical power for detecting associations with ambient solar radiation and of  
408 interactions of genetic polymorphisms with UVR exposure. Finally, cataract status was determined  
409 only on the basis of cataract surgery, rather than on the presence of lens opacities, which were not  
410 available in our study because of lack of pupil dilation. This may have caused misclassification of  
411 cataract status and therefore may have biased the estimates.

412 In conclusion, our study confirms the high risk for cataract in subjects exposed to high ambient solar  
413 radiation. It also suggests a U-shaped association of early AMD with UVR exposure, with an increased  
414 risk for both low- and high-exposures. This will need to be confirmed in future, possibly prospective,  
415 studies. The reasons for a potentially increased risk for early AMD with low solar radiation remain  
416 unclear and need to be further studied.

417

#### 418 References:

- 419 1. Resnikoff S, Pascolini D, Etya'ale D, et al. Global data on visual impairment in the year 2002. *Bull*  
420 *World Health Organ.* 2004; 82: 844-851.
- 421 2. Asbell PA, Dualan I, Mindel J, Brocks D, Ahmad M and Epstein S. Age-related cataract. *Lancet.*  
422 2005; 365: 599-609.

423 3. Lim LS, Mitchell P, Seddon JM, Holz FG and Wong TY. Age-related macular degeneration. *Lancet*.  
424 2012; 379: 1728-1738.

425 4. Sui GY, Liu GC, Liu GY, et al. Is sunlight exposure a risk factor for age-related macular  
426 degeneration? A systematic review and meta-analysis. *Br J Ophthalmol*. 2013; 97: 389-394.

427 5. McCarty CA and Taylor HR. A review of the epidemiologic evidence linking ultraviolet radiation and  
428 cataracts. *Dev. Ophthalmol*. 2002; 35: 21-31.

429 6. Wu J, Seregard S and Algvere PV. Photochemical damage of the retina. *Surv Ophthalmol*. 2006; 51:  
430 461-481.

431 7. Brilliant LB, Grasset NC, Pokhrel RP, et al. Associations among cataract prevalence, sunlight hours,  
432 and altitude in the Himalayas. *Am. J. Epidemiol*. 1983; 118: 250-264.

433 8. Hiller R, Sperduto RD and Ederer F. Epidemiologic associations with cataract in the 1971-1972  
434 National Health and Nutrition Examination Survey. *Am. J. Epidemiol*. 1983; 118: 239-249.

435 9. Taylor HR, West SK, Rosenthal FS, et al. Effect of ultraviolet radiation on cataract formation. *N*.  
436 *Engl. J. Med*. 1988; 319: 1429-1433.

437 10. Rosmini F, Stazi MA, Milton RC, Sperduto RD, Pasquini P and Maraini G. A dose-response effect  
438 between a sunlight index and age-related cataracts. Italian-American Cataract Study Group. *Ann*.  
439 *Epidemiol*. 1994; 4: 266-270.

440 11. West SK, Duncan DD, Munoz B, et al. Sunlight exposure and risk of lens opacities in a population-  
441 based study: The Salisbury Eye Evaluation Project. *Jama*. 1998; 280: 714-718.

442 12. Sreenivas V, Prabhakar AK, Badrinath SS, et al. A rural population based case-control study of  
443 senile cataract in India. *J. Epidemiol*. 1999; 9: 327-336.

444 13. Delcourt C, Carriere I, Ponton Sanchez A, et al. Light exposure and the risk of cortical, nuclear, and  
445 posterior subcapsular cataracts: the Pathologies Oculaires Liées à l'Age (POLA) Study. *Arch*.  
446 *Ophthalmol*. 2000; 118: 385-392.

447 14. Taylor HR, Munoz B, West S, Bressler NM, Bressler SB and Rosenthal FS. Visible light and risk of  
448 age-related macular degeneration. *Trans. Am. Ophthalmol. Soc*. 1990; 88: 163-173.

449 15. Cruickshanks KJ, Klein R and Klein BE. Sunlight and age-related macular degeneration. The Beaver  
450 Dam Eye Study. *Arch. Ophthalmol*. 1993; 111: 514-518.

451 16. Cruickshanks KJ, Klein R, Klein BE and Nondahl DM. Sunlight and the 5-year incidence of early  
452 age-related maculopathy: the beaver dam eye study. *Arch. Ophthalmol*. 2001; 119: 246-250.

453 17. Tomany SC, Cruickshanks KJ, Klein R, Klein BE and Knudtson MD. Sunlight and the 10-year  
454 incidence of age-related maculopathy: the Beaver Dam Eye Study. *Arch. Ophthalmol*. 2004; 122: 750-  
455 757.

456 18. Group EDC-cS. Risk factors for neovascular age-related macular degeneration. The Eye Disease  
457 Case-Control Study Group. *Arch. Ophthalmol*. 1992; 110: 1701-1708.

458 19. Khan JC, Shahid H, Thurlby DA, et al. Age related macular degeneration and sun exposure, iris  
459 colour, and skin sensitivity to sunlight. *Br J Ophthalmol*. 2006; 90: 29-32.

460 20. Pham TQ, Rochtchina E, Mitchell P, Smith W and Wang JJ. Sunlight-related factors and the 10-  
461 year incidence of age-related maculopathy. *Ophthalmic Epidemiol*. 2009; 16: 136-141.

462 21. Darzins P, Mitchell P and Heller RF. Sun exposure and age-related macular degeneration. An  
463 Australian case-control study. *Ophthalmology*. 1997; 104: 770-776.

464 22. Delcourt C, Carriere I, Ponton Sanchez A, Fourrey S, Lacroux A and Papoz L. Light exposure and  
465 the risk of age-related macular degeneration: the Pathologies Oculaires Liees a l'Age (POLA) study.  
466 *Arch. Ophthalmol*. 2001; 119: 1463-1468.

467 23. Delcourt C, Korobelnik JF, Barberger-Gateau P, et al. Nutrition and Age-Related Eye Diseases: The  
468 ALIENOR (Antioxydants, Lipides Essentiels, Nutrition et Maladies Oculaires) Study. *J Nutr Health*  
469 *Aging*. 2010; 14: 854-861.

470 24. Group CS. Vascular factors and risk of dementia: design of the Three-City Study and baseline  
471 characteristics of the study population. *Neuroepidemiology*. 2003; 22: 316-325.

472 25. Bird AC, Bressler NM, Bressler SB, et al. An international classification and grading system for age-  
473 related maculopathy and age-related macular degeneration. The International ARM Epidemiological  
474 Study Group. *Surv Ophthalmol*. 1995; 39: 367-374.

475 26. Klein R, Klein BE, Knudtson MD, et al. Prevalence of age-related macular degeneration in 4  
476 racial/ethnic groups in the multi-ethnic study of atherosclerosis. *Ophthalmology*. 2006; 113: 373-380.  
477 27. Chakravarthy U, McKay GJ, de Jong PT, et al. ARMS2 increases the risk of early and late age-  
478 related macular degeneration in the European Eye Study. *Ophthalmology*. 2013; 120: 342-348.  
479 28. Despret DD, Klaver CC, Witteman JC, et al. Complement factor H polymorphism, complement  
480 activators, and risk of age-related macular degeneration. *Jama*. 2006; 296: 301-309.  
481 29. Seddon JM, Gensler G and Rosner B. C-reactive protein and CFH, ARMS2/HTRA1 gene variants are  
482 independently associated with risk of macular degeneration. *Ophthalmology*. 2010; 117: 1560-1566.  
483 30. Delcourt C, Delyfer MN, Rougier MB, et al. Associations of Complement Factor H and Smoking  
484 with Early Age-Related Macular Degeneration: The ALIENOR Study. *Invest Ophthalmol Vis Sci*. 2011;  
485 52: 5955-5962.  
486 31. Delcourt C, Delyfer MN, Rougier MB, et al. ARMS2 A69S Polymorphism and the Risk for Age-  
487 Related Maculopathy: The ALIENOR Study. *Arch Ophthalmol*. 2012; 130: 1077-1078.  
488 32. Czernichow S, Bruckert E, Oppert JM, et al. Intake of added oils and fats among middle-aged  
489 French adults: relationships with educational level and region of residence. *J Am Diet Assoc*. 2005;  
490 105: 1889-1894.  
491 33. Ewbank DC. The APOE gene and differences in life expectancy in Europe. *J Gerontol A Biol Sci Med*  
492 *Sci*. 2004; 59: 16-20.  
493 34. Al Delaimy WK, Van Kappel AL, Ferrari P, et al. Plasma levels of six carotenoids in nine European  
494 countries: report from the European Prospective Investigation into Cancer and Nutrition (EPIC).  
495 *Public. Health. Nutr*. 2004; 7: 713-722.  
496 35. Feart C, Jutand MA, Larrieu S, et al. Energy, macronutrient and fatty acid intake of French elderly  
497 community dwellers and association with socio-demographic characteristics: data from the Bordeaux  
498 sample of the Three-City Study. *Br J Nutr*. 2007; 98: 1046-1057.  
499 36. Merle B, Delyfer MN, Korobelnik JF, et al. Dietary omega-3 Fatty acids and the risk for age-related  
500 maculopathy: the alienor study. *Invest Ophthalmol Vis Sci*. 2011; 52: 6004-6011.  
501 37. Zeger SL, Liang KY and Albert PS. Models for longitudinal data: a generalized estimating equation  
502 approach. *Biometrics*. 1988; 44: 1049-1060.  
503 38. Lucas RM, McMichael AJ, Armstrong BK and Smith WT. Estimating the global disease burden due  
504 to ultraviolet radiation exposure. *Int J Epidemiol*. 2008; 37: 654-667.  
505 39. Chowdhury R, Kunutsor S, Vitezova A, et al. Vitamin D and risk of cause specific death: systematic  
506 review and meta-analysis of observational cohort and randomised intervention studies. *BMJ*. 2014;  
507 348: g1903.  
508 40. Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007; 357: 266-281.  
509 41. Mora JR, Iwata M and von Andrian UH. Vitamin effects on the immune system: vitamins A and D  
510 take centre stage. *Nat Rev Immunol*. 2008; 8: 685-698.  
511 42. Parekh N, Chappell RJ, Millen AE, Albert DM and Mares JA. Association between vitamin D and  
512 age-related macular degeneration in the Third National Health and Nutrition Examination Survey,  
513 1988 through 1994. *Arch Ophthalmol*. 2007; 125: 661-669.  
514 43. Millen AE, Voland R, Sondel SA, et al. Vitamin D status and early age-related macular  
515 degeneration in postmenopausal women. *Arch Ophthalmol*. 2011; 129: 481-489.  
516 44. Morrison MA, Silveira AC, Huynh N, et al. Systems biology-based analysis implicates a novel role  
517 for vitamin D metabolism in the pathogenesis of age-related macular degeneration. *Hum Genomics*.  
518 2011; 5: 538-568.  
519 45. Seddon JM, Reynolds R, Shah HR and Rosner B. Smoking, dietary betaine, methionine, and  
520 vitamin d in monozygotic twins with discordant macular degeneration: epigenetic implications.  
521 *Ophthalmology*. 2011; 118: 1386-1394.  
522 46. Lucas R, McMichael T, Smith W and Armstrong B. Solar Ultraviolet Radiation - Global burden of  
523 disease from solar ultraviolet radiation. Geneva. WHO, 2006.  
524 47. West SK, Rosenthal FS, Bressler NM, et al. Exposure to sunlight and other risk factors for age-  
525 related macular degeneration see comments. *Arch. Ophthalmol*. 1989; 107: 875-879.

- 526 48. Mitchell P, Smith W and Wang JJ. Iris color, skin sun sensitivity, and age-related maculopathy: The  
527 Blue Mountains Eye Study. *Ophthalmology*. 1998; 105: 1359-1363.
- 528 49. Fletcher AE, Bentham GC, Agnew M, et al. Sunlight exposure, antioxidants, and age-related  
529 macular degeneration. *Arch Ophthalmol*. 2008; 126: 1396-1403.
- 530 50. Sparrow JR and Boulton M. RPE lipofuscin and its role in retinal pathobiology. *Exp Eye Res*. 2005;  
531 80: 595-606.
- 532 51. Augood CA, Vingerling JR, de Jong PT, et al. Prevalence of age-related maculopathy in older  
533 Europeans: the European Eye Study (EUREYE). *Arch Ophthalmol*. 2006; 124: 529-535.
- 534 52. Vingerling JR, Dielemans I, Hofman A, et al. The prevalence of age-related maculopathy in the  
535 Rotterdam Study. *Ophthalmology*. 1995; 102: 205-210.
- 536 53. Friedman DS, O'Colmain BJ, Munoz B, et al. Prevalence of age-related macular degeneration in  
537 the United States. *Arch Ophthalmol*. 2004; 122: 564-572.
- 538 54. Kim NR, Kang JH, Kwon OW, Lee SJ, Oh JH and Chin HS. Association between complement factor  
539 H gene polymorphisms and neovascular age-related macular degeneration in Koreans. *Invest*  
540 *Ophthalmol Vis Sci*. 2008; 49: 2071-2076.
- 541 55. Sofat R, Casas JP, Webster AR, et al. Complement factor H genetic variant and age-related  
542 macular degeneration: effect size, modifiers and relationship to disease subtype. *Int J Epidemiol*.  
543 2012; 41: 250-262.
- 544
- 545

Table 1. Light exposure-related characteristics of participants according to lifetime ambient total UVR exposure of the Alienor Study (Bordeaux, France, 2006-2008) (N=837)

	Total UVR, kJ/cm <sup>2</sup>			p-value
	≤ 39.649 (N=209)	]39.649-40.173[ (N=419)	≥40.173 (N=209)	
Mean total UVR, kJ/cm <sup>2</sup> (sd)	38.375 (1.26)	39.887 (0.08)	42.718 (2.29)	
Place of birth (n (%)) (N=837)				
South of France	66 (31.58)	392 (93.56)	102 (48.80)	P<0.0001
Central France	52 (24.88)	9 (2.15)	14 (6.70)	
Northern France	88 (42.11)	12 (2.86)	14 (6.70)	
Southern Europe	0 (0.00)	4 (0.95)	17 (8.13)	
Northern Africa	0 (0.00)	1 (0.24)	57 (27.27)	
Other	3 (1.44)	1 (0.24)	5 (2.39)	
Time spent at place of birth, years (median (range)) (N=837)				
South of France	68 (39-89)	79 (55-94)	74 (35-92)	P<0.0001
Central France	22 (2-58)	13 (1-34)	20 (4-37)	P=0.11
Northern France	23.5 (3-69)	3.5 (1-42)	12.5 (1-31)	0.001
Southern Europe		2.5 (1-6)	17 (2-31)	0.03
Northern Africa		1 (1-1)	32 (17-48)	P<0.0001
Other	3 (1-6)	3 (3-3)	18 (1-30)	0.13

Table 2. Socio-demographic, medical and genetic characteristics of participants according to lifetime ambient total UVR exposure of the Alienor Study (Bordeaux, France, 2006-2008) (N=837)

	Ambient total UVR, kJ/cm <sup>2</sup>			p-value
	≤ 39.649 (N=209)	]39.649- 40.173[ (N=419)	≥40.173 (N=209)	
Age, years (mean (sd)) (N=837)	80.50 (4.42)	80.21 (4.49)	79.96 (4.44)	0.46
Male gender (n (%)) (N=837)	87 (41.63)	133 (31.74)	89 (42.58)	0.008
Education (n (%)) (N=837)				
None or primary school	48 (22.97)	130 (31.00)	54 (25.84)	P<0.0001
Secondary school	34 (16.27)	137 (32.70)	63 (30.14)	(global)
High school	25 (11.96)	39 (9.31)	23 (11.00)	
University	102 (48.80)	113 (26.97)	69 (33.01)	
Diabetes (n (%)) (N=763)	23 (12.17)	53 (13.80)	21 (11.05)	0.63
Hypertension (n (%)) (N=837)	161 (77.03)	318 (75.89)	152 (72.73)	0.56
Asthma (n (%)) (N=828)	21 (10.14)	36 (8.67)	14 (6.80)	0.47
Oral corticosteroid use (n (%)) (N=837)	3 (1.44)	2 (0.48)	4 (1.91)	0.22
CFH Y402 H (n (%)) (N=766)				
TT (low AMD risk)	79 (40.93)	188 (49.21)	86 (45.03)	0.05
TC (intermediate AMD risk)	91 (47.15)	143 (37.43)	90 (47.12)	(global)
CC (high AMD risk)	23 (11.92)	51 (13.35)	15 (7.85)	
ARMS2 A69S (n (%)) (N=703)				
GG (low AMD risk)	118 (66.67)	232 (66.48)	112 (63.28)	0.91
GT (intermediate AMD risk)	53 (29.94)	103 (29.51)	59 (33.33)	(global)
TT (high AMD risk)	6 (3.39)	14 (4.01)	6 (3.39)	

Table 3. Lifestyle and nutritional characteristics of participants according to lifetime ambient total UVR exposure of the Alienor Study (Bordeaux, France, 2006-2008) (N=837)

	Ambient total UVR, kJ/cm <sup>2</sup>			p-value
	≤ 39.649 (N=209)	]39.649- 40.173[ (N=419)	≥40.173 (N=209)	
<b>Smoking (n (%)) (N=828)</b>				
Non smoker	111 (54.15)	296 (71.15)	138 (66.67)	0.0005
< 20 pack-years	47 (22.93)	68 (16.35)	31 (14.98)	
≥ 20 pack-years	47 (22.93)	52 (12.50)	38 (18.36)	
<b>Physical activity (n (%)) (N=837)</b>				
Low	29 (13.88)	83 (19.81)	31 (14.83)	0.20
Medium	112 (53.59)	226 (53.94)	108 (51.67)	
High	47 (22.49)	76 (18.14)	43 (20.57)	
No answer	21 (10.05)	34 (8.11)	27 (12.92)	
Body mass index, kg/m <sup>2</sup> (mean (sd)) (N=829)	25.90 (4.00)	26.17 (3.97)	26.61 (3.65)	0.17
<b>Fasting plasma measurements, mmol/l (mean (sd)) :</b>				
Glucose (N=759)	5.16 (1.03)	5.13 (1.12)	5.13 (1.22)	0.96
Total cholesterol (N=787)	5.83 (0.91)	5.83 (1.03)	5.72 (0.92)	0.37
HDL- cholesterol (N=786)	1.60 (0.40)	1.61 (0.39)	1.57 (0.39)	0.66
LDL- cholesterol (N=785)	3.66 (0.80)	3.67 (0.91)	3.61 (0.79)	0.73
Triglycerides (N=786)	1.25 (0.61)	1.23 (0.60)	1.18 (0.48)	0.37
<b>Dietary intake (mean (sd)):</b>				
Total energy Kcal/day (N=797)	1835.7 (574.1)	1678.2 (513.4)	1704.5 (535.3)	0.003
Vitamin C, mg/day (N=797)	89.62 (56.98)	81.95 (60.96)	92.10 (82.66)	0.15
Vitamin E, mg/day (N=797)	6.72 (4.72)	6.30 (4.51)	6.70 (4.43)	0.45
Lutein and zeaxanthin, mg/day (N=781)	0.60 (0.92)	0.55 (0.91)	0.71 (0.90)	0.14
Omega 3 fatty acids, mg/day (N=797)	0.48 (1.20)	0.43 (1.00)	0.41 (1.16)	0.79

Table 4. Associations of cataract extraction with lifetime ambient UVR exposure in the Alienor Study (Bordeaux, France, 2006-2008) (Odds-ratios (OR) and 95 % confidence intervals (CI))

	Cataract surgery				Odds-ratio*	P
	No		Yes			
	(N=846 eyes)		(N=542 eyes)			
	N	%	N	%		
Total UV (kJ/cm <sup>2</sup> )						
≤39.649 (n=350 eyes)	230	65.71	120	34.29	0.75 (0.51-1.12)	0.16
]39.649-40.173[ (n=696 eyes)	430	61.78	266	38.22	1.00 (ref)	
≥40.173 (n=342 eyes)	186	54.39	156	45.61	1.53 (1.04-2.26)	0.03
UVA (kJ/cm <sup>2</sup> )						
≤38.617 (n=350 eyes)	230	65.71	120	34.29	0.75 (0.51-1.12)	0.16
]38.617-39.131[ (n=696 eyes)	430	61.78	266	38.22	1.00 (ref)	
≥39.131 (n=342 eyes)	186	54.39	156	45.61	1.53 (1.04-2.26)	0.03
UVB (kJ/cm <sup>2</sup> )						
≤1.028 (n=350 eyes)	230	65.71	120	34.29	0.75 (0.51-1.10)	0.14
]1.028 -1.042[ (n=696 eyes)	430	61.78	266	38.22	1.00 (ref)	
≥1.042 (n=342 eyes)	186	54.39	156	45.61	1.53 (1.04-2.26)	0.03


\* estimated using multivariate Generalized Estimating Equations (GEE) logistic regression models, adjusted for age, gender, educational level, smoking, diabetes, oral corticosteroid, asthma, total energy intake and CFH Y402H.

Table 5. Associations of AMD with lifetime ambient UVR exposure in the Alienor Study (Bordeaux, France, 2006-2008) (Odds-ratios (OR) and 95 % confidence intervals (CI))


	<b>Early AMD</b>	<b>P</b>	<b>Late AMD</b>	<b>P</b>
	<b>OR (CI)*</b>		<b>OR (CI)*</b>	
	<b>(N=238 eyes)</b>		<b>(N=49 eyes)</b>	
<b>Total UV (kJ/cm<sup>2</sup>)</b>				
≤39.649 (n=300 eyes)	1.69 (1.06-2.69)	0.03	1.03 (0.33-3.26)	0.95
]39.649-40,173[ (n=559 eyes)	1.00 (ref)		1.00 (ref)	
≥40.173 (n=295 eyes)	1.59 (1.04-2.44)	0.03	1.11 (0.36-3.36)	0.86
<b>UVA (kJ/cm<sup>2</sup>)</b>				
≤38.617 (n=300 eyes)	1.69 (1.06-2.69)	0.03	1.03 (0.33-3.26)	0.95
]38.617-39.131[ (n=559 eyes)	1.00 (ref)		1.00 (ref)	
≥39.131 (n=295 eyes)	1.59 (1.04-2.44)	0.03	1.11 (0.36-3.36)	0.86
<b>UVB (kJ/cm<sup>2</sup>)</b>				
≤1.028 (n=302 eyes)	1.66 (1.04-2.64)	0.03	1.03 (0.33-3.25)	0.96
]1.028 -1.042[ (n=557 eyes)	1.00 (ref)		1.00 (ref)	
≥1.042 (n=295 eyes)	1.58 (1.03-2.42)	0.04	1.11 (0.36-3.36)	0.86

\* estimated using multivariate Generalized Estimating Equations (GEE) logistic regression models, age, gender, educational level, smoking, CFH Y402H and ARMS2 A69S polymorphisms, cataract extraction, dietary intake of total energy and of omega3 fatty acids.

## Total UV



## UVA



## UVB




Figure 1. Distribution of lifetime ambient total UV, UVA and UVB radiations exposure in participants of the Alienor Study (Bordeaux, 2006-2008)