1 **Intended category:** Review

2 **Title:** Compost and *Legionella longbeachae* – An emerging infection?

3 **Authors:** Sandra L. Currie and Tara K. Beattie

4 **Abstract**

5 Human disease caused by *Legionella* species is dominated by *Legionella pneumophila*, the main
6 causative agent in cases of Legionnaires' disease. However other species are known to cause infection,
7 e.g. *Legionella longbeachae* causes an equivalent number of cases of disease as *L. pneumophila* in
8 Australia and New Zealand. Infection with *L. longbeachae* is commonly associated with exposure to
9 composts and potting soils, and cases of infection with this organism have been increasing in Europe
10 over the past 10 years. The increase in incidence may be linked to factors such as increased awareness
11 of clinical presentation, or due to changing formulation of growing media, although it should be noted
12 that the presence of *Legionella* species in growing media does not correlate with the number of cases
13 currently seen. This is likely due to the variables associated with infection, for example, host factors such
14 as smoking or underlying health conditions, or difference in growing media storage or climate,
15 especially warm humid conditions, which may affect survival and growth of these organisms in the
16 growing media environment. There are numerous unknowns in this area and collaboration between
17 growing media manufacturers and researchers, as well as more awareness among diagnosing clinicians,
18 laboratory staff and the general public is necessary to reduce risk. More research is needed before
19 definitive conclusions can be drawn: *L. pneumophila* research currently dominates the field and it is
20 likely that the overreliance on diagnostic techniques such as the Urinary Antigen Test which is specific
21 for *L. pneumophila* Sg 1, is detrimental to the diagnosis of *L. longbeachae* infection.

22 **Introduction**

23 The term Legionnaires' disease was first coined in 1976, after 182 attendees at a meeting of the
24 American Legion showed symptoms of a mystery illness. It took a further 6 months before the causative
25 agent in the outbreak, a Gram negative rod, was identified and named *Legionella pneumophila* (1). Since
26 then, over 50 species of *Legionella* have been identified, with 64 serotypes.

27 Despite the numerous pathogenic species found, *L. pneumophila* is the main causative agent of human
28 disease. An international-collaborative study by Yu and colleagues, comparing the most common
29 causative agents in cases of sporadic community-acquired legionellosis from the USA (72.2% of cases
30 reviewed), Italy (12.6%), Switzerland (6.1%), New Zealand (4.3%) and Australia (4.7%), cited *L.*
31 *pneumophila* as the agent responsible for 91.5% of cases of Legionellosis; the second most commonly
32 isolated species, *Legionella longbeachae*, was present in 3.9% of cases (2). Despite the low
33 representation worldwide, *L. longbeachae* plays a significant role in the burden of legionellosis in the
34 southern hemisphere; in the study by Yu, 14 of the 20 *L. longbeachae* isolates came from Australia and
35 New Zealand (2). A review of legionellosis survey data in Southern Australia from 1996 to 2000 reported
36 that 42% of cases were attributable to *L. longbeachae*, compared with 51% due to *L. pneumophila* (3). In
37 Western Australia, between 1999 and 2010, 87% of diagnosed cases of Legionnaires' disease were
38 caused by *L. longbeachae*, whereas only 9% of cases were caused by *L. pneumophila* (4). Similarly, in
39 New Zealand, the Ministry of Health found that, in 2011, *L. longbeachae* was responsible for more cases
40 than *L. pneumophila*, with 42% and 30% of laboratory-reported cases of infection, respectively (5).
41 Human infection with *L. longbeachae* has also been noted in the USA (6), Japan (7) and Thailand, where
42 Phares et al. found that *L. longbeachae* was responsible for 5% of clinically defined cases of pneumonia
43 in a rural district, whereas *L. pneumophila* was not reported (8).

44 Historically, the incidence of infection with *L. longbeachae* in Europe has been low; however, as noted
45 by Whiley and Bentham, the number of cases of infection appears to be increasing (9). In 2012, Lindsay
46 et al noted that *L. longbeachae* had been cited as the causative agent in only 11 cases of infection in the
47 UK since 1984, seven of these occurred in Scotland (10). Further work revealed that between 2008 to
48 date, 26 cases of *L. longbeachae* infection had been detected in Scotland; in most cases the patient had
49 been in contact with commercially available growing media before the onset of symptoms (11) (personal
50 communication, Dr Kevin Pollock/Ross Campbell, Health Protection Scotland). In addition, an atypical
51 case of *L. longbeachae* cutaneous infection in a female patient in the UK was recently described (12).

52 This study presents a review of the literature currently available examining *Legionella* spp in the
53 compost habitat, particularly the status of *L. longbeachae* infection. Work was completed using Web of
54 Knowledge and PubMed searches including, but not limited to, individual and combinations of the
55 following terms: *Legionella*, Legionnaires disease, Pontiac fever, *Legionella longbeachae*, soil, compost,
56 growing media, garden, amoeba, PCR, diagnosis, biofilm, source, water. Searches were not truncated by
57 date, although the relatively recent discovery of this organism reduces the amount of searchable
58 literature available.

59 **Infection**

60 Infection with *Legionella* spp can be symptomatic or asymptomatic: hospital patients and healthy
61 individuals have been shown to experience increased antibody titres to *Legionella* antigens without
62 showing clinical signs of infection (13, 14). Symptomatic infection will generally present as legionellosis
63 in one of two distinct clinical manifestations: Pontiac fever (PF)—a self-limiting influenza-like illness; or
64 Legionnaires' disease (LD)—a more serious pneumonia that can be fatal. However there have also been
65 a number of atypical manifestations of *Legionella* infection, e.g. cutaneous infection (15) caused by *L.*

66 *pneumophila* Sg 8, prosthetic joint infection caused by *L. micdadei* (16) and a septic foot infection (17)
67 and endocarditis (18) both caused by *L. longbeachae*.

68 A number of symptoms, appearing after a 2-10 day incubation period, are associated with LD, including
69 malaise, shortness of breath, fever and diarrhoea; this is the most serious form of the disease and, on
70 average, is fatal in 10% of cases (19). Cases of infection can be community, nosocomial or travel-related;
71 in 2013 in the UK, the majority of cases were either community acquired (179 of 331) or travel-related
72 (148 of 331) (20). Pontiac fever is a less serious manifestation of infection, with flu-like symptoms
73 appearing 1-2 days after exposure, and resolving without intervention within a week (21). Unlike LD,
74 where both immunosuppression and increased age are risk factors for infection, PF does not appear to
75 discriminate between adults and children, healthy or immunocompromised individuals (22). Indeed,
76 exposure to a PF source is more likely to result in illness than exposure to a LD source. Information for
77 clinicians from the Centers for Disease Control and Prevention (CDC) shows that when exposed to the
78 source of LD, <5% individuals become ill, compared with >90% of those exposed to the source of PF (23).
79 The reason why exposure to *Legionella* spp results in different clinical manifestation remains unclear.
80 Rowbotham (24) suggested that pathogenesis of LD is caused by the invasion and replication of
81 *Legionella* bacteria within human cells, whereas PF is due to hypersensitivity caused by an unknown
82 component of the bacteria or an amoebal host.

83 Although around 40-50% of identified *Legionella* spp have been shown as agents of human disease
84 including *L. pneumophila*, *L. longbeachae*, *L. bozemanii*, *L. micdadei* and *L. anisa* (25, 26), many of these
85 are identified rarely in clinical samples, and others have only been identified once. There does not
86 appear to be a difference between species of *Legionella* and their ability to cause PF; *L. pneumophila*
87 and *L. longbeachae* have both been responsible agents in outbreaks (27, 28). Comparisons between the

88 clinical presentation of LD caused by *L. pneumophila* and *L. longbeachae* showed no significant
89 difference in symptoms observed between the two species (29).

90 When comparing the genomes of *L. pneumophila* and *L. longbeachae* there are a number of similarities,
91 perhaps indicating why these are the most successful pathogens in the *Legionella* genus. Gomez-Valero
92 et al found 124 genes specific to *L. pneumophila* and *L. longbeachae* which “increase successful infection
93 of mammalian cells” when comparing their genomes with those of *Legionella micdadei*, *Legionella*
94 *hackeliae* and *Legionella fallonii* (a *Legionella*-like amoebal pathogen designated LLAP-10), species much
95 less likely to cause disease in humans (30). However, unlike *L. pneumophila*, *L. longbeachae* lacks flagella
96 and produces a capsule (31), which along with a chemotaxis system and sequences for cellulolytic
97 enzymes in the *L. longbeachae* genome, but not the *L. pneumophila* genome, is likely to help its survival,
98 for example, in the potting soil environment, and from host defences (31). While *L. longbeachae* appear
99 to have adapted to soil life, it is also likely that *Legionellae* survival in compost and the composting
100 process is aided by an association with soil-dwelling free living amoebal host species, which may provide
101 a niche habitat away from the potentially harmful environment. Such protective symbiosis has been
102 noted before, for example, *Acanthamoeba* spp, which are often used for co-culture work (32), can both
103 protect and revive *L. pneumophila* after treatment with sodium hypochlorite (33). It should be noted
104 though that limited work has investigated such symbiotic relationships for *L. longbeachae*.

105 **Diagnosis and Treatment**

106 Fast accurate diagnosis is key to successful treatment of disease. There are numerous techniques
107 available for the diagnosis of *Legionella* spp infection, including the urinary antigen test (UAT),
108 serological testing, PCR and culture from patient samples. Culture on buffered charcoal yeast extract
109 agar (BCYE) is seen as the “gold-standard” in identification of *Legionella* spp; however, colony growth

110 can take 3-10 days which is much slower than other available methods and is undesirable in a clinical
111 setting where fast diagnosis is preferred (34). This is likely one of the reasons why 79% (5162/6601) of
112 cases in Europe in 2013 were identified by the UAT compared with only 11% (720/6601) identified by
113 culture (20). The UAT is only specific for *L. pneumophila* Sg 1 and may be a contributing factor in the late
114 diagnosis of infections caused by non-Sg1 *L. pneumophila* and other species of *Legionella*. In addition,
115 Thalanayar et al. showed that the urine test is not accurate in all cases; these authors found a negative
116 result when serum levels showed a positive reaction to *L. pneumophila* Sg1 and elevation from 1:64 to
117 1:1024 (35). Cases of *L. longbeachae* infections have been seen in Australia since 1989 (36) and it is likely
118 that this species is tested for more widely here than in the Northern Hemisphere due to increased
119 awareness amongst clinicians. Likewise, the recent increase in *L. longbeachae* infection seen across
120 Europe may also be linked to increased clinical awareness following media reports highlighting patient
121 case studies, clusters and research in this field. As well as incorrect or slow diagnosis of LD, the self-
122 limiting nature of PF means that it is unlikely to be properly diagnosed unless an outbreak occurs (37).

123 A cluster of *L. longbeachae* infection occurred in Scotland during summertime 2013 and four out of six
124 of these cases were initially identified using PCR in the NHS Lothian region; the diagnostic lab had
125 implemented *Legionella* spp PCR testing for all severe community acquired pneumonia (CAP) patients in
126 2010 (11). Work by Murdoch et al suggests that PCR diagnosis using primers targeting a *Legionella*
127 specific region of the 16S rDNA gene may be more effective even than the preferred culture method
128 (38). When comparing data on Legionellosis two years before and two years after the introduction of
129 PCR testing for *Legionella* spp on all respiratory specimens, the authors found a fourfold increase in
130 diagnosis of *Legionella* spp infection when moving from culture to PCR diagnosis (38). PCR is suitable in
131 the relatively fast identification of *Legionella* spp, and does not have the species limitations of the UAT.

132 The British Thoracic Society recommends the use of this technique over serological testing where
133 available (39). Use of this method as a preliminary identification technique prior to the culture of
134 samples for typing and confirmation may be beneficial in faster diagnosis of this disease in the future. In
135 the UK, pneumonia affects up to 11 in 1,000 adults each year (40) and can be caused by a number of
136 different bacteria, viruses and fungi. Lamoth and Greub noted while reviewing literature on respiratory
137 tract infections that the aetiological agent in 50% of CAP and 75% of nosocomial pneumonia remains
138 unknown and it is possible that a change in diagnostic practice could lead to identification of more cases
139 than currently observed (41).

140 Due to the slow nature of such diagnosis it is possible that the correct antibiotic regimen may not be
141 administered in a timely fashion leading to poorer patient outcomes, extended hospital stays and
142 inevitably escalating treatment costs. In addition, resistance of a variety of bacteria to all classes of
143 antibiotics has been seen to be increasing over time. The removal of the sources of infection is
144 preferable to overreliance on antibiotics for treatment, leaving drugs for patients who are most
145 seriously infected.

146 **Source**

147 *L. longbeachae* was first isolated in 1981, from a clinical sample taken from a patient with CAP (42).
148 Subsequent cases of LD where *L. longbeachae* was the aetiological agent have been widely linked to
149 gardening (10, 43, 28); cases reported range in severity from an outbreak of PF (28) to LD requiring
150 treatment in an intensive care unit (ICU) (11). The link between gardening and *L. longbeachae* was first
151 made by Steele et al, who isolated the organism from potting soils in South Australia after an outbreak
152 affecting 23 people identified gardening as a major risk factor for infection (44). Since then, *L.*
153 *longbeachae* has been isolated from compost and potting mixes in Japan (45), Switzerland (46), Greece

154 (47), Scotland (48), and the USA (6), but has not been isolated from water, unlike other species of
155 *Legionella*.

156 The high microbial diversity in growing media means that *Legionella* can be difficult to culture due to
157 inhibition by other organisms, plate overgrowth and insufficient agar media, and it may be the case that
158 sources of infection, other than water, have been overlooked in the past due to this fastidious nature of
159 *Legionella* spp. Increased identification of *Legionella* from this environment may also be due to the
160 changing composition of composts, for example, the reduction of peat content in the UK (49). It is
161 possible that variety in compost composition affects the conditions and subsequently different species
162 survival in growing media. Steele et al (50) isolated *Legionella* including *L. longbeachae* from potting
163 composts and green wastes, but not from peat alone. Two similar studies did not isolate *Legionella* spp
164 from 100% peat samples (45, 47). A report for the South Australian guidance committee noted that the
165 source of *Legionella* spp in compost is inconclusive, but that plants and trees may be a source of these
166 organisms (51). It is important to note that *L. longbeachae* was not isolated in a study looking at the
167 prevalence of *Legionellae* at compost making facilities and green waste storage plants in Switzerland
168 (52) but was isolated in a more recent Swiss study (53). Differences in detection may be due to the high
169 limit of detection of *Legionella* spp from environmental samples and subsequent growth during the
170 composting process.

171 Often the source of sporadic Legionellosis infection is not discovered. Of six cases of *L. longbeachae*
172 infection identified in Taiwan 2006-2010, only two identified specific soil exposure (54). In addition,
173 composts and soils should not be ruled out as a source of infection for other species of *Legionella*. *L.*
174 *pneumophila* Sg1 has been isolated from compost and soil samples (46-48, 55), and Wallis and Robinson

175 associated a case of *L. pneumophila* infection with soil (56). *L. pneumophila* pneumonia described by
176 Thacker et al was also thought to have soil as a source (57).

177 **Transmission**

178 The main route of transmission for LD is widely regarded as through the inhalation or aspiration of
179 water aerosols contaminated with *L. pneumophila* (26). For infection linked to compost use, there is
180 more debate. There have been suggestions that *Legionella* spp may be able to enter the body through
181 open abrasions in the skin (58, 59), while Steele et al suggested that *L. longbeachae* leaches out of
182 potting mix after watering, and may be present in any aerosols formed during the watering process,
183 which could be inhaled by the gardener (44). Work by Doyle et al found that an aerosolized Australian
184 clinical isolate of *L. longbeachae* Sg1 was lethal to 3 out of 5 exposed Guinea pigs, and lung tissue
185 showed similar characteristics to infection with *L. pneumophila* Sg1 upon post mortem examination (60),
186 suggesting that aerosolization would be a viable route of infection.

187 Inhalation or aspiration of live bacterial cells, contaminated dust or soil particles (61, 62), or protozoa
188 containing the bacteria (63) are also potential routes of infection. Rowbotham suggested that amoebae
189 or vesicles released from amoebae could prevent dehydration of legionellae and through inhalation
190 could provide a large dose of the bacteria to a potential host (64). Work by Cabello-Vílchez et al provides
191 support for this theory as *Acanthamoeba* spp were isolated from 21 (28.4%) of 74 nasal swabs taken
192 from healthy individuals in Peru (65), and another study found amoebae-resisting bacteria after
193 amoebal co-culture of human nasal swabs (7 out of 444 samples)(66). Berk et al also described the
194 release of respirable vesicles containing live clusters of *L. pneumophila* by *Acanthamoeba polyphaga*
195 and *Acanthamoeba castellanii* (67). Although Cramp et al described a cluster of PF attributed to
196 aerosolized potting mix, the source of infection remains unclear, as contaminated soil, dust, water,

197 protozoa and bacteria may all have been present in the air (28). However, this does support the theory
198 that the inhalation of aerosols consisting of contaminated water or compost particles is the most likely
199 route of infection, as does the evidence that this is the method for transmission of *Legionella* spp found
200 in water. Proximity to dripping hanging flower pots was found to be a predictor for infection and
201 aerosolization suggested as a likely mode of transmission (68). Conza et al isolated *L. pneumophila* and
202 Free-living amoeba (FLA) from 10.6% (5/47) and 19.1% (9/47) of bioaerosol samples, respectively,
203 collected at composting facilities; however the authors did not isolate *L. pneumophila* and FLA
204 simultaneously from the same samples, including potential intracellular *Legionella* spp (53). The
205 evidence suggests that transmission occurs when live *Legionella* spp, or contaminated compost particles
206 or water droplets are aerosolized when the growing media is handled, when bags are opened or when
207 the material is watered.

208 **Risk Factors**

209 The number of cases of *L. longbeachae* reported does not tally with the frequency with which the
210 organism is isolated from growing media. The potential for greenhouse storage to increase levels of
211 legionellae in growing media was noted, based on observations of amoebal enrichment and a
212 preliminary study, by Lindsay et al (10). A limited increase of legionellae was seen in some but not all
213 amoebal enrichment studies (45, 48) which may suggest that these species increase in numbers in warm
214 humid conditions, for example as could be provided in a greenhouse. Recent work examining a cluster
215 of six *L. longbeachae* infections did not identify a common growing media product or manufacturer, but
216 did isolate the organism in growing media from 5 out of 6 cases (11). It was noted that growing media
217 had been stored inside the house, greenhouse, car, polytunnel, shed or garage of the infected
218 individuals. This, combined with the higher than normal temperatures seen in Scotland during the time

219 that this cluster occurred, leads the authors to suggest that climatic conditions and storage of the
220 growing media may have enabled high levels of growth, leading to increased risk of human infection. An
221 analysis of 1676 community-acquired cases in England and Wales between 1993 – 2008 identified a
222 higher risk of sporadic LD after warm wet weather (69), and wet, warm and humid weather was linked
223 to the occurrence of legionellosis in metropolitan Philadelphia between 1995-2003 (70).

224 O'Connor et al highlights that presence of *Legionella* spp in growing media does not necessarily indicate
225 that those handling it will become infected, and also that education of potential risk factors and hand
226 washing before eating, drinking and smoking was shown to decrease incidence of infection (68).

227 **Conclusions**

228 While *L. pneumophila* Sg1 is the main causative agent of LD, *L. longbeachae* is responsible for a
229 significant burden of legionellosis infection in the southern hemisphere, particularly Australia and New
230 Zealand; in animal studies, Australian strains of *L. longbeachae* were more virulent than strains from
231 elsewhere (71), which may help to account for the discrepancy in infection rate between Australia and
232 other countries. However there has been an apparent increase in cases of infection caused by *L.*
233 *longbeachae* in UK in the last ten years. This may be linked to factors such as increased awareness of
234 clinical presentation, or due to changing formulation of growing media. *Legionella* spp were isolated
235 from 62.5% (15/24) of UK compost samples (48), but the prevalence in compost does not correlate with
236 number of cases currently seen. Both *L. pneumophila* and *L. longbeachae* may be adapted to infect
237 mammalian cells better than other species (30), however as these species are not the most commonly
238 found in growing media, the potential for infection via this route is low, i.e. the presence of *Legionella*
239 spp in growing media may not be indicative of the risk of infection.

240 The variables involved in *Legionella* related infection linked to compost are summarised in Figure 1.
241 More research is needed before definitive conclusions can be drawn. *L. pneumophila* research
242 dominates the field; a crude search using ISI Web of Science gives 369 hits and 29,632 hits when
243 searching for “*longbeachae*” and “*pneumophila*” respectively. There are numerous unknowns in this
244 area and collaboration between growing media manufacturers and researchers, as well as more
245 awareness among diagnosing clinicians, laboratory staff and the general public, is necessary. It is likely
246 that specific conditions are needed before infection occurs, including: host factors such as smoking or
247 underlying health conditions; storage and climate, especially warm humid conditions; transmission of
248 infective agent through method of compost use; and presence or absence of pathogenic strains and
249 their host species in the growing media environment, which may be impacted by composition. McDade
250 highlights the importance that recognition and pursuit of anomalies in routine investigation plays in new
251 discoveries, and the potential pitfalls of sticking to a standard diagnostic algorithm (72). This may well
252 be true of the current system: while the importance of *L. pneumophila* Sg1 as an aetiological agent
253 cannot be denied, it is likely that the overreliance on the Urinary Antigen Test is detrimental to the
254 diagnosis of *L. longbeachae* infection.

255

256

257

258

259

260

261 Figure 1 Variable factors related to the occurrence and recording of *Legionella* spp infections linked to
 262 compost use

264 **References**

- 265 1. McDade JE, Shepard CC, Fraser DW, Tsai TR, Redus MA, Dowdle WR. Legionnaires Disease - Isolation
266 of a bacterium and demonstration of its role in other respiratory disease. N Engl J Med.
267 1977;297(22):1197-1203.
- 268 2. Yu VL, Plouffe JF, Pastoris MC, Stout JE, Schousboe M, Widmer A, et al. Distribution of Legionella
269 species and serogroups isolated by culture in patients with sporadic community-acquired legionellosis:
270 An international collaborative survey. J Infectious Dis. 2002;186(1):127-128.
- 271 3. Li JS, O'Brien ED, Guest C. A review of national legionellosis surveillance in Australia, 1991 to 2000.
272 Communicable diseases intelligence quarterly report. 2002;26(3):461-468.
- 273 4. Commission for Occupational Safety and Health. *Code of practice prevention and control of*
274 *Legionnaires' disease*. Report for Government of Western Australia, Department of Commerce,
275 Department of Mines and Petroleum 2010 ISBN 978-1-920836-49-8
- 276 5. The Institute of Environmental Science and Research Ltd (IESRNZ). Notifiable and other diseases in
277 New Zealand: Annual Report 2011,
278 http://www.surv.esr.cri.nz/surveillance/annual_surveillance.php?we_objectID=2963 (April 2012
279 accessed 16/06/2015).
- 280 6. Duchin JS, Koehler J, Kobayashi JM, Rakita RM, Olson K, Hampson NB, et al. Legionnaires' diseases
281 associated with potting soil: California, Oregon, and Washington, May-June 2000. MMWR.
282 2000;49(34):777-778.

283 7. Okazaki M, Umeda B, Koide M, Saito A. Legionella longbeachae pneumonia in a gardener.
284 Kansenshogaku Zasshi. 1998;72(10):1076-1079.

285 8. Phares CR, Wangroongsarb P, Chantira S, Paveenkitiporn W, Tondella M-L, Benson RF, et al.
286 Epidemiology of severe pneumonia caused by Legionella longbeachae, Mycoplasma pneumoniae, and
287 Chlamydia pneumoniae: 1-year, population-based surveillance for severe pneumonia in Thailand. Clin
288 Infect Dis. 2007;45(12):E147-E55.

289 9. Whiley H, Bentham R. Legionella longbeachae and Legionellosis. Emerg Infect Dis. 2011;17(4):579-
290 583.

291 10. Lindsay DSJ, Brown AW, Brown DJ, Pravinkumar SJ, Anderson E, Edwards GFS. Legionella
292 longbeachae serogroup 1 infections linked to potting compost. J Med Microbiol. 2012;61(2):218-222.

293 11. Potts A, Donaghy M, Marley M, Othieno R, Stevenson J, Hyland J, et al. Cluster of Legionnaires'
294 disease cases caused by Legionella longbeachae serogroup 1, Scotland, August to September 2013. Euro
295 Surveill. 2013;18(50):9-13.

296 12. Grimstead D, Tucker D, Harris K, Turner D. Cutaneous Legionella longbeachae Infection in
297 Immunosuppressed Woman, United Kingdom. *Emerging Infectious Diseases* 2015; 21(8):1426–8.

298 13. Grove ID, Lawson PJ, Burgess JS, Moran JL, O'Fathartaigh MS, Winslow WE. An outbreak of Legionella
299 longbeachae infection in an intensive care unit? J Hosp Infect. 2002;52(4):250-258.

300 14. Rudbeck M, Molbak K, Uldum SA. Dynamics of Legionella antibody levels during 1 year in a healthy
301 population. Epidemiol Infect. 2009;137(7):1013-1018.

302 15. Padrnos LJ, Blair JE, Kusne S, DiCaudo DJ, Mikhael JR. Cutaneous legionellosis: Case report and review
303 of the medical literature. *Transplant Infectious Disease* 2014; 16(2): 307–14.

304 16. Fernández-Cruz A, Marín M, Castelo L, Usubillaga R, Martín-Rabadán P, Bouza E *et al.* *Legionella*
305 *micdadei*, a new cause of prosthetic joint infection. *Journal of Clinical Microbiology* 2011; 49(9): 3409–
306 10.

307 17. Dugar M, Rankin WA, Rowe E, Smith MD. "My foot hurts": a flare of rheumatoid arthritis? *Med J*
308 *Australia*. 2009;190(7):392-393.

309 18. Leggieri N, Gouriet F, Thuny F, Habib G, Raoult D, Casalta J-P. *Legionella longbeachae* and
310 Endocarditis. *Emerg Infect Dis*. 2012;18(1):95-97.

311 19. Beauté J, de Jong B. European Centre for Disease Prevention and Control. Legionnaires' disease in
312 Europe, 2010. Stockholm: ECDC; 2012.

313 20. Beauté J, Robesyn E. European Centre for Disease Prevention and Control. Legionnaires' disease in
314 Europe, 2013. Stockholm: ECDC; 2015.

315 21. Kaufmann AF, McDade JE, Patton CM, Bennett JV, Skaliy P, Feeley JC, et al. PONTIAC FEVER -
316 ISOLATION OF THE ETIOLOGIC AGENT (LEGIONELLA-PNEUMOPHILA) AND DEMONSTRATION OF ITS
317 MODE OF TRANSMISSION. *Am J Epidemiol*. 1981;114(3):337-347.

318 22. Goldberg DJ, Wrench JG, Collier PW, Emslie JA, Fallon RJ, Forbes GI, et al. LOCHGOILHEAD FEVER -
319 OUTBREAK OF NON-PNEUMONIC LEGIONELLOSIS DUE TO LEGIONELLA-MICDADEI. *Lancet*.
320 1989;1(8633):316-318.

321 23. Centers for Disease Control and Prevention. Legionella (Legionnaires' Disease and Pontiac Fever) for
322 clinicians <http://www.cdc.gov/legionella/clinicians.html> (Feb 2013 accessed 12/06/2015)

323 24. Rowbotham TJ. PONTIAC FEVER EXPLAINED. *Lancet*. 1980a;2(8201):969.

324 25. Muder RR, Yu VL. Infection due to Legionella species other than L-pneumophila. *Clin Infect Dis*.
325 2002;35(8):990-998.

326 26. Fields BS, Benson RF, Besser RE. Legionella and Legionnaires' disease: 25 years of investigation. *Clin*
327 *Microbiol Rev*. 2002;15(3):506-526.

328 27. Benin AL, Benson RF, Arnold KE, Fiore AE, Cook PG, Williams LK, et al. An outbreak of travel-
329 associated legionnaires disease and pontiac fever: The need for enhanced surveillance of travel-
330 associated legionellosis in the United States. *J Infect Dis*. 2002;185(2):237-243.

331 28. Cramp GJ, Harte D, Douglas NM, Graham F, Schousboe M, Sykes K. An outbreak of Pontiac fever due
332 to Legionella longbeachae serogroup 2 found in potting mix in a horticultural nursery in New Zealand.
333 *Epidemiol Infect*. 2010;138(1):15-20.

334 29. Amodeo MR, Murdoch DR, Pithie AD. Legionnaires' disease caused by Legionella longbeachae and
335 Legionella pneumophila: comparison of clinical features, host-related risk factors, and outcomes. *Clin*
336 *Microbiol Infect*. 2010;16(9):1405-1407.

337 30. Gomez-Valero L, Rusniok C, Rolando M, Neou M, Dervins-Ravault D, Demirtas J, et al. Comparative
338 analyses of Legionella species identifies genetic features of strains causing Legionnaires' disease.
339 *Genome Biology*. 2014;15(11).

340 31. Cazalet C, Gomez-Valero L, Rusniok C, Lomma M, Dervins-Ravault D, Newton HJ, et al. Analysis of the
341 *Legionella longbeachae* Genome and Transcriptome Uncovers Unique Strategies to Cause Legionnaires'
342 Disease. Plos Genet. 2010;6(2).

343 32. Rowbotham TJ. ISOLATION OF LEGIONELLA-PNEUMOPHILA FROM CLINICAL SPECIMENS VIA
344 AMEBAS, AND THE INTERACTION OF THOSE AND OTHER ISOLATES WITH AMEBAS. J Clin Pathol.
345 1983;36(9):978-986.

346 33. Garcia MT, Jones S, Pelchocaz C, Millar RD, Abu Kwaik Y. Acanthamoeba polyphaga resuscitates
347 viable non-culturable Legionella pneumophila after disinfection. Environ Microbiol. 2007;9(5):1267-
348 1277.

349 34. Hayden RT, Uhl JR, Qian X, Hopkins MK, Aubry MC, Limper AH, et al. Direct detection of Legionella
350 species from bronchoalveolar lavage and open lung biopsy specimens: Comparison of LightCycler PCR, in
351 situ hybridization, direct fluorescence antigen detection, and culture. J Clin Microbiol. 2001;39(7):2618-
352 2626.

353 35. Thalanayar P, Mahendiran S, Ho J, Holguin F. Legionella pneumonia with diffuse macular purpuric
354 rash and negative urine test: a diagnostic dilemma. Chest 2014; 145(3 Suppl); 119A

355 36. Lim I, Sangster N, Reid DP, Lanser JA. LEGIONELLA-LONGBEACHAE PNEUMONIA - REPORT OF 2
356 CASES. Med J Australia. 1989;150(10):599-601.

357 37. Nicolay N, Boland M, Ward M, Hickey L, Collins C, Lynch M, et al. Investigation of Pontiac-like illness
358 in office workers during an outbreak of Legionnaires' disease, 2008. Epidemiol Infect.
359 2010;138(11):1667-1673.

360 38. Murdoch DR, Podmore RG, Anderson TP, Barratt K, Maze MJ, French KE, et al. Impact of Routine
361 Systematic Polymerase Chain Reaction Testing on Case Finding for Legionnaires' Disease: A Pre-Post
362 Comparison Study. Clin Infect Dis. 2013;57(9):1275-1281.

363 39. British Thoracic Society (2009) Guidelines for the Management of Community Acquired Pneumonia
364 in Adults: A Quick Reference Guide, British Thoracic Society Reports, 1 (3), p4- 11. [https://www.brit-](https://www.brit-thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-guide-bts-guidelines-for-the-management-of-community-acquired-pneumonia-in-adults/)
365 [thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-](https://www.brit-thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-guide-bts-guidelines-for-the-management-of-community-acquired-pneumonia-in-adults/)
366 [guide-bts-guidelines-for-the-management-of-community-acquired-pneumonia-in-adults/](https://www.brit-thoracic.org.uk/document-library/clinical-information/pneumonia/adult-pneumonia/a-quick-reference-guide-bts-guidelines-for-the-management-of-community-acquired-pneumonia-in-adults/) (2009
367 accessed 16/06/15)

368 40. Marshall H. NICE guidelines to family doctors on diagnosis of pneumonia. Lancet Resp Med.
369 2015;3(1):17-.

370 41. Lamoth F, Greub G. Amoebal pathogens as emerging causal agents of pneumonia. FEMS Microbiol
371 Rev. 2010; 34: 260-280.

372 42. McKinney RM, Porschen RK, Edelstein PH, Bissett ML, Harris PP, Bondell SP, et al. LEGIONELLA-
373 LONGBEACHAE SPECIES NOVA, ANOTHER ETIOLOGIC AGENT OF HUMAN PNEUMONIA. Ann Intern Med.
374 1981;94(6):739-743.

375 43. den Boer JW, Yzerman EPF, Jansen R, Bruin JP, Verhoef LPB, Neve G, et al. Legionnaires' disease and
376 gardening. Clin Microbiol Infect. 2007;13(1):88-91.

377 44. Steele TW, Lanser J, Sangster N. ISOLATION OF LEGIONELLA-LONGBEACHAE SEROGROUP-1 FROM
378 POTTING MIXES. Appl Environ Microbiol. 1990a;56(1):49-53.

379 45. Koide M, Arakaki N, Saito A. Distribution of *Legionella longbeachae* and other legionellae in Japanese
380 potting soils. *J Infect Chemother*. 2001;7(4):224-227.

381 46. Casati S, Gioria-Martinoni A, Gaia V. Commercial potting soils as an alternative infection source of
382 *Legionella pneumophila* and other *Legionella* species in Switzerland. *Clin Microbiol Infect*.
383 2009;15(6):571-575.

384 47. Velonakis EN, Kiouisi IM, Koutis C, Papadogiannakis E, Babatsikou F, Vatopoulos A. First isolation of
385 *Legionella* species, including *L. pneumophila* serogroup 1, in Greek potting soils: possible importance for
386 public health. *Clin Microbiol Infect*. 2010;16(6):763-766.

387 48. Currie SL, Beattie TK, Knapp CW, Lindsay DSJ. *Legionella* spp. in UK composts-a potential public
388 health issue? *Clin Microbiol Infect*. 2014;20(4):O224-O229.

389 49. HM Government, UK. White Paper. The natural choice: securing the value of nature.
390 <http://www.officialdocuments.gov.uk/document/cm80/8082/8082.pdf>. (2011 accessed 16/06/2015)

391 50. Steele TW, Moore CV, Sangster N. DISTRIBUTION OF *LEGIONELLA*-*LONGBEACHAE* SEROGROUP-1
392 AND OTHER *LEGIONELLAE* IN POTTING SOILS IN AUSTRALIA. *Appl Environ Microbiol*. 1990b;56(10):2984-
393 2988.

394 51. Broadbent C (1996) Guidance for the Control of *Legionella*. South Australian Guidance Report,
395 National Environmental Health Forum Monographs Water Series No. 1, 1996.

396 52. Casati S, Conza L, Bruin J, Gaia V. Compost facilities as a reservoir of *Legionella pneumophila* and
397 other *Legionella* species. *Clinical Microbiology and Infection* 2010; 16: 945–7.

398 53. Conza L, Pagani SC, Gaia V. Presence of Legionella and free-living Amoebae in composts and
399 bioaerosols from composting facilities. *PLoS One* 2013; 8(7): e68244.

400 54. Wei S-H, Tseng L-R, Tan J-K, Cheng C-Y, Hsu Y-T, Cheng E-T, et al. Legionnaires' disease caused by
401 Legionella longbeachae in Taiwan, 2006-2010. *Int J Infect Dis.* 2014;19:95-97.

402 55. van Heijnsbergen E, Husman AMdR, Lodder WJ, Bouwknecht M, van Leeuwen AED, Bruin JP, et al.
403 Viable Legionella pneumophila bacteria in natural soil and rainwater puddles. *J Appl Microbiol.*
404 2014;117(3):882-890.

405 56. Wallis L, Robinson P. Soil as a source of Legionella pneumophila serogroup 1 (Lp1). *Aust N Z J Public*
406 *Health.* 2005;29(6):518-520.

407 57. Thacker SB, Bennett JV, Tsai TF, Fraser DW, McDade JE, Shepard CC, et al. OUTBREAK IN 1965 OF
408 SEVERE RESPIRATORY ILLNESS CAUSED BY LEGIONNAIRES DISEASE BACTERIUM. *J Infect Dis.*
409 1978;138(4):512-519.

410 58. Patten SM, Sur E, Sundaram R, Weinhardt B. Dangers in the garden. *Lancet.* 2010;376(9743):844

411 59. Lowry PW, Blankenship RJ, Gridley W, Troup NJ, Tompkins LS. A CLUSTER OF LEGIONELLA-STERNAL
412 WOUND INFECTIONS DUE TO POSTOPERATIVE TOPICAL EXPOSURE TO CONTAMINATED TAP WATER.
413 *New Engl J Med.* 1991;324(2):109-113.

414 60. Doyle RM, Steele TW, McLennan AM, Parkinson IH, Manning PA, Heuzenroeder MW. Sequence
415 analysis of the mip gene of the soilborne pathogen Legionella longbeachae. *Infect Immun.*
416 1998;66(4):1492-1499.

417 61. de Jong B, Zucs P. Legionella, springtime and potting soils. *Euro Surveill.* 2010;15(8):2-3.

418 62. Cameron S, Roder D, Walker C, Feldheim J. Epidemiologic characteristics of Legionella infection in
419 South-Australia - Implications for disease-control. Aust N Z J Med. 1991;21(1):65-70.

420 63. Atlas RM. Legionella: from environmental habitats to disease pathology, detection and control.
421 Environ Microbiol. 1999;1(4):283-293.

422 64. Rowbotham TJ. Preliminary-report on the pathogenicity of *Legionella pneumophila* for fresh-water
423 and soil amebas. J Clin Pathol. 1980b;33(12):1179-1183.

424 65. Cabello-Vilchez AM, Martin-Navarro CM, Lopez-Arencibia A, Reyes-Batlle M, Gonzalez AC, Guerra H,
425 et al. Genotyping of potentially pathogenic Acanthamoeba strains isolated from nasal swabs of healthy
426 individuals in Peru. Acta Tropica. 2014;130:7-10.

427 66. Greub G, La Scola B, Raoult D. Amoebae-resisting bacteria isolated from human nasal swabs by
428 amoebal coculture. Emerg Infect Dis. 2004;10(3):470-477.

429 67. Berk SG, Ting RS, Turner GW, Ashburn RJ. Production of respirable vesicles containing live Legionella
430 pneumophila cells by two Acanthamoeba spp. Appl Environ Microbiol. 1998;64(1):279-286.

431 68. O'Connor BA, Carman J, Eckert K, Tucker G, Givney R, Cameron S. Does using potting mix make you
432 sick? Results from a Legionella longbeachae case-control study in South Australia. Epidemiol Infect.
433 2007;135(1):34-39.

434 69. Halsby KD, Joseph CA, Lee JV, Wilkinson P. The relationship between meteorological variables and
435 sporadic cases of Legionnaires' disease in residents of England and Wales. Epidemiol Infect.
436 2014;142(11):2352-2359.

437 70. Fisman DN, Lim S, Wellenius GA, Johnson C, Britz P, Gaskins M, et al. It's not the heat, it's the
438 humidity: Wet weather increases Legionellosis risk in the greater Philadelphia metropolitan area. *J Infect*
439 *Dis.* 2005;192(12):2066-2073.

440 71. Doyle RM, Cianciotto NP, Banvi S, Manning PA, Heuzenroeder MW. Comparison of virulence of
441 *Legionella longbeachae* strains in guinea pigs and U937 macrophage-like cells. *Infect Immun.*
442 2001;69(9):5335-5344.

443 72. McDade JE. Legionnaires' disease 25 years later: Lessons learned. In Marre et al (ed.) *Legionella*.
444 ASM Press, Washington, D.C. 2002; 1–10.

445