

'CBT-I in Cancer: We Know It Works, So Why Are We Waiting?'

Leanne Fleming, Ph.D.^{1*}

Kenneth MacMahon, Ph.D.²

¹ Division of Psychology

University of the West of Scotland

Paisley, PA1 2BE

Scotland, UK

Tel: +44(0)141 849 4088

Email: Leanne.Fleming@uws.ac.uk

*Corresponding author

² Section of Clinical Psychology

School of Health in Social Science

University of Edinburgh

Edinburgh, EH8 9AG

Tel: +44(0) 131 651 3960

E-mail: ken.macmahon@ed.ac.uk

Abstract

Insomnia is one of the most frequently reported and debilitating difficulties associated with cancer. Recent decades have seen a move from pharmacological interventions for insomnia, to non-pharmacological, cognitive behavioural therapies (CBT-I). Numerous clinical trials have established the effectiveness of CBT-I in the general population and more recently, in those with insomnia associated with cancer. However, despite these promising outcomes, the availability of such therapies remains limited across cancer services. Recent years have seen developments to widen access to CBT-I, including the use of internet-based resources. Such developments may offer a useful means of overcoming the availability and access issues of CBT-I for those with insomnia associated with cancer.

Keywords: cancer; sleep; insomnia; cognitive behaviour therapy; intervention; psychological

Introduction

Despite its pervasiveness and chronicity, insomnia in cancer care is under-reported by patients, overlooked by clinicians and as a consequence, poorly managed. This is partly due to the belief that insomnia is a transient response to cancer diagnosis and treatment, which will self-resolve. However, insomnia which remains untreated has consequences for psychological and physical wellbeing. Diminished quality of life, impaired mood, heightened pain and increased fatigue are all common responses to persistent insomnia disorder, and many of those living with cancer experience unremitting insomnia which impacts upon daytime functioning. Evidence demonstrates that cognitive behaviour therapy (CBT) should be the first-line intervention for insomnia disorder and patients have expressed their acceptance of this treatment approach. However, despite this evidence, CBT is not routinely available to cancer populations, perhaps due to uncertainty around when and how it could most usefully be made available.

Cancer and insomnia prevalence

According to the World Health Organisation, cancer is among the leading causes of morbidity and mortality worldwide. Approximately 14 million new cases were recorded in 2012 and this figure is expected to rise by around 70% over the next 2 decades (1). In particular, breast cancer incidence has increased from a rate of 103/100,000 in 1975 to 126/100,000 in 2007 and there is an increased incidence of colorectal cancer in developed countries (2). Both breast and colorectal cancers have been linked to circadian disruption (3) and their epidemiology correlates with regions where circadian disruption, through night-shift working, for example, is most

prevalent (4). Although tentative, this evidence may suggest a circadian disruption-based model for cancer, particularly breast and colorectal cancers, which should be further explored.

Fortunately, due to screening programmes, earlier disease detection and improving treatment regimes, cancer survival rates are also increasing. Recent estimates indicate that around half of those diagnosed with cancer are expected to survive for at least 10 years and in the UK alone, cancer survival rates have doubled in the last 40 years (5) However these improvements in rates of survivorship are not without cost. Evidence shows that many cancer survivors experience a significant symptom burden (3), particularly at the end of active treatment when treatment side effects often fail to remit (4).

One of the most commonly reported side effects of cancer diagnosis / treatment is insomnia, which refers to difficulty initiating or maintaining sleep at least three nights per week for at least three months, accompanied by impaired daytime functioning (5). Epidemiological data from cancer populations report that 25-69% report difficulty sleeping, with 18-29% reporting sleep disturbance that meets diagnostic criteria for insomnia disorder (9–11). However, despite increasing awareness of the pervasiveness of insomnia within cancer groups, scientific reports of insomnia prevalence remain variable and wide-ranging. This is partly due to the utilisation of different insomnia definitions, measurements and timing of assessments (8). However, recent amendments to the diagnostic criteria for insomnia disorder (8) and the publication of clinical screening tools based upon these new criteria (12) should result in improvements to the consistency of prevalence estimates in future studies.

Evolution of insomnia in cancer

Although insomnia is a prevalent problem for cancer patients, little information is available about its long-term course and limited studies have focused on insomnia as a persistent disorder in oncology groups. However, scientific understanding of the evolution of insomnia has been applied in the context of cancer, and etiological factors including insomnia predisposition, precipitation and perpetuation have been proposed (13). One recent report on the natural history of insomnia in cancer patients reported data on incidence, persistence, remission and relapse of insomnia over the 18-month period following diagnosis in a group of patients with mixed cancer types (11). The study revealed high rates of insomnia at baseline (59%), including 28% with insomnia disorder. Insomnia became less prevalent over the 18-month assessment period but did remain pervasive at the end of the study (36%). Approximately 15% of patients experienced a first incidence of insomnia during the study and around 20% experienced relapse. Patients categorised as having an insomnia disorder were much less likely to experience remission than those who had insomnia symptoms. Those meeting criteria for insomnia disorder at baseline retained that sleep status for the duration of the study, highlighting its persistence over time. This study provides a useful analysis of the evolution of insomnia disorder during the cancer diagnosis / treatment / rehabilitation phases and indicates the point(s) at which cancer patients may be most vulnerable to developing a persistent complaint of poor sleep.

The elevated prevalence of insomnia amongst cancer patients is partly explained by heightened psychological distress following diagnosis (15). However, the side-effects from active cancer treatment are also known to contribute to the deterioration of good sleep (16–18). A large scale prospective study of cancer patients undergoing

chemotherapy, found that 39.8% reported moderate or severe insomnia symptoms after their initial chemotherapy session. Furthermore, two-thirds of patients reported that insomnia symptoms persisted throughout their subsequent cycles of chemotherapy (9). In particular, sleep maintenance problems are the most frequently reported sleep complaint during active treatment, with 63.3% of patients experiencing fragmented sleep patterns (19).

Once established, insomnia often becomes a chronic, unremitting complaint. RCT data from a cancer survivor population reported a median insomnia duration of greater than 2 years (20). Similarly, a cross-sectional survey of 982 cancer patients found that 31% reported insomnia, of which 75% had a chronic disorder (i.e., lasting six months or more). Therefore, it seems that insomnia often takes a chronic course in many cancer patients (21). This pattern of persistence may be explained by individual responses to sleep disturbance that influence its maintenance over time (e.g. maladaptive sleep behaviour, faulty beliefs). These individual responses are potentially more commonly observed in cancer populations than in the general population, due to their focus on restoration and recovery. In order to cope with the physical demands of active treatment (i.e. surgery, chemotherapy, radiotherapy), cancer patients may be required to rest more than usual during the day. Qualitative reports from cancer survivor populations indicate that they implement earlier bedtimes, later rising times and increased daytime napping during their cancer treatment (22). These maladaptive sleep habits encourage the development of an irregular sleep-wake pattern, which may ultimately lead to de-synchronisation of the sleep-wake cycle and contribute to the development of a chronic insomnia disorder (13). Therefore, disease-specific factors (such as psychological distress, active cancer treatment) may contribute to the precipitation of insomnia, whilst poor sleep

behaviours arising as a result of attempts to combat these disease specific factors, may contribute to its maintenance.

Consequences of insomnia in cancer

Insomnia disorder is not only prevalent and persistent, but is consequential for both psychological and physical wellbeing. Research indicates that good sleep quality, characterised by consistent and distinguishable sleep/wake patterns, reduces depressive symptomatology, enhances quality of life and improves physical outcomes (23–25). However, due to the high prevalence of insomnia following diagnosis, cancer patients are especially vulnerable to the negative impact of sleep disruption. Also, insomnia often occurs alongside other symptoms like fatigue, anxiety and low mood (18,26–28), causing further impairment to quality of life. Liu et al found that breast cancer patients who reported multiple symptoms (symptom clusters) prior to the onset of active treatment, experienced poorer sleep, more fatigue and reduced mood during cancer treatment (26). Therefore one symptom may contribute to the maintenance and exacerbation of the others (29).

Associations between commonly occurring symptoms have also been explored in the general population. For example, epidemiological data indicates that insomnia contributes to the development of depressive disorder (27) and individuals with cancer and insomnia experience reduced functioning, increased pain and more fatigue than those without insomnia (30,31). Sharma et al, reported that almost one third of cancer patients described sleep problems reaching clinical levels, which were strongly related to symptoms of psychological distress (28). Therefore, given the increasing number of cancer survivors who are likely to have comorbid symptoms (including insomnia), there is a considerable burden on healthcare providers to

understand and implement symptom management strategies to cope with these increasing demands.

Cognitive Behaviour Therapy for insomnia (CBT-I) in cancer

The consequences of insomnia can be severe enough to warrant medical intervention, with 25% of cancer patients regularly taking sleeping tablets. However, this long term pharmacotherapy is undesirable to patients (32). There is considerable evidence that intervening at the psychosocial level, using cognitive behavioural therapy, is effective for insomnia associated with cancer (20,33–40) and is a treatment approach that patients prefer (22).

Cognitive behavioural therapy for insomnia (CBT-I) is a multicomponent intervention that was developed to target the specific cognitive, physiological and behavioural aspects that characterise the insomnia disorder. Essentially, CBT-I comprises five potential elements, including stimulus control, sleep restriction, sleep hygiene, cognitive restructuring and relaxation training. Evidence demonstrates that CBT-I can be successfully administered using a range of delivery methods, including groups (41,42), telephone (43), online (44,45) and self-help manuals (46). Importantly, the effectiveness of CBT-I has been demonstrated for insomnia that occurs alongside complex psychiatric and medical conditions as well as for insomnia without those comorbidities.

To date, nine controlled trials of CBT-I in cancer patients have been conducted (summarised in Table I) (20,33,35–40,47–49). The largest of these trials using face-to-face CBT-I was conducted by Espie et al (2008) (20). This study recruited 150 cancer patients (mixed tumour sites) to either a group delivered, nurse led CBT-I intervention or to a treatment as usual (TAU) condition. Significant

improvements in subjectively reported sleep onset latency, wake time after sleep onset and sleep efficiency were noted in the CBT-I group, with large between-group effect sizes of -0.86, -0.97, and 1.09, respectively. These results represented a reduction of approximately 1 hour in sleep onset latency and time spent awake during the night. There was no change in the TAU group and results were sustained at 6 month follow-up assessment. Alongside these improvements in sleep outcomes, improvements to quality of life, mood and levels of cancer-related fatigue were also reported in the CBT-I condition (29).

[Table 1 here](#)

Recent Advances in CBT Delivery

Although numerous studies have identified CBT as an effective and well-tolerated therapy for insomnia associated with cancer, this therapeutic approach is not always widely available to patients. Limited availability of therapists with sufficient expertise and geographical distance from treatment centres can prove to be barriers. This is something that is not limited to those with cancer: treatment for sleep disorders in general is often piece-meal across services (50). However, a concomitant physical illness can bring with it further difficulties in accessing treatment. The diagnosis of cancer can bring significant financial burdens, meaning that the cost of additional travel to treatment centres may be prohibitive (51). Furthermore, the physical effects of both the illness and its treatment may preclude others from seeking additional treatment.

The issues of availability and access have been explored widely in the general literature for psychological therapies, with increasing attempts, over recent decades, to train non-psychologists to deliver psychological therapies. In particular, attention has been paid to developing the skills of nursing members of the multi-disciplinary health team (52). A similar process has taken place within the general insomnia literature, with a substantial focus on widening access to CBT for insomnia through dissemination of treatment programmes to a wider pool of health professionals, such as community nursing staff (53). Such efforts provide a wider pool of therapists, thereby increasing the availability of the therapy and potentially, reducing the cost of delivery. A further step in this process has been the use of guided self-help materials (often utilising the internet) as a means of delivering psychological therapies. Positive outcomes for these have been found in the general literature for psychological disorders, as well as in studies specific to insomnia (44,54,55). The availability of differing treatment delivery methods allows the opportunity for a stepped-care model of care within services, thereby increasing access and reducing costs (50).

Similar trends in the delivery of CBT for insomnia associated with cancer have emerged in recent years, with studies exploring options for providing psychological treatment to individuals with cancer and sleep disorders. The composition of these interventions typically involves a series of treatment sessions, with material that covers the same areas as would be provided in face-to-face treatment. The difference in these cases is that treatment is delivered at distance, either through video-recorded sessions (50), or through on-line sessions (35). In Savard et al's (2014) study, the opportunity for telephone contact with a therapist was available, but somewhat surprisingly, only a limited number (8%) of participants chose to do this.

This study established that video-based self-help materials can have significant benefits in resolving symptoms of insomnia associated with cancer (48), albeit that outcomes for remission rates and relapse rates (for insomnia) were poorer in the video-based treatment group, than a comparison face-to-face group. Ritterberg et al (2012) also found evidence of efficacy in the use of their internet-based therapy, to an extent that matched those found in studies of face-to-face delivered therapy with cancer survivors (35). Although it is acknowledged that aspects of direct treatment (in particular, the therapeutic relationship) may be difficult to recreate in distance methods, the accessibility and cost benefits of such models should not be discounted. Indeed, for many individuals, the only viable treatment option would be through distance methods. Clearly, the counterbalance to this will be individuals' access to the necessary technology, such as internet connections, and the ease of use of programmes. Nonetheless, work to overcome these barriers and explore this additional treatment modality is likely to be a focus of research in future years.

Conclusion

A clear focus on informing the implementation of CBT for insomnia as part of general psychological care within the oncology settings should be a research priority. In current clinical practice, the only help available for the troubling, chronic problem of insomnia has been the attempted short term fix of hypnotic medication, which most cancer patients find particularly undesirable (22). However, CBT for insomnia is an evidence-based solution that is not only safe and effective, but also has the potential to make long term improvements to health and wellbeing in cancer populations.

CBT is an effective and enduring treatment for chronic insomnia disorder, but so far the evidence for CBT has not changed practice. Uncertainty remains about how CBT for insomnia could be integrated into existing oncology care settings, specifically with regard to when it would be most useful for the patient. However, it is suggested that the most substantive barrier to integration of CBT for insomnia into practice is availability. Therapist availability, geographical distance and financial resource implications may limit patient access to treatment.

Recent years have seen the development of several models of treatment delivery for CBT for insomnia. This provides the potential for a stepped-care model, something that may be key to increasing integration of insomnia treatment within wider cancer services. Moving forward, the research focus must be to improve the management of severe and intrusive sleep disturbance that is common in cancer by determining the methods by which it would be most suitable to provide CBT intervention to these patients.

References

1. Stewart BW, Wild CP. World Cancer Report 2014. World Health Organization: Geneva. 2014.
2. Parkin DM. Is the recent fall in incidence of post-menopausal breast cancer in UK related to changes in use of hormone replacement therapy? *Eur J Cancer*. 2009;45(9):1649–53.
3. Davis S, Mirick DK. Circadian disruption, shift work and the risk of cancer: A summary of the evidence and studies in Seattle. *Cancer Causes and Control*. 2006. p. 539–45.
4. Shanmugam V, Wafi A, Al-Taweel N, Büsselberg D. Disruption of circadian rhythm increases the risk of cancer, metabolic syndrome and cardiovascular disease. *J Local Glob Heal Sci [Internet]*. 2013;(2013):3. Available from: <http://www.qscience.com/doi/abs/10.5339/jlghs.2013.3>

5. Siegel R, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. *CA Cancer J Clin* [Internet]. 2014;64(1):9–29. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24399786>
6. Armes J, Crowe M, Colbourne L, Morgan H, Murrells T, Oakley C, et al. Patients' supportive care needs beyond the end of cancer treatment: a prospective, longitudinal survey. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2009.
7. Santin O, Mills M, Treanor C, Donnelly M. A comparative analysis of the health and well-being of cancer survivors to the general population. *Support Care Cancer*. 2012;
8. Association AP. DSM 5 [Internet]. *American Journal of Psychiatry*. 2013. 991 p. Available from: <http://ajp.psychiatryonline.org/article.aspx?articleID=158714>
<http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:DSM-5#0>
9. Palesh OG, Roscoe JA, Mustian KM, Roth T, Savard J, Ancoli-Israel S, et al. Prevalence, demographics, and psychological associations of sleep disruption in patients with cancer: University of Rochester Cancer Center-community clinical oncology program. *J Clin Oncol*. 2010;28:292–8.
10. Fiorentino L, McQuaid JR, Liu L, Natarajan L, He F, Cornejo M, et al. Individual cognitive behavioral therapy for insomnia in breast cancer survivors: a randomized controlled crossover pilot study. *Nat Sci Sleep* [Internet]. 2009;2010:1–8. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2953254&tool=pmcentrez&rendertype=abstract>
11. Garland SN, Johnson JA, Savard J, Gehrman P, Perlis M, Carlson L, et al. Sleeping well with cancer: A systematic review of cognitive behavioral therapy for insomnia in cancer patients. *Neuropsychiatric Disease and Treatment*. 2014. p. 1113–23.
12. Espie CA, Kyle SD, Hames P, Gardani M, Fleming L, Cape J. The Sleep Condition Indicator: a clinical screening tool to evaluate insomnia disorder. *BMJ Open* [Internet]. 2014;4:e004183. Available from: <http://bmjopen.bmj.com/content/4/3/e004183.short>
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3964344&tool=pmcentrez&rendertype=abstract>
13. Savard J, Morin CM. Insomnia in the context of cancer: A review of a neglected problem. *Journal of Clinical Oncology*. 2001. p. 895–908.
14. Savard J, Ivers H, Villa J, Caplette-Gingras A, Morin CM. Natural course of insomnia comorbid with cancer: An 18-month longitudinal study. *J Clin Oncol*. 2011;29:3580–6.

15. Savard J, Savard MH. Insomnia and cancer: Prevalence, nature, and nonpharmacologic treatment. *Sleep Medicine Clinics*. 2013. p. 373–87.
16. Savard J, Savard MH. Insomnia and cancer: Prevalence, nature, and nonpharmacologic treatment. *Sleep Medicine Clinics*. 2013. p. 373–87.
17. Miakowski C, Lee K, Dunn L, Dodd M, Aouizerat BE, West C. Sleep wake circadian activity rhythm parameters and fatigue in oncology patients prior to initiation of radiation therapy. *Cancer Nurs*. 2011;34(4):255–68.
18. Ancoli-Israel S, Liu L, Marler MR, Parker BA, Jones V, Sadler GR, et al. Fatigue, sleep, and circadian rhythms prior to chemotherapy for breast cancer. *Support Care Cancer* [Internet]. 2006;14(3):201–9. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16010529
19. Savard M-H, Savard J, Trudel-Fitzgerald C, Ivers H, Quesnel C. Changes in self-reported hot flashes and their association with concurrent changes in insomnia symptoms among women with breast cancer. *Menopause*. 2011;18(9):985–93.
20. Espie CA, Fleming L, Cassidy J, Samuel L, Taylor LM, White CA, et al. Randomized controlled clinical effectiveness trial of cognitive behavior therapy compared with treatment as usual for persistent insomnia in patients with cancer. *J Clin Oncol* [Internet]. 2008;26:4651–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18591549>
21. Davidson JR, MacLean AW, Brundage MD, Schulze K. Sleep disturbance in cancer patients. *Soc Sci Med*. 2002;
22. Fleming L, Gillespie S, Espie CA. The development and impact of insomnia on cancer survivors: A qualitative analysis. *Psychooncology*. 2010;19(9):991–6.
23. Lis CG, Gupta D, Grutsch JF. The relationship between insomnia and patient satisfaction with quality of life in cancer. *Support Care Cancer*. 2008;16(3):261–6.
24. Mormont MC, Waterhouse J, Bleuzen P, Giacchetti S, Jami A, Bogdan A, et al. Marked 24-h rest/activity rhythms are associated with better quality of life, better response, and longer survival in patients with metastatic colorectal cancer and good performance status. *Clin Cancer Res*. 2000;6(8):3038–45.
25. Mormont MC, Waterhouse J. Contribution of the rest-activity circadian rhythm to quality of life in cancer patients. *Chronobiol Int*. 2002;19(1):313–23.
26. Liu L, Fiorentino L, Rissling M, Natarajan L, Parker BA, Dimsdale JE, et al. Decreased health-related quality of life in women with breast cancer is associated with poor sleep. *Behav Sleep Med* [Internet]. 2013;11(3):189–206. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23205513>

27. Riemann D, Voderholzer U. Primary insomnia: A risk factor to develop depression? *J Affect Disord.* 2003;76(1-3):255–9.
28. Sharma N, Hansen CH, OConnor M, Thekkumpurath P, Walker J, Kleiboer A, et al. Sleep problems in cancer patients: Prevalence and association with distress and pain. *Psychooncology.* 2012;21(9):1003–9.
29. Fleming L, Randell K, Harvey CJ, Espie CA. Does cognitive behaviour therapy for insomnia reduce clinical levels of fatigue, anxiety and depression in cancer patients? *Psychooncology.* 2014;23(6):679–84.
30. Malone M, Harris AL, Luscombe DK. Assessment of the impact of cancer on work, recreation, home management and sleep using a general health status measure. *J R Soc Med.* 1994;87(7):386–9.
31. Redd WH, Silberfarb PM, Andersen BL, Andrykowski MA, Bovbjerg DH, Burish TG, et al. Physiologic and psychobehavioral research in oncology. *Cancer.* 1991;67(3 Suppl):813–22.
32. Morin CM, Gaulier B, Barry T, Kowatch RA. Patients' acceptance of psychological and pharmacological therapies for insomnia. *Sleep.* 1992;15(4):302–5.
33. Savard J, Simard S, Ivers H, Morin CM. Randomized study on the efficacy of cognitive-behavioral therapy for insomnia secondary to breast cancer, part II: Immunologic effects. *J Clin Oncol.* 2005;23:6097–106.
34. Davidson JR, Waisberg JL, Brundage MD, MacLean AW. Nonpharmacologic group treatment of insomnia: a preliminary study with cancer survivors. *Psychooncology [Internet].* 2001;10:389–97. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11536417>
35. Berger AM, Kuhn BR, Farr LA, Von Essen SG, Chamberlain J, Lynch JC, et al. One-year outcomes of a behavioral therapy intervention trial on sleep quality and cancer-related fatigue. *J Clin Oncol.* 2009;27(35):6033–40.
36. Berger AM, Kuhn BR, Farr LA, Lynch JC, Agrawal S, Chamberlain J, et al. Behavioral therapy intervention trial to improve sleep quality and cancer-related fatigue. *Psychooncology.* 2009;18(6):634–46.
37. Fiorentino L, McQuaid JR, Liu L, Natarajan L, He F, Cornejo M, et al. Individual cognitive behavioral therapy for insomnia in breast cancer survivors: A randomized controlled crossover pilot study. *Nat Sci Sleep [Internet].* 2010;2:1–8. Available from: <http://www.scopus.com/inward/record.url?eid=2-s2.0-80053924244&partnerID=40&md5=e044b721a67963cf619c4f48fc3a596f>
38. Ritterband LM, Bailey ET, Thorndike FP, Lord HR, Farrell-Carnahan L, Baum LD. Initial evaluation of an Internet intervention to improve the sleep of cancer survivors with insomnia. *Psychooncology.* 2012;21(7):695–705.

39. Epstein DR, Dirksen SR. Randomized trial of a cognitive-behavioral intervention for insomnia in breast cancer survivors. *Oncol Nurs Forum*. 2007;34(5):E51–9.
40. Savard J, Simard S, Ivers H, Morin CM. Randomized study on the efficacy of cognitive-behavioural therapy for insomnia secondary to breast cancer, part I: Sleep and psychological effects. *J Clin Oncol* [Internet]. 2005;23:6083–96. Available from: <Go to ISI>://000231606300033
41. Jansson M, Linton SJ. Cognitive-behavioral group therapy as an early intervention for insomnia: A randomized controlled trial. *J Occup Rehabil*. 2005;15(2):177–90.
42. Verbeek IH, Konings GM, Aldenkamp AP, Declerck AC, Klip EC. Cognitive behavioral treatment in clinically referred chronic insomniacs: group versus individual treatment. *Behavioral sleep medicine*. 2006.
43. Bastien CH, Morin CM, Ouellet M-C, Blais FC, Bouchard S. Cognitive-behavioral therapy for insomnia: comparison of individual therapy, group therapy, and telephone consultations. *Journal of consulting and clinical psychology*. 2004.
44. Espie C, Kyle S, Williams C, Ong J, Douglas N, Hames P, et al. A Randomized, Placebo-Controlled Trial of Online Cognitive Behavioral Therapy for Chronic Insomnia Disorder Delivered via an Automated Media-Rich Web Application. *Sleep*. 2013;35:769–81.
45. Vincent N, Lewycky S. Logging on for better sleep: RCT of the effectiveness of online treatment for insomnia. *Sleep*. 2009;32(6):807–15.
46. Mimeault V, Morin CM. Self-help treatment for insomnia: bibliotherapy with and without professional guidance. *Journal of consulting and clinical psychology*. 1999.
47. Matthews EE, Berger AM, Schmiede SJ, Cook PF, McCarthy MS, Moore CM, et al. Cognitive Behavioral Therapy for Insomnia Outcomes in Women After Primary Breast Cancer Treatment: A Randomized, Controlled Trial. *Oncol Nurs Forum* [Internet]. 2014;41(3):A1–13. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24650832>
48. Savard J, Ivers H, Savard M-H, Morin CM. Is a video-based cognitive behavioral therapy for insomnia as efficacious as a professionally administered treatment in breast cancer? Results of a randomized controlled trial. *Sleep* [Internet]. 2014;37(8):1305–14. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25083010>
49. Garland SN, Carlson LE, Stephens AJ, Antle MC, Samuels C, Campbell TS. Mindfulness-based stress reduction compared with cognitive behavioral therapy for the treatment of insomnia comorbid with cancer: A randomized, partially blinded, noninferiority trial. *J Clin Oncol*. 2014;32(5):449–57.

50. Espie CA. "Stepped care": a health technology solution for delivering cognitive behavioral therapy as a first line insomnia treatment. *Sleep*. 2009;32(12):1549–58.
51. Brooks J, Wilson K, Amir Z. Additional financial costs borne by cancer patients: A narrative review. *Eur J Oncol Nurs*. 2011;15(4):302–10.
52. Moss-Morris R, Dennison L, Chalder T, Landau S. A randomised controlled trial (RCT) of cognitive behavioural therapy for adjusting to multiple sclerosis. Does it work and for whom does it work? *Psychosom Med* [Internet]. 2011;73(3). Available from: Available from Highwire Press in http://link.worldcat.org/?rft.institution_id=130125&spage=A119&pkgName=freeart&issn=0033-3174&linkclass=to_article&jKey=www.psychosomaticmedicine.org&issue=3&provider=highwire&date=2011-04&aulast=Moss-Morris+R.&atitle=A+ra
53. Espie CA, MacMahon KMA, Kelly H-L, Broomfield NM, Douglas NJ, Engleman HM, et al. Randomized clinical effectiveness trial of nurse-administered small-group cognitive behavior therapy for persistent insomnia in general practice. *Sleep*. 2007;30(5):574–84.
54. Andersson G, Titov N. Advantages and limitations of Internet-based interventions for common mental disorders. *World Psychiatry*. 2014;13(1):4–11.
55. Northouse L, Schafenacker A, Barr KLC, Katapodi M, Yoon H, Brittain K, et al. A Tailored Web-Based Psychoeducational Intervention for Cancer Patients and Their Family Caregivers. *Cancer Nurs* [Internet]. 2014; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24945270>

Key References

- **Northouse, L., Schafenacker, A., K.L.C., B., Katapodi, M., Yoon, H., Brittain, K., et al. A Tailored Web-Based Psychoeducational Intervention for Cancer Patients and Their Family Caregivers. *Cancer Nursing*, 37(5), 321 - 330.**

The authors describe a novel internet-based approach to psycho-education for cancer survivors and family members. Tailored materials were developed and presented via the internet for family dyads (cancer survivor and significant other) to complete. The intervention was generally well-received and effective.

- **Ritterband LM, Bailey ET, Thorndike FP, Lord HR, Farrell-Carnahan L, Baum LD. Initial evaluation of an Internet intervention to improve the sleep of cancer survivors with insomnia. *Psychooncology*, 2012;21(7):695–705.**

This paper describes a wait-list control study of internet-based CBT-I for cancer survivors with insomnia. The intervention significantly reduced sleep

difficulties, with effect sizes similar to those seen in face-to-face interventions. The authors highlight the potential future role of internet-based therapies.

- **Savard, J., Ivers, H., Savard, M. H., & Morin, C. M. (2014). Is a Video-Based Cognitive Behavioral Therapy for Insomnia as Efficacious as a Professionally Administered Treatment in Breast Cancer? Results of a Randomized Controlled Trial. *Sleep*, 37(5).**

The authors present a three-group randomized controlled trial of CBT for insomnia, comparing face-to-face delivery, with video-based therapy and a treatment-as-usual group. Positive effects were found in both the video and face-to-face groups, although gains were more substantial and sustained in the latter group.