

Peter W. Tinning¹, Aimee J.P.M. Franssen², Shehla U. Hridi², Trevor J. Bushell² and Gail McConnell¹

¹ Centre for Biophotonics, Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, United Kingdom.

² Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, United Kingdom.

Introduction

Cytosolic Ca²⁺ plays an integral role in cells and the study of its dynamics can reveal much about biological processes [1]. Fura-2 can provide quantitative data on cytosolic Ca²⁺ changes by exciting at 340 nm and 380 nm and taking the ratio of the emission at both wavelengths [2].

Traditionally for this type of imaging an arc lamp had to be used for illumination as LEDs of the appropriate wavelengths were not available [3]. LEDs hold advantages over arc lamps by exhibiting high amplitude stability and the ability to rapidly switch between wavelengths. We aimed to test a new 340/380 nm LED system for use in ratiometric Fura-2 AM Ca²⁺ imaging and present results using tsA-201 cells and hippocampal neurons.

Methods

- Specimens were washed three times with HEPES-buffered saline solution and loaded with Fura-2 AM for 60 minutes at 37 °C. They were then washed a further three times before imaging.
- HBS Control solution:** - 1 litre of distilled water containing (in mM): NaCl, 140; KCl, 5; MgCl₂, 2; HEPES, 10; D-glucose, 10; CaCl₂, 2 at a pH of 7.4.
- Power at the specimen plane:** - 340 nm: 1.35 mW
380 nm: 1.40 – 3.08 mW
- LED exposure time:** - 0.5 Hz imaging: 100 ms
24.39 Hz video-rate imaging: 20.5 ms
- Emission detection:** - Hamamatsu ORCA-Flash 4.0 CMOS camera with a binning n = 2




Figure 1: Schematic diagram of Olympus BX50 microscope and imaging apparatus

0.5 Hz ratiometric Fura-2 AM Ca²⁺ imaging of drug-mediated responses in tsA-201 cells and hippocampal neurons




Figure 2: A) 340 and 380 images of 1 μM Fura-2 AM loaded tsA-201 cells
Figure 2: B) 340 and 380 images of 1 μM Fura-2 AM loaded hippocampal neurons




Figure 3: A) Representative traces of pharmacologically induced Ca²⁺ concentration changes in tsA-201 cells and C) hippocampal neurons. B) Average Ca²⁺ increase for each stimuli in tsA-201 cells (n = 572) and D) hippocampal neuron (n = 388). Ca²⁺ increases are in agreement with previous experiments in the same cells illuminated by arc lamps [4 - 8]

Average fluctuations in basal Ca²⁺ levels




Figure 4: Example plot of baseline Ca²⁺ fluctuations in hippocampal neurons

The fluctuations are below the theoretical 5 – 10 nM precision of Fura-2 [9]

Ca²⁺ imaging of tsA-201s loaded with lower concentrations of Fura-2AM




Figure 5: Ca²⁺ responses obtained from tsA-201 cells loaded with different concentrations of Fura-2 AM

No significant change in obtained trypsin (100 nM) Ca²⁺ response

Advantages of using lower concentrations:

- Increasing the number of uses from the vial
- Reduced cost
- Improving cell viability
- Cells live longer

Fura-2 ratiometric imaging of synaptically-driven Ca²⁺ events in hippocampal neurons




Figure 6: Synaptically-driven Ca²⁺ events captured at a rate of A) 0.5 Hz in a single neuron and B) 24.39 Hz in two neurons

Conclusions

- This is the first application of a truly 340/380 nm LED illuminator for Fura-2 ratiometric Ca²⁺ imaging of live cell specimens with a precision that is only limited by the response of Fura-2.
- Using this illuminator it is now possible to use Fura-2 AM dye concentrations as low as 250 nM offering both an economical and cell viability advantage.
- Video rate imaging of synaptically-driven Ca²⁺ events combines high temporal and spatial resolution to obtain higher throughput information than previously possible [10].
- This LED illuminator combines optimum excitation and high stability wavelength switching to free Fura-2 imaging from illumination problems experienced in the past.