

Upper limits on gravitational wave emission from 78 radio pulsars

B. Abbott,¹⁴ R. Abbott,¹⁴ R. Adhikari,¹⁴ J. Agresti,¹⁴ P. Ajith,² B. Allen,⁵⁰ R. Amin,¹⁸ S. B. Anderson,¹⁴ W. G. Anderson,⁵⁰ M. Arain,³⁹ M. Araya,¹⁴ H. Armandula,¹⁴ M. Ashley,⁴ S. Aston,³⁸ P. Aufmuth,³⁵ C. Aulbert,¹ S. Babak,¹ S. Ballmer,¹⁴ H. Bantilan,⁸ B. C. Barish,¹⁴ C. Barker,¹⁶ D. Barker,¹⁶ B. Barr,⁴⁰ P. Barriga,⁴⁹ M. Barton,⁴⁰ K. Bayer,¹⁵ J. Betzwieser,¹⁵ P. T. Beyersdorf,²⁷ B. Bhawal,^{14, a} I. A. Bilenko,²¹ G. Billingsley,¹⁴ R. Biswas,⁵⁰ E. Black,¹⁴ K. Blackburn,¹⁴ L. Blackburn,¹⁵ D. Blair,⁴⁹ B. Bland,¹⁶ J. Bogenstahl,⁴⁰ L. Bogue,¹⁷ R. Bork,¹⁴ V. Boschi,¹⁴ S. Bose,⁵¹ P. R. Brady,⁵⁰ V. B. Braginsky,²¹ J. E. Brau,⁴³ M. Brinkmann,² A. Brooks,³⁷ D. A. Brown,¹⁴ A. Bullington,³⁰ A. Bunkowski,² A. Buonanno,⁴¹ O. Burmeister,² D. Busby,¹⁴ R. L. Byer,³⁰ L. Cadonati,¹⁵ G. Cagnoli,⁴⁰ J. B. Camp,²² J. Cannizzo,²² K. Cannon,⁵⁰ C. A. Cantley,⁴⁰ J. Cao,¹⁵ L. Cardenas,¹⁴ G. Castaldi,⁴⁶ C. Cepeda,¹⁴ E. Chalkey,⁴⁰ P. Charlton,⁹ S. Chatterji,¹⁴ S. Chelkowski,² Y. Chen,¹ F. Chiadini,⁴⁵ N. Christensen,⁸ J. Clark,⁴⁰ P. Cochrane,² T. Cokelaer,⁷ R. Coldwell,³⁹ R. Conte,⁴⁵ D. Cook,¹⁶ T. Corbitt,¹⁵ D. Coyne,¹⁴ J. D. E. Creighton,⁵⁰ R. P. Croce,⁴⁶ D. R. M. Crooks,⁴⁰ A. M. Cruise,³⁸ A. Cumming,⁴⁰ J. Dalrymple,³¹ E. D'Ambrosio,¹⁴ K. Danzmann,^{35, 2} G. Davies,⁷ D. DeBra,³⁰ J. Degallaix,⁴⁹ M. Degree,³⁰ T. Demma,⁴⁶ V. Dergachev,⁴² S. Desai,³² R. DeSalvo,¹⁴ S. Dhurandar,¹³ M. Díaz,³³ J. Dickson,⁴ A. Di Credico,³¹ G. Diederichs,³⁵ A. Dietz,⁷ E. E. Doomes,²⁹ R. W. P. Drever,⁵ J.-C. Dumas,⁴⁹ R. J. Dupuis,¹⁴ J. G. Dwyer,¹⁰ P. Ehrens,¹⁴ E. Espinoza,¹⁴ T. Etzel,¹⁴ M. Evans,¹⁴ T. Evans,¹⁷ S. Fairhurst,^{7, 14} Y. Fan,⁴⁹ D. Fazi,¹⁴ M. M. Fejer,³⁰ V. Fiumara,⁴⁵ L. S. Finn,³² N. Fotopoulos,⁵⁰ A. Franzen,³⁵ K. Y. Franzen,^{39, b} A. Freise,³⁸ R. E. Frey,⁴³ T. Fricke,⁴⁴ P. Fritschel,¹⁵ V. V. Frolov,¹⁷ M. Fyffe,¹⁷ V. Galdi,⁴⁶ J. Garofoli,¹⁶ I. Gholami,¹ J. A. Giaime,^{17, 18} S. Giampanis,⁴⁴ K. D. Giardino,¹⁷ K. Goda,¹⁵ E. Goetz,⁴² L. Goggin,¹⁴ G. González,¹⁸ S. Gossler,⁴ A. Grant,⁴⁰ S. Gras,⁴⁹ C. Gray,¹⁶ M. Gray,⁴ J. Greenhalgh,²⁶ A. M. Gretarsson,¹¹ R. Grosso,^{33, c} H. Grote,² S. Grunewald,¹ M. Guenther,¹⁶ R. Gustafson,⁴² B. Hage,³⁵ D. Hammer,⁵⁰ C. Hanna,¹⁸ J. Hanson,¹⁷ J. Harms,² G. Harry,¹⁵ E. Harstad,⁴³ T. Hayler,²⁶ J. Heefner,¹⁴ I. S. Heng,⁴⁰ A. Heptonstall,⁴⁰ M. Heurs,² M. Hewitson,² S. Hild,³⁵ E. Hirose,³¹ D. Hoak,¹⁷ D. Hosken,³⁷ J. Hough,⁴⁰ D. Hoyland,³⁸ S. H. Huttner,⁴⁰ D. Ingram,¹⁶ E. Innerhofer,¹⁵ M. Ito,⁴³ Y. Itoh,⁵⁰ A. Ivanov,¹⁴ B. Johnson,¹⁶ W. W. Johnson,¹⁸ D. I. Jones,^{40, d} G. Jones,⁷ R. Jones,⁴⁰ L. Ju,⁴⁹ P. Kalmus,¹⁰ V. Kalogera,²⁴ D. Kasprzyk,³⁸ E. Katsavounidis,¹⁵ K. Kawabe,¹⁶ S. Kawamura,²³ F. Kawazoe,²³ W. Kells,¹⁴ D. G. Keppel,¹⁴ F. Ya. Khalili,²¹ C. Kim,²⁴ P. King,¹⁴ J. S. Kissel,¹⁸ S. Klimentko,³⁹ K. Kokeyama,²³ V. Kondrashov,¹⁴ R. K. Kopparapu,¹⁸ D. Kozak,¹⁴ B. Krishnan,¹ P. Kwee,³⁵ P. K. Lam,⁴ M. Landry,¹⁶ B. Lantz,³⁰ A. Lazzarini,¹⁴ M. Lei,¹⁴ J. Leiner,⁵¹ V. Leonhardt,²³ I. Leonor,⁴³ K. Libbrecht,¹⁴ P. Lindquist,¹⁴ N. A. Lockerbie,⁴⁷ M. Longo,⁴⁵ M. Lormand,¹⁷ M. Lubinski,¹⁶ H. Lück,^{35, 2} B. Machenschalk,¹ M. MacInnis,¹⁵ M. Mageswaran,¹⁴ K. Mailand,¹⁴ M. Malec,^{35, e} V. Mandic,¹⁴ S. Marano,⁴⁵ S. Márka,¹⁰ J. Markowitz,¹⁵ E. Maros,¹⁴ I. Martin,⁴⁰ J. N. Marx,¹⁴ K. Mason,¹⁵ L. Matone,¹⁰ V. Matta,⁴⁵ N. Mavalvala,¹⁵ R. McCarthy,¹⁶ D. E. McClelland,⁴ S. C. McGuire,²⁹ M. McHugh,²⁰ K. McKenzie,⁴ S. McWilliams,²² T. Meier,³⁵ A. Melissinos,⁴⁴ G. Mendell,¹⁶ R. A. Mercer,³⁹ S. Meshkov,¹⁴ E. Messaritaki,¹⁴ C. J. Messenger,⁴⁰ D. Meyers,¹⁴ E. Mikhailov,¹⁵ S. Mitra,¹³ V. P. Mitrofanov,²¹ G. Mitselmakher,³⁹ R. Mittleman,¹⁵ O. Miyakawa,¹⁴ S. Mohanty,³³ G. Moreno,¹⁶ K. Mossavi,² C. MowLowry,⁴ A. Moylan,⁴ G. Mueller,³⁹ S. Mukherjee,³³ H. Müller-Ebhardt,² J. Munch,³⁷ P. Murray,⁴⁰ E. Myers,¹⁶ J. Myers,¹⁶ G. Newton,⁴⁰ A. Nishizawa,²³ K. Numata,²² B. O'Reilly,¹⁷ R. O'Shaughnessy,²⁴ D. J. Ottaway,¹⁵ H. Overmier,¹⁷ B. J. Owen,³² Y. Pan,⁴¹ M. A. Papa,^{1, 50} V. Parameshwaraiah,¹⁶ P. Patel,¹⁴ M. Pedraza,¹⁴ S. Penn,¹² V. Pierro,⁴⁶ I. M. Pinto,⁴⁶ M. Pitkin,⁴⁰ H. Pletsch,⁵⁰ M. V. Plissi,⁴⁰ F. Postiglione,⁴⁵ R. Prix,¹ V. Quetschke,³⁹ F. Raab,¹⁶ D. Rabeling,⁴ H. Radkins,¹⁶ R. Rahkola,⁴³ N. Rainer,² M. Rakhmanov,³² S. Ray-Majumder,⁵⁰ V. Re,^{38, f} H. Rehbein,² S. Reid,⁴⁰ D. H. Reitze,³⁹ L. Ribichini,² R. Riesen,¹⁷ K. Riles,⁴² B. Rivera,¹⁶ N. A. Robertson,^{14, 40} C. Robinson,⁷ E. L. Robinson,³⁸ S. Roddy,¹⁷ A. Rodriguez,¹⁸ A. M. Rogan,⁵¹ J. Rollins,¹⁰ J. D. Romano,⁷ J. Romie,¹⁷ R. Route,³⁰ S. Rowan,⁴⁰ A. Rüdiger,² L. Ruet,¹⁵ P. Russell,¹⁴ K. Ryan,¹⁶ S. Sakata,²³ M. Samidi,¹⁴ L. Sancho de la Jordana,³⁶ V. Sandberg,¹⁶ V. Sannibale,¹⁴ S. Saraf,²⁵ P. Sarin,¹⁵ B. S. Sathyaprakash,⁷ S. Sato,²³ P. R. Saulson,³¹ R. Savage,¹⁶ P. Savov,⁶ S. Schediwy,⁴⁹ R. Schilling,² R. Schnabel,² R. Schofield,⁴³ B. F. Schutz,^{1, 7} P. Schwinberg,¹⁶ S. M. Scott,⁴ A. C. Searle,⁴ B. Sears,¹⁴ F. Seifert,² D. Sellers,¹⁷ A. S. Sengupta,⁷ P. Shawhan,⁴¹ D. H. Shoemaker,¹⁵ A. Sibley,¹⁷ J. A. Sidles,⁴⁸ X. Siemens,¹⁴ D. Sigg,¹⁶ S. Sinha,³⁰ A. M. Sintes,^{36, 1} B. Slagmolen,⁴ J. Slutsky,¹⁸ J. R. Smith,² M. R. Smith,¹⁴ K. Somiya,^{2, 1} K. A. Strain,⁴⁰ D. M. Strom,⁴³ A. Stuver,³² T. Z. Summerscales,³ K.-X. Sun,³⁰ M. Sung,¹⁸ P. J. Sutton,¹⁴ H. Takahashi,¹ D. B. Tanner,³⁹ R. Taylor,¹⁴ R. Taylor,⁴⁰ J. Thacker,¹⁷ K. A. Thorne,³² K. S. Thorne,⁶ A. Thüring,³⁵ K. V. Tokmakov,⁴⁰ C. Torres,³³ C. Torrie,⁴⁰ G. Traylor,¹⁷ M. Trias,³⁶ W. Tyler,¹⁴ D. Ugolini,³⁴ K. Urbanek,³⁰ H. Vahlbruch,³⁵ M. Vallisneri,⁶ C. Van Den Broeck,⁷ M. Varvella,^{14, g} S. Vass,¹⁴ A. Vecchio,³⁸ J. Veitch,⁴⁰ P. Veitch,³⁷ A. Villar,¹⁴ C. Vorvick,¹⁶ S. P. Vyachanin,²¹ S. J. Waldman,¹⁴ L. Wallace,¹⁴ H. Ward,⁴⁰ R. Ward,¹⁴ K. Watts,¹⁷ A. Weidner,² M. Weinert,² A. Weinstein,¹⁴ R. Weiss,¹⁵ S. Wen,¹⁸ K. Wette,⁴

J. T. Whelan,¹ S. E. Whitcomb,¹⁴ B. F. Whiting,³⁹ C. Wilkinson,¹⁶ P. A. Willems,¹⁴ L. Williams,³⁹
B. Willke,^{35,2} I. Wilmot,²⁶ W. Winkler,² C. C. Wipf,¹⁵ S. Wise,³⁹ A. G. Wiseman,⁵⁰ G. Woan,⁴⁰ D. Woods,⁵⁰
R. Wooley,¹⁷ J. Worden,¹⁶ W. Wu,³⁹ I. Yakushin,¹⁷ H. Yamamoto,¹⁴ Z. Yan,⁴⁹ S. Yoshida,²⁸ N. Yunes,³²
M. Zanolin,¹⁵ J. Zhang,⁴² L. Zhang,¹⁴ C. Zhao,⁴⁹ N. Zotov,¹⁹ M. Zucker,¹⁵ H. zur Mühlen,³⁵ and J. Zweizig¹⁴
(The LIGO Scientific Collaboration, <http://www.ligo.org>)

M. Kramer⁵² and A. G. Lyne⁵²

- ¹Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-14476 Golm, Germany
²Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-30167 Hannover, Germany
³Andrews University, Berrien Springs, MI 49104 USA
⁴Australian National University, Canberra, 0200, Australia
⁵California Institute of Technology, Pasadena, CA 91125, USA
⁶Caltech-CaRT, Pasadena, CA 91125, USA
⁷Cardiff University, Cardiff, CF2 3YB, United Kingdom
⁸Carleton College, Northfield, MN 55057, USA
⁹Charles Sturt University, Wagga Wagga, NSW 2678, Australia
¹⁰Columbia University, New York, NY 10027, USA
¹¹Embry-Riddle Aeronautical University, Prescott, AZ 86301 USA
¹²Hobart and William Smith Colleges, Geneva, NY 14456, USA
¹³Inter-University Centre for Astronomy and Astrophysics, Pune - 411007, India
¹⁴LIGO - California Institute of Technology, Pasadena, CA 91125, USA
¹⁵LIGO - Massachusetts Institute of Technology, Cambridge, MA 02139, USA
¹⁶LIGO Hanford Observatory, Richland, WA 99352, USA
¹⁷LIGO Livingston Observatory, Livingston, LA 70754, USA
¹⁸Louisiana State University, Baton Rouge, LA 70803, USA
¹⁹Louisiana Tech University, Ruston, LA 71272, USA
²⁰Loyola University, New Orleans, LA 70118, USA
²¹Moscow State University, Moscow, 119992, Russia
²²NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA
²³National Astronomical Observatory of Japan, Tokyo 181-8588, Japan
²⁴Northwestern University, Evanston, IL 60208, USA
²⁵Rochester Institute of Technology, Rochester, NY 14623, USA
²⁶Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX United Kingdom
²⁷San Jose State University, San Jose, CA 95192, USA
²⁸Southeastern Louisiana University, Hammond, LA 70402, USA
²⁹Southern University and A&M College, Baton Rouge, LA 70813, USA
³⁰Stanford University, Stanford, CA 94305, USA
³¹Syracuse University, Syracuse, NY 13244, USA
³²The Pennsylvania State University, University Park, PA 16802, USA
³³The University of Texas at Brownsville and Texas Southmost College, Brownsville, TX 78520, USA
³⁴Trinity University, San Antonio, TX 78212, USA
³⁵Universität Hannover, D-30167 Hannover, Germany
³⁶Universitat de les Illes Balears, E-07122 Palma de Mallorca, Spain
³⁷University of Adelaide, Adelaide, SA 5005, Australia
³⁸University of Birmingham, Birmingham, B15 2TT, United Kingdom
³⁹University of Florida, Gainesville, FL 32611, USA
⁴⁰University of Glasgow, Glasgow, G12 8QQ, United Kingdom
⁴¹University of Maryland, College Park, MD 20742 USA
⁴²University of Michigan, Ann Arbor, MI 48109, USA
⁴³University of Oregon, Eugene, OR 97403, USA
⁴⁴University of Rochester, Rochester, NY 14627, USA
⁴⁵University of Salerno, 84084 Fisciano (Salerno), Italy
⁴⁶University of Sannio at Benevento, I-82100 Benevento, Italy
⁴⁷University of Strathclyde, Glasgow, G1 1XQ, United Kingdom
⁴⁸University of Washington, Seattle, WA, 98195
⁴⁹University of Western Australia, Crawley, WA 6009, Australia
⁵⁰University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA
⁵¹Washington State University, Pullman, WA 99164, USA
⁵²University of Manchester, Jodrell Bank Observatory, Macclesfield, Cheshire, SK11 9DL, United Kingdom
- (Dated: February 8, 2007)

We present upper limits on the gravitational wave emission from 78 radio pulsars based on data from the third and fourth science runs of the LIGO and GEO 600 gravitational wave detectors. The data from both runs have been combined coherently to maximise sensitivity. For the first time pulsars within binary (or multiple) systems have been included in the search by taking into account the signal modulation due to their orbits. Our upper limits are therefore the first measured for 56 of these pulsars. For the remaining 22, our results improve on previous upper limits by up to a factor of 10. For example, our tightest upper limit on the gravitational strain is 3.2×10^{-25} for PSR J1603–7202, and the equatorial ellipticity of PSR J2124–3358 is less than 10^{-6} . Furthermore, our strain upper limit for the Crab pulsar is only three times greater than the fiducial spin-down limit.

PACS numbers: 04.80.Nn, 95.55.Ym, 97.60.Gb, 07.05.Kf

I. INTRODUCTION

This paper details the results of a search for gravitational wave signals from known radio pulsars in data from the third and fourth LIGO and GEO 600 science runs (denoted S3 and S4). These runs were carried out from 31st October 2003 to 9th January 2004 and 22nd February to 23rd March, 2005 respectively. We have applied, and extended, the search technique of Dupuis and Woan [1] to generate upper limits on the gravitational wave amplitude from a selection of known radio pulsars, and infer upper limits on their equatorial ellipticities. The work is a natural extension our previous work given in Refs. [2, 3].

A. Motivation

To emit gravitational waves a pulsar must have some mass (or mass-current) asymmetry around its rotation axis. This can be achieved through several mechanisms such as elastic deformations of the solid crust or core or distortion of the entire star by an extremely strong misaligned magnetic field (see Sec. III of Ref. [4] for a recent review). Such mechanisms generally result in a triaxial neutron star which, in the quadrupole approximation and with rotation and angular momentum axes aligned, would produce gravitational waves at twice the rotation frequency. These waves would have a characteristic strain amplitude at the Earth (assuming optimal orientation of the rotation axis) of

$$h_0 = \frac{16\pi^2 G}{c^4} \frac{\varepsilon I_{zz} \nu^2}{r}, \quad (1.1)$$

where ν is the neutron star's spin frequency, I_{zz} its principal moment of inertia, $\varepsilon = (I_{xx} - I_{yy})/I_{zz}$ its equatorial ellipticity, and r its distance from Earth [5].

Known pulsars provide an enticing target for gravitational wave searches as their positions and frequencies are generally well-known through radio or X-ray observations. As a result the signal search covers a much smaller parameter space than is necessary when searching for signals from unknown sources, giving a lower significance threshold. In addition, the deterministic nature of the waves allows a building up of the signal-to-noise ratio by observing coherently for a considerable time. The main drawback in a search for gravitational waves from the majority of known pulsars is that the level of emission is likely to be lower than can be detected with current detector sensitivities.

Using existing radio measurements, and some reasonable assumptions, it is possible to set an upper limit on the gravitational wave amplitude from a pulsar based purely on energy conservation arguments. If one assumes that the pulsar is an isolated rigid body and that the observed spin-down of the pulsar is due to the loss of rotational kinetic energy as gravitational radiation (i.e., $dE_{\text{rot}}/dt = 4\pi^2 I_{zz} \nu \dot{\nu}$) then the gravitational wave amplitude at the Earth (assuming optimal orientation of the rotation axis) would be

$$h_{\text{sd}} = \left(\frac{5 G I_{zz} |\dot{\nu}|}{2 c^3 r^2 \nu} \right)^{1/2}. \quad (1.2)$$

Of course these assumptions may not hold, but it would be surprising if neutron stars radiated significantly more gravitational energy than this. With these uncertainties in mind, searches such as the one described in this paper place *direct* upper limits on gravitational wave emission from rotating neutron stars, and these limits are already approaching the regime of astrophysical interest.

B. Previous results

Before the advent of large-scale interferometric detectors, there was only a limited ability to search for gravitational waves from known pulsars. Resonant mass gravitational wave detectors are only sensitive in a relatively narrow band around their resonant frequency and so cannot be used to target objects radiating outside that band. A specific attempt to search for gravitational waves from the Crab pulsar at a frequency of ~ 60 Hz was, however,

^aCurrently at Google, Inc.

^bCurrently at KLA-Tencor

^cCurrently at Universität Erlangen-Nürnberg

^dCurrently at University of Southampton

^eCurrently at LD Didactic, Gundelfingen

^fCurrently at University of Trento

[§]Permanent Address: Laboratoire de l'Accélérateur Linéaire, Orsay

made with a specially designed aluminium quadrupole antenna [6, 7] giving a 1σ upper limit of $h_0 \leq 2 \times 10^{-22}$. A search for gravitational waves from what was then the fastest millisecond pulsar, PSR J1939+2134, was conducted by Hough *et al* [8] using a split bar detector, producing an upper limit of $h_0 < 10^{-20}$.

The first pulsar search using interferometer data was carried out with the prototype 40 m interferometer at Caltech by Hereld [9]. The search was again for gravitational waves from PSR J1939+2134, and produced upper limits of $h_0 < 3.1 \times 10^{-17}$ and $h_0 < 1.5 \times 10^{-17}$ for the first and second harmonics of the pulsar’s rotation frequency.

A much larger sample of pulsars is accessible to broadband interferometers. As of the beginning of 2005 the Australia Telescope National Facility (ATNF) online pulsar catalogue [10] listed¹ 154 millisecond and young pulsars, all with rotation frequencies > 25 Hz (gravitational wave frequency > 50 Hz) that fall within the design band of the LIGO and GEO 600 interferometers, and the search for their gravitational waves has developed rapidly since the start of data-taking runs in 2002. Data from the first science run (S1) were used to perform a search for gravitational waves at twice the rotation frequency from PSR J1939+2134 [2]. Two techniques were used in this search: one a frequency domain, frequentist search, and the other a time domain, Bayesian search which gave a 95% credible amplitude upper limit of 1.4×10^{-22} , and an ellipticity upper limit of 2.9×10^{-4} assuming $I_{zz} = 10^{38}$ kg m².

Analysis of data from the LIGO S2 science run set upper limits on the gravitational wave amplitude from 28 radio pulsars [3]. To do this, new radio timing data were obtained to ensure the pulsars’ rotational phases could be predicted with the necessary accuracy and to check that none of the pulsars had glitched. These data gave strain upper limits as low as a few times 10^{-24} , and several ellipticity upper limits less than 10^{-5} . The Crab pulsar was also studied in this run, giving an upper limit a factor of ~ 30 greater than the spin-down limit considered above. Prior to this article these were the most sensitive studies made. Preliminary results for the same 28 pulsars using S3 data were given in Dupuis (2004) [11], and these are expanded below.

In addition to the above, data from the LIGO S2 run have been used to perform an all-sky (i.e., non-targeted) search for continuous wave signals from isolated sources, and a search for a signal from the neutron star within the binary system Sco-X1 [4]. An all-sky continuous wave search using the distributed computing project Einstein@home² has also been performed on S3 data [14]. These searches use the same search algorithms, are fully coherent and are ongoing using data from more recent

(and therefore more sensitive) runs. Additional continuous wave searches using incoherent techniques are also being performed on LIGO data [15, 16].

Unfortunately the pulsar population is such that most have spin frequencies that fall below the sensitivity band of current detectors. In the future, the low-frequency sensitivity of VIRGO [12] and Advanced LIGO [13] should allow studies of a significantly larger sample of pulsars.

C. The signal

Following convention, we model the observed phase evolution of a pulsar using a Taylor expansion about a fixed epoch time t_0 :

$$\phi(T) = \phi_0 + 2\pi \left\{ \nu_0(T - t_0) + \frac{1}{2} \dot{\nu}_0(T - t_0)^2 + \frac{1}{6} \ddot{\nu}_0(T - t_0)^3 + \dots \right\}, \quad (1.3)$$

where ϕ_0 is the initial (epoch) spin phase, ν_0 and its time derivatives are the pulsar spin frequency and spin-down coefficients at t_0 , and T is the pulsar proper time.

The expected signal in an interferometer from a triaxial pulsar is

$$h(t) = \frac{1}{2} F_+(t; \psi) h_0 (1 + \cos^2 \iota) \cos 2\phi(t) + F_\times(t; \psi) h_0 \cos \iota \sin 2\phi(t), \quad (1.4)$$

where $\phi(t)$ is the phase evolution in the detector time t , F_+ and F_\times are the detector antenna patterns for the plus and cross polarisations of gravitational waves, ψ is the wave polarisation angle, and ι is the angle between the rotation axis of the pulsar and the line-of-sight. A gravitational wave impinging on the interferometer will be modulated by Doppler, time delay and relativistic effects caused by the motions of the Earth and other bodies in the solar system. Therefore we need to transform the ‘arrival time’ of a wave-crest at the detector, t , to its arrival time at the solar system barycentre (SSB) t_b via

$$t_b = t + \delta t = t + \frac{\mathbf{r} \cdot \hat{\mathbf{n}}}{c} + \Delta_{E_\odot} + \Delta_{S_\odot}, \quad (1.5)$$

where \mathbf{r} is the position of the detector with respect to the SSB, $\hat{\mathbf{n}}$ is the unit vector pointing to the pulsar, Δ_{E_\odot} is the special relativistic Einstein delay, and Δ_{S_\odot} is the general relativistic Shapiro delay [17]. Although pulsars can be assumed to have a large velocity with respect to the SSB, it is conventional to ignore this Doppler term and set $t_b = T$, as its proper motion is generally negligible (see §VIA for cases where this assumption is not the case). For pulsars in binary systems, there will be additional time delays due to the binary orbit, discussed in §III B.

¹ The catalogue is continually updated and as such now contains more objects.

² <http://einstein.phys.uwm.edu>

The S3 and S4 runs used all three LIGO interferometers (H1 and H2 at the Hanford Observatory in Washington, and L1 at the Livingston Observatory in Louisiana) in the USA and the GEO 600 interferometer in Hannover, Germany. GEO 600 did not run for all of S3, but had two main data taking periods between which improvements were made to its sensitivity. All these detectors had different duty factors and sensitivities.

A. LIGO

For S3 the H1 and H2 interferometers maintained relatively high duty factors of 69.3% and 63.4% respectively. The L1 interferometer was badly affected by anthropogenic seismic noise sources during the day and thus had a duty factor of only 21.8%.

Between S3 and S4 the L1 interferometer was upgraded with better seismic isolation. This greatly reduced the amount of time the interferometer was thrown out of its operational state by anthropogenic noise, and allowed it to operate successfully during the day, with a duty factor of 74.5% and a longest lock stretch of 18.7h. The H1 and H2 interferometers also both improved their duty factors to 80.5% and 81.4%, with longest lock stretches of almost a day. The typical strain sensitivities of all the interfer-




FIG. 1: Median strain amplitude spectral density curves for the LIGO and GEO 600 interferometers during the S4 run.

ometers during S4 can be seen in Fig. 1. This shows the LIGO detectors reach their best sensitivities at about 150 Hz, whilst GEO 600 achieves its best sensitivity at its tuned frequency of 1 kHz.

During S3 GEO 600 was operated as a dual-recycled Michelson interferometer tuned to have greater sensitivity to signals around 1 kHz. The first period of GEO 600 participation in S3 was between 5th to 11th November 2003, called S3I, during which the detector operated with a 95.1% duty factor. Afterwards, GEO 600 was taken offline to allow further commissioning work aimed at improving sensitivity and stability. Then from 30th December 2003 to 13th January 2004 GEO 600 rejoined S3, called S3II, with an improved duty factor of 98.7% and with more than one order of magnitude improvement in peak sensitivity. During S3 there were five locks of longer than 24 hours and one lock longer than 95 hours. For more information about the performance of GEO 600 during S3 see Ref. [18].

GEO 600 participated in S4 from 22nd February to 24th March 2005, with a duty factor of 96.6%. It was operated in essentially the same optical configuration as in S3. With respect to S3, the sensitivity was improved more than an order of magnitude over a wide frequency range, and close to two orders of magnitude around 100 Hz. For more information about GEO 600 during S4 see Ref. [19].

C. Data quality

When a detector is locked on resonance and all control loops are in their nominal running states and there are no on-site work activities that are known to compromise the data, then the data are said to be *science mode*. All science mode data are not of sufficient quality to be analysed however, and may be flagged for exclusion. Examples of such data quality flags are ones produced for epochs of excess seismic noise, and the flagging of data corrupted by overflows of photodiode ADCs. For this analysis we use all science mode data for which there is no corresponding data quality flag. For S3 this gives observation times of 45.5 days for H1; 42.1 days for H2; and 13.4 days for L1. For S4 this gives observation times of 19.4 days for H1; 22.5 days for H2; and 17.1 days for L1.

III. THE SEARCH METHOD

Our search method involves heterodyning the data using the phase model $\phi(t)$ to precisely unwind the phase evolution of the expected signal, and has been discussed in detail in Ref. [1]. After heterodyning, the data are low-pass filtered, using a ninth order Butterworth filter with a knee frequency of 0.5 Hz, and re-binned from the raw data sample rate of 16384 Hz to 1/60 Hz i.e., one sample per minute. The motion of the detector within the solar system modulates the signal and this is taken into account within the heterodyne by using a time delay given in Eq. (1.5), which transforms the signal to the SSB. Signals from binary pulsar systems have an extra

modulation term in the signal, as discussed briefly below, and these we targeted for the first time in S3/S4.

We infer the pulsar signal parameters, denoted $\mathbf{a} = (h_0, \phi_0, \cos \iota, \psi)$, from their (Bayesian) posterior probability distribution (pdf) over this parameter space, assuming Gaussian noise. The data are broken up into time segments over which the noise can be assumed stationary and we analytically marginalise over the unknown noise floor, giving a Student-t likelihood for the parameters for each segment (see Ref. [1] for the method). Combining the segments gives an overall likelihood of

$$p(\{B_k\}|\mathbf{a}) \propto \prod_j^M \left(\sum_{k=1+\sum_{i=1}^{j-1} m_i}^{\sum_{i=1}^j m_i} |B_k - y_k|^2 \right)^{-m_j}, \quad (3.1)$$

where each B_k is a heterodyned sample with a sample rate of one per minute, M is the number of segments into which the whole data set has been cut, m_j is the number of data points in the j th segment, and y_k , given by

$$y_k = \frac{1}{4} F_+(t_k; \psi) h_0 (1 + \cos^2 \iota) e^{i2\phi_0} - \frac{i}{2} F_\times(t_k; \psi) h_0 \cos \iota e^{i2\phi_0}, \quad (3.2)$$

is the gravitational wave signal model evaluated at t_k , the time corresponding to the k 'th heterodyned sample. In Ref. [3] the value of m_j was fixed at 30 to give 30 min data segments, and data that was contiguous only on shorter timescales, and which could not be fitted into one of these segments, was thrown out. In the analysis presented here we have allowed segment lengths to vary from 5 to 30 min, so we maximise the number of 30-minute segments whilst also allowing shorter segments at the end of locked stretches to contribute. The likelihood in Eq. (3.1) assumes that the data is stationary over each of these 30 minute (or smaller) segments. This assumption holds well for our data. Large outliers can also be identified and vetoed from the data, for example those at the beginning of a data segment caused by the impulsive ringing of the low-pass filter applied after the data is heterodyned.

The prior probabilities for each of the parameters are taken as uniform over their respective ranges. Upper limits on h_0 are set by marginalising the posterior over the nuisance parameters and then calculating the $h_0^{95\%}$ value that bounds the cumulative probability for the desired credible limit of 95%:

$$0.95 = \int_0^{h_0^{95\%}} p(h_0|\{B_k\}) dh_0. \quad (3.3)$$

A. Combining data

In the search of Ref. [3] the combined data from the three LIGO interferometers were used to improve the sensitivity of the search. This was done by forming the joint

likelihood from the three *independent* data sets:

$$p(B_k|\mathbf{a})_{\text{Joint}} = p(B_k|\mathbf{a})_{\text{H1}} \cdot p(B_k|\mathbf{a})_{\text{H2}} \cdot p(B_k|\mathbf{a})_{\text{L1}}. \quad (3.4)$$

This is valid provided the data acquisition is coherent between detectors and supporting evidence for this is presented in §V. It is of course a simple matter to extend Eq. (3.4) to include additional likelihood terms from other detectors, such as GEO 600.

In this analysis we also combine data sets from two different science runs. This is appropriate because S3 and S4 had comparable sensitivities over a large portion of the spectrum. Provided the data sets maintain phase coherence between runs, this combination can simply be achieved by concatenating the data sets from the two runs together for each detector.

An example of the posterior pdfs for the four unknown pulsar parameters of PSR J0024–7204C (each marginalised over the three other parameters) is shown in Fig. 2. The pdfs in Fig. 2 are from the joint analysis




FIG. 2: The marginalised posterior pdfs for the four unknown pulsar parameters h_0 , ϕ_0 , $\cos \iota$ and ψ , for PSR J0024–7204C using the joint data from the three LIGO detectors over S3 and S4.

of the three LIGO detectors using the S3 and S4 data, all combined coherently. The shaded area in the h_0 posterior shows the area containing 95% of the probability as given by Eq. (3.3). In this example the posterior on h_0 is peaked at $h_0 = 0$, though any distribution that is credibly close to zero is consistent with $h_0 = 0$. Indeed an upper limit can formally be set even when the bulk of the probability is well away from zero (see the discussion of hardware injections in §V).

B. Binary models

Our previous known pulsar searches [2, 3] have excluded pulsars within binary systems, despite the majority of pulsars within our detector band being in such systems. To address this, we have included an additional time delay to transform from the binary system barycentre (BSB) to pulsar proper time, which is a stationary reference frame with respect to the pulsar. The code for this is based on the widely used radio pulsar timing software TEMPO [20]. The algorithm and its testing are discussed more thoroughly in Ref. [21].

There are five principal parameters describing a Keplerian orbit: the time of periastron, T_0 , the longitude of periastron, ω_0 , the eccentricity, e , the period, P_b , and the projected semi-major axis, $x = a \sin i$. These describe the majority of orbits very well, although to fully describe the orbit of some pulsars requires additional relativistic parameters. The basic transformation and binary models below are summarised by Taylor and Weisberg [17] and Lange *et al.* [22], and are those used in TEMPO. The transformation from SSB time t_b to pulsar proper time T follows the form of Eq. (1.5) and is

$$t_b = T + \Delta_R + \Delta_E + \Delta_S, \quad (3.5)$$

where Δ_R is the Roemer time delay giving the propagation time across the binary orbit, Δ_E is the Einstein delay and gives gravitational redshift and time dilation corrections, and Δ_S is the Shapiro delay and gives the general relativistic correction (see Ref. [17] for definitions of these delays).

The majority of binary pulsars can be described by three orbital models: the Blandford-Teukolsky (BT) model, the low eccentricity (ELL1) model, and the Damour-Deruelle (DD) model (see Refs. [17, 20, 22] for further details of these models). These different models make different assumptions about the system and/or are specialised to account for certain system features. For example, the ELL1 model is used in cases where the eccentricity is very small, and therefore periastron is very hard to define, in which case the time and longitude of periastron will be highly correlated and have to be reparameterised to the Laplace-Lagrange parameters [22]. When a binary pulsar's parameters are estimated from radio observations using TEMPO the different models are used accordingly. These models can be used within our search to calculate all the associated time delays and therefore correct the signal to the pulsar proper time, provided we have accurate model parameters for the pulsar.

IV. PULSAR SELECTION

The noise floor of the LIGO detectors increases rapidly below about 50 Hz, so pulsar targets were primarily selected on their frequency. The choice of a 50 Hz gravitational wave frequency cut-off (pulsar spin frequency

of 25 Hz) is somewhat arbitrary, but it also loosely reflects the split between the population of fast (millisecond/recycled and young) pulsars and slow pulsars.

All 154 pulsars with spin frequencies > 25 Hz were taken from the ATNF online pulsar catalogue [10] (described in Ref. [23]). The accuracy of these parameters varies for each pulsar and is dependent on the time span, density of observations and the noise level of the timing observations. Clearly it is important to ensure that parameter uncertainties do not lead to unacceptable phase errors in the heterodyne. Pulsars are not perfect clocks, so the epoch of the parameters is also important as more recent measurements will better reflect the current state of the pulsar. Importantly, there is near-continuous monitoring of the Crab pulsar at Jodrell Bank Observatory, and as such its parameters are continuously updated [24].

Precise knowledge of the phase evolution of each target pulsar is vital for our analysis, and possible effects that may lead to a departure from the simple second-order Taylor expansion are discussed below.

A. Pulsar timing

Using TEMPO, we obtained the parameters of 75 pulsars from the regular observation programs carried out at Jodrell Bank Observatory and the Parkes Telescope (see Ref. [25] for details of the techniques used for this). For 37 of these the timings spanned the period of S3. These same model parameters were used to extrapolate the pulsar phases to the period of S4. The effect of parameter uncertainties on this extrapolation is discussed in §IV B, but is only important in its effect on the extrapolated phase. For those pulsars observed during S3 the interpolation is taken to be free from significant error.

The parameters for 16 additional pulsars (for which new timings were not available) were taken directly from the ATNF catalogue, selected using criteria described in the following section. The parameters of the X-ray pulsar PSR J0537–6910 were taken from Ref. [26] and those for the Crab pulsar from the Jodrell Bank monthly ephemeris [24]. The remaining 61 pulsars (from the original list of 154) were not timed with sufficient confidence and were excluded from the search. This included many of the newly discovered pulsars (for example the 21 millisecond pulsars in the Terzan 5 globular cluster [27]) for which accurate timing solutions have yet to be published. We therefore had a catalogue of 93 timed pulsars for our gravitational wave search.

B. Error propagation in source parameters

The impact of parameter uncertainties on the search was assessed for both the S3 and S4 runs. At some level there are positional, frequency and frequency derivative uncertainties for all the target pulsars, and for pulsars in binary system there are also uncertainties associated

with all the binary orbital parameters. Some of these uncertainties are correlated, for example the error on frequency could affect the accuracy of the first frequency derivative, and the binary time of periastron and longitude of periastron are also highly correlated.

We took a ‘worst-case scenario’ approach by adding and subtracting the quoted uncertainties from the best-fit values of all the parameters to determine the combination which gave a maximum phase deviation, when propagated over the period of the run (either S3 or S4), from the best fit phase value calculated over the same time period. For example if we assume $\phi(t_{S3})$ given by Eq. (1.3) (ignoring for simplicity the ϕ_0 and $\dot{\nu}$ terms) is the best fit phase over the time span of S3, t_{S3} , the maximum phase uncertainty is

$$\Delta\phi_{\text{err}} = \max \left[\left| \phi(t_{S3}) \pm 2\pi \left\{ (\nu \pm \sigma_\nu)(t_{S3} \pm \sigma_{t_{S3}}) + \frac{1}{2}(\dot{\nu} \pm \sigma_{\dot{\nu}})(t_{S3} \pm \sigma_{t_{S3}})^2 + \dots \right\} \right| \right], \quad (4.1)$$

where the σ s are the uncertainties on the individual parameters. Correlations between the parameters mean that this represents an upper limit to the maximum phase uncertainty, sometimes greatly overestimating its true value.

There are 12 pulsars with overall phase uncertainty $> 30^\circ$ in S3, which we take as the threshold of acceptability. A 30° phase drift could possibly give a factor of $\sim 1 - \cos 30^\circ = 0.13$ in loss of sensitivity for a signal. Nine of these are in binary systems (PSRs J0024–7204H, J0407+1607, J0437–4715, J1420–5625, J1518+0205B, J1709+2313, J1732–5049, J1740–5340 and J1918–0642) and in five of these T_0 and ω_0 contribute most to the phase uncertainty. For the three isolated pulsars (PSRs J0030+0451, J0537–6910, and J1721–2457) the phase error is dominated by uncertainties in frequency and/or position.

Applying the same criterion to the time-span of S4 we find that PSR J1730–2304 rises above the limit. For this pulsar its parameter uncertainties do not affect it for the S3 analysis as it was timed over this period, however when extrapolating over the time of the S4 run the uncertainties become non-negligible.

In total there are 13 pulsars rejected over the combined run. This highly conservative parameter check reduces our 93 candidate pulsars to 80.

C. Timing noise

Pulsars are generally very stable rotators, but there are phenomena which can cause deviations in this stability, generically known as timing noise. The existence of timing noise has been clear since the early days of pulsar astronomy and appears as a random walk in phase, frequency or frequency derivative of the pulsar about the regular spin-down model given in Eq. (1.3) [28]. The strength of this effect was quantified in Ref. [28] as an

activity parameter A , referenced to that of the Crab pulsar, and in Ref. [29] as a *stability parameter* Δ_8 . A is based on the logarithm of the ratio of the rms residual phase of the pulsar, after removal of the timing model, to that of the Crab pulsar over an approximately three-year period. Δ_8 is not based on the stochastic nature of the Crab pulsar’s timing noise and is defined for a fixed time (10^8 s) as

$$\Delta_8 = \log \left(\frac{1}{6\nu} |\ddot{\nu}| \times (10^8 \text{ s})^3 \right). \quad (4.2)$$

This assumes that the measured value of $\ddot{\nu}$ is dominated by the timing noise rather than the pulsar’s intrinsic second spin-down derivative. Although generally true, this assumption is not valid for the Crab pulsar and PSR J0537–6910, where a non-timing noise dominated $\ddot{\nu}$ can be measured between glitches³. This quantity relates to the pulsar clock error caused by timing noise. The value of $\ddot{\nu}$ is so small as to be unmeasurable for most pulsars, although an upper limit can often be defined. Arzoumanian *et al.* [29] deduce, by eye, a linear relationship between Δ_8 and $\log \dot{P}$ of

$$\Delta_8 = 6.6 + 0.6 \log \dot{P}, \quad (4.3)$$

where $\dot{P} = -\dot{\nu}/\nu^2$ is the period derivative.

As defined, Δ_8 is a somewhat imprecise indicator of the timing noise, not least because the time span of 10^8 seconds chosen by Arzoumanian *et al.* was simply the length of their data set. A preferred measure may simply be the magnitude and sign of \dot{P} , but we shall continue to use the Δ_8 parameter as our timing noise magnitude estimate for the current analysis. A thorough study of timing noise, comparing and contrasting the various measures used, will be given in Ref. [30] (also see Refs. [25, 31]).

There is a definite correlation between the Δ_8 parameters, spin-down rate and age. Young pulsars, like the Crab pulsar, generally show the most timing noise. The categorisation of the type of timing noise (i.e., phase, frequency or frequency derivative) in Ref. [28] allowed them to ascribe different processes for each. The majority of pulsars studied showed frequency-type noise, possibly a result of random fluctuations in the star’s moment of inertia. The actual mechanism behind the process is still unknown, with Cordes and Greenstein [32] positing and then ruling out several mechanisms inconsistent with observations.

Timing noise intrinsically linked to motions of the electromagnetic emission source or fluctuations in the magnetosphere, rather than the rotation of the pulsar, is important in the search for gravitational waves as it may allow the relative phase of the electromagnetic and gravitational signals to drift. The implications of timing noise

³ These two pulsars are among the most prolific glitchers, and in any global fit to their parameters the value of $\ddot{\nu}$ would most likely be swamped by the glitch events.

in this context are discussed by Jones [33]. He gives three categories of timing noise, not necessarily related to the three types of timing noise given by Cordes and Helfand [28], having different effects on any search. If all parts of the neutron star are strongly coupled on short timescales, there should be no difference between the electromagnetic phase and gravitational wave phase. If the timing noise were purely a magnetospheric fluctuation then phase wandering caused by timing noise would not be seen in the gravitational wave emission. The third possibility, whereby the electromagnetic emission source wanders with respect to the mass quadrupole, could result from a weak exchange of angular momentum between the parts of the star responsible for electromagnetic and gravitational wave emission. Jones describes the ratio of the electromagnetic and gravitational timing noise phase residuals ($\Delta\Phi$) by a parameter $\alpha = \Delta\Phi_{\text{gw}}/\Delta\Phi_{\text{em}}$, with the three types of timing noise described above corresponding to $\alpha = 1, 0$ and $-I_{\text{em}}/I_{\text{gw}}$ respectively, where the I s represent the moments of inertia of the electromagnetic and gravitational wave producing components. In principle this factor could be included as another search parameter. However, given the cost of including an extra parameter in this search, and given that it is plausible that that all parts of a neutron star are tightly coupled on the timescales of interest here, we will assume rigid coupling between the two components, i.e set $\alpha = 1$, corresponding to the gravitational and electromagnetic signals remaining perfectly in phase.

The Crab pulsar is regularly monitored [24] on timescales that are sufficiently short to allow its timing noise to be effectively removed using a second heterodyne procedure [34]. Like the Crab pulsar, PSR J0537–6910 is young, has a high glitch rate and also shows high levels of timing noise [26]. Unfortunately, unlike the Crab pulsar, we have no regular ephemeris for it that covers our data set, and timing irregularities are likely to be too great for historical data to be of use. We therefore have excluded PSR J0537–6910 from the analysis. For less noisy pulsars we still need a method of estimating the effect of timing noise on phase evolution that does not rely on continuous observation. One such estimate is the Δ_8 parameter given by Eq. (4.2), which can provide a measure of the cumulative phase error. For those pulsars with a measured $\ddot{\nu}$ we use this estimate to obtain a corresponding value of Δ_8 as shown in Fig. 3. This should provide a reasonable estimate of the timing noise over the timespan of the pulsar observation. Again we apply our criterion that cumulative phase errors of $> 30^\circ$ are unacceptable. In Fig. 3 there are four pulsars (those with the four largest Δ_8 values), with measured $\ddot{\nu}$, for which this is the case, and therefore timing noise could be a problem (having already noted the Crab pulsar and PSR J0537–6910 as exceptions): PSRs J1748–2446A, J1823–3021A, J1913+1011, and J1952+3252. For pulsars with no measured $\ddot{\nu}$ we use the approximate linear relation between the period derivative \dot{P} and Δ_8 given in Eq. (4.3). The low \dot{P} values for these pulsars implies that timing noise will be




FIG. 3: The values of Δ_8 for our selection of pulsars with measured $\ddot{\nu}$.

negligible.

In addition to the above, there are some pulsars in globular clusters for which there is no $\ddot{\nu}$ and for which \dot{P} is negative ($\ddot{\nu}$ is positive), so no value of Δ_8 can be assigned either through Eqs. (4.2) or (4.3). For these pulsars the value of $\ddot{\nu}$ (and therefore $\ddot{\nu}$) must be rather small to have been affected by motions within the cluster (discussed more in §VI), so timing noise should again be negligible.

For pulsars which were re-timed over the period of S3 timing noise will be negligible (for the S3 analysis at least) as any timing noise, which usually has variations on time-scales of several months to years, will have been absorbed in the parameter estimation. PSRs J1748–2446A and J1823–3021A were re-timed over S3 meaning that their S3 results will stand, although the other two will not. However, being conservative we will remove all four pulsars with large values of Δ_8 , and PSR J0537–6910, in which timing noise could be problematic, from the S4 and joint analysis. Note that PSR J0537–6910 is vetoed by both the parameter error criterion and our timing noise criterion.

This reduces our final number of well parameterised pulsar targets to 78 for the S3 analysis and 76 for the S4 and joint analyses. The 76 pulsars include 21 of the 28 from the previous study of Abbott *et al.* [3], and so through our selection criterion we lose the following 7 previously analysed pulsars: PSRs J0030+0451, J1721–2457, J1730–2304, J1823–3021A, J1910–5959B, J1913+1011, and J1952+3252. The same selection rules were not applied over S2, especially of note was that no timing noise criterion was considered, which accounts for three of the pulsars we lose between the two analyses. Also our 30° rule was strictly applied, which the other four pulsars just exceeded.

The analysis was actually performed on all 93 timed

pulsars mentioned above, however the various parameter uncertainties preclude us setting upper limits on a total of 15 of these.

V. HARDWARE INJECTIONS

For analysis validation purposes simulated gravitational wave signals for a variety of sources (bursts, pulsars, inspirals and stochastic) have been mechanically injected into the LIGO interferometers during science runs. During S2 two pulsar signals were injected [3]. This was increased to 10 injections in the LIGO instruments for S3 and 12 for S4 covering a wider range of signal parameters. Extracting and understanding these injections has been invaluable in validating the analysis.

The hardware injection signals are produced using software (under LALApps [35]), which was largely developed independently of the extraction code. However, the codes do share the same solar system barycentring and detector antenna response function routines, both of which have been extensively checked against other sources (e.g. checks against TEMPO in Refs. [1, 21]).

The signals were added into each of the three LIGO detectors via the position control signal going to the end test mass in one arm. Control signals in the digital servos that maintain optical cavities on resonance were summed with fake pulsar waveforms, modulating mirror positions to mimic the effect of a real spinning compact object (i.e. differential length motions with frequency and amplitude modulations appropriate for a given sky position, frequency and spin-down). Furthermore, as the digital fake waveforms have to be converted to analog coil currents of suspended optics, the injected waveforms have to be divided by the transfer function of the output chain (predominantly the pendulum), in order to produce the desired differential length response of the cavity.

The extraction of these injections is described in detail in Appendix A. They show the relative phase consistency between the detectors over the course of a run. This means that a joint analysis combining the data from all detectors is valid. The injection plots (see Figs. 7 and 8) show what we would expect our posterior plots to look like given a detection i.e. strongly peaked pdfs with very small probability at $h_0 = 0$, as compared to those in Fig. 2 where h_0 peaks at zero.

VI. RESULTS

A. Upper limits

Here we present 95% degree-of-belief upper limits on the amplitude of gravitational waves (h_0) from the 78 pulsars identified above. The value of h_0 is independent of any assumptions about the neutron star other than it is emitting gravitational waves at twice its rotation frequency. The results will also be presented in terms

of the pulsars' equatorial ellipticity ε , which under the assumption of triaxiality is related to h_0 via Eq. (1.1) by

$$\varepsilon = 0.237 \left(\frac{h_0}{10^{-24}} \right) \left(\frac{r}{1 \text{ kpc}} \right) \left(\frac{1 \text{ Hz}}{\nu} \right)^2 \left(\frac{10^{38} \text{ kg m}^2}{I_{zz}} \right). \quad (6.1)$$

To obtain an upper limit on ε from that for h_0 we assume a fiducial moment of inertia value of $I_{zz} = 10^{38} \text{ kg m}^2$. We discuss below in Sec. VI B the effect of relaxing this assumption. Pulsar distances are taken from the ATNF catalogue [10] and are generally derived from the radio dispersion measures, with errors estimated to be of order 20%, although in some cases even this can be an underestimate. A critical review of pulsar distance measurements can be found in Ref. [36].

All upper limit results from the individual S3 and S4 runs along with results from the combined run, with and without GEO 600 included, are given in Appendix B in Tables III and IV. The GEO 600 data only provides comparable sensitivities to LIGO at frequencies greater than 1000 Hz, and are therefore only used in the search for PSR J1939+2134 (at the time the fastest known millisecond pulsar) in S3, and additionally PSR J1843–1113 in S4 and the combined run. Inclusion of GEO 600 does not significantly change the joint upper limits for these pulsars. For the majority of pulsars the lowest upper limits come from the combined S3/S4 data set, although for 7 pulsars (PSRs J0024–7204I, J0024–7204S, J0024–7204U, J1045–4509, J1802–2124, J1910–5959D and J1911+0101B) the S4 results alone provide a lower limit. The combined S3 and S4 run results are presented in histogram form in Fig. 4. Fig. 5 shows the results compared to a joint LIGO S4 upper limit estimate curve, taken as the best sensitivity during S4. The joint upper limit sensitivity curve for the three detectors can be estimated by combining the detector one-sided power spectral densities (PSDs) via

$$S(f) = \left(\frac{T_{\text{obs H1}}}{S_h(f)_{\text{H1}}} + \frac{T_{\text{obs H2}}}{S_h(f)_{\text{H2}}} + \frac{T_{\text{obs L1}}}{S_h(f)_{\text{L1}}} \right)^{-1}, \quad (6.2)$$

$$h_0^{95\%} = 10.8 \sqrt{S(f)},$$

where $S_h(f)$ is the PSD and T_{obs} is each detector's live time (using the associated duty factor of each interferometer during the run). The factor of 10.8 is given in Ref. [1] and was calculated through simulations with Gaussian noise⁴.

The results are also compared to the upper limit deduced from the observed spin-down via Eq. (1.2) making the assumption that all rotational energy is lost through

⁴ In Ref. [3] a similar plot to Fig. 5 is shown for the S2 data using a factor of 11.4 in the relation between the upper limit and PSD. This definition comes from using the \mathcal{F} -statistic search method and setting a 1% false alarm rate and 10% false dismissal rate for signals given the underlying detector PSD [2]




FIG. 4: Histograms of the log of amplitude, ellipticity, and ratio of spin-down to gravitational wave upper limits for the combined LIGO S3 and S4 run.




FIG. 5: The combined S3 and S4 upper limit results on the amplitude of gravitational waves for 76 pulsars using LIGO data compared to the joint sensitivity curve.

gravitational wave emission. The spin-down limit is seen as a natural crossing point after which gravitational-wave data, including upper limits, have a likely bearing on the nature of the neutron star. The spin-down upper limit will obviously depend on $\dot{\nu}$. This value however, can be masked by radial and transverse motions of the object (see Ref. [37] for discussion of these effects). The Shklovskii effect [38], in which the pulsar has a large transverse velocity v , will cause an apparent rate of change in the pulsar’s period of

$$\dot{P}_S = \frac{v^2}{rc} P. \quad (6.3)$$

Its $1/r$ dependence makes this effect more prominent for nearby pulsars. In the ATNF catalogue [10] values of the

intrinsic period derivative $\dot{P}_{\text{int}} = \dot{P} - \dot{P}_S$ can be obtained where this effect has been corrected for. This provides a measure of intrinsic (rather than apparent) spin-down⁵ and, when available, is used in the spin-down ratio results.

The observed value of \dot{P}_{obs} will also differ from its intrinsic value, \dot{P}_{int} , if the pulsar is accelerating – a likely scenario in the gravitational field of a globular cluster [37]. Any line-of-sight component to the acceleration, $a_{||}$, will give an observed value of

$$\dot{P}_{\text{obs}} = \dot{P}_{\text{int}} + \frac{a_{||}}{c} P \quad (6.4)$$

where P is the spin period [39]. These effects can cause pulsars to have apparent spin-ups (seen in quite a large number of globular cluster pulsars) although are only strong enough to greatly affect pulsars with intrinsically small period derivatives. There are still many globular clusters for which the radial accelerations have not been measured, therefore no firm spin-down upper limit can be set, making the direct gravitational wave results a unique limit.

Highlights of the combined S3/S4 results include the tightest strain upper limit set on a pulsar of $h_0^{95\%} = 3.2 \times 10^{-25}$ for PSR J1603–7207, the smallest ellipticity at $\epsilon = 8.5 \times 10^{-7}$ for PSR J2124–3358, and the closest upper limit to the spin-down limit at a ratio of 2.7 for the Crab pulsar (PSR J0534+2200).

⁵ Note that the heterodyne procedure still needs to make use of the measured spin-down rather than the intrinsic spin-down as these Doppler effects will have the same effect on the gravitational waves.

B. Dependence on the moment of inertia

The pulsar ellipticity results detailed above assume a moment of inertia of 10^{38} kg m^2 , which is the standard fiducial number used in the literature. However, modern theoretically computed equations of state (EOS) generally predict somewhat larger moments of inertia for stars more massive than $1 M_\odot$, a group which includes all neutron stars with measured masses (see Ref. [40]). Therefore the dependence on moment of inertia should be considered.

Bejger, Bulik, and Haensel [41] give an overview of the theoretical expectations for the moment of inertia. Their Fig. 2 plots the moment of inertia vs. mass for several theoretically predicted types of EOS. The maximum moment of inertia they find (after varying the mass of the star) is 2.3 times the fiducial value, with stars of $1.4 M_\odot$ having moments of inertia 1.2–2.0 except for one outlying type of EOS. Typically the maximum moment of inertia occurs for a neutron star mass of $1.7 M_\odot$ or more. Recently masses greater than $1.6\text{--}1.7 M_\odot$ with 95% confidence have been measured [27, 42] for some systems, making this reasonable to consider. More recently Lackey [43] found the highest moment of inertia to be $3.3 \times 10^{38} \text{ kg m}^2$ for EOS G4 of Lackey, Nayyar, and Owen [44]. This is a relativistic mean-field EOS similar to the GNH family considered by Bejger, Bulik, and Haensel [41] but contains no exotic phases of matter such as hyperons or quarks. Consequently, we consider the range of theoretically predicted moments of inertia to be approximately $1\text{--}3 \times 10^{38} \text{ kg m}^2$.

There have been recent attempts to infer neutron star moments of inertia from observations. Bejger and Haensel [45, 46] derived a value for the Crab pulsar’s moment of inertia by equating the spin-down power to the observed electromagnetic luminosity and inferred acceleration of the nebula. However, this (extremely high) value is dominated by the assumptions about the highly uncertain mass and mass distribution of the nebula as well as the relativistic wind from the pulsar, and thus cannot yet be considered to give a reliable value. The double pulsar system J0737-3039 shows great promise for tighter measurements of the moment of inertia (and constraints on the EOS) in the near future [41, 47, 48, 49]. However, for the moment we are left with the theoretical range quoted above.

As suggested in Ref. [50], instead of using Eq. (6.1) to set a limit on ε assuming a value of I_{zz} , one can use it to set a limit on the neutron star quadrupole moment $\approx I_{zz}\varepsilon$ without relying on any assumption about I_{zz} . The limit on the quadrupole moment can then be used to help define an exclusion region in the $I\text{--}\varepsilon$ plane. This exclusion region allows one to read off an upper limit on ε as a function of the EOS-dependent moment of inertia. The spin-down can also be used to provide exclusion regions via the relation

$$I_{zz} = \frac{5|\dot{\nu}|c^5}{512G\nu^5} \frac{1}{\varepsilon^2}. \quad (6.5)$$

Theoretical contributions to the exclusion regions come from predictions of the maximum moment of inertia and ellipticity. In terms of the exclusion region, our observational upper limits on h_0 are far from contributing except for the Crab pulsar, to which we now turn.

C. The Crab pulsar - PSR J0534+2200

Of the known radio pulsars, the Crab pulsar has often been considered one of the most promising sources of gravitational waves. This is due to its youth and large spin-down rate, leading to a relatively large spin-down upper limit several orders of magnitude higher than for most other pulsars. The high rate of glitching in the pulsar also provides possible evidence of asymmetry. One glitch model favoured for the Crab pulsar involves a change in the pulsar ellipticity, and breaking of the crust, as the star settles to its new equilibrium state as it spins down [37]. In the 1970s, estimates of gravitational wave strains were spurred on by the experimenters producing novel technologies which allowed the possibility of probing these low strains, with Zimmermann [51] producing estimates of gravitational wave strains from the Crab pulsar ranging from $h_0 \approx 2 \times 10^{-25} - 10^{-29}$.

The first searches for gravitational waves from the Crab pulsar were carried out using specially designed resonant bar detectors, with frequencies of around 60 Hz [6]. The most recent result using such a bar was from 1993 and gave a 1σ upper limit of $h_0 \leq 2 \times 10^{-22}$ [7]. This upper limit was passed in the LIGO S2 run, which gave $h_0^{95\%} = 4.1 \times 10^{-23}$ [3]. Using Eq. (1.2), and taking $I_{zz} = 10^{38} \text{ kg m}^2$ and $r = 2 \text{ kpc}$, gives a spin-down upper limit for the Crab pulsar of $h_0 < 1.4 \times 10^{-24}$, about a factor of 30 below the S2 observational upper limit. However, the S2 limit on the Crab was at the time the closest approach to the spin-down limit obtained for any pulsar.

Our new results for the Crab pulsar (and the other 77 targets) are shown in Table III. The results improve by up to an order of magnitude over those from the S2 run, and the majority of this improvement was between the S2 and S3 runs. The results for the Crab pulsar over the S2, S3 and S4 runs are plotted on the $I\text{--}\varepsilon$ plane in Fig. 6. The solid lines in Fig. 6 mark the lower boundaries of exclusion regions on this plane using our upper limits obtained for the different runs. The dashed black line marks the lower boundary of the upper limit from spin-down as given in Eq. (6.5). The dashed horizontal grey lines give lower and upper bounds on the moment of inertia of $1\text{--}3 \times 10^{38} \text{ kg m}^2$, as given by our arguments in §VIB. It can be seen that our experimental results currently only beat the spin-down limit for moments of inertia at values greater than double the maximum of our theoretical range. However, over this range the ratio of the gravitational to spin-down upper limit ranges from 2.7 at the lowest value to only 1.5 at the largest value.

The spin-down limit in fact over-estimates the




FIG. 6: The moment of inertia–ellipticity plane for the Crab pulsar over the S2, S3 and S4 runs. The areas to the right of the solid lines are the experimentally excluded regions. The horizontal lines represent theoretical upper and lower limits on the moment of inertia as mentioned in §VIB. Theoretical upper limits on the ellipticity are much more uncertain, the highest being a few times 10^{-4} .

strongest possible signal because we know that much of the spin-down energy of the Crab goes into powering the nebula through electromagnetic radiation and relativistic particle winds. Thus it is interesting to ask how far we would need to beat the spin-down limit to have a chance of detecting a signal allowing for what is known about the non gravitational wave spin-down. Palomba [52] uses the observed braking index 2.51 of the Crab pulsar with a simple model of spin-down through gravitational radiation (braking index 5) combined with some other mechanism (braking index a free parameter) to place an upper limit of about $\epsilon \leq 3 \times 10^{-4}$. This is about 2.5 times lower than the spin-down limit and nearly 7 times lower than our result (for $I_{zz} = 10^{38} \text{ kg m}^2$).

The Crab pulsar experienced two glitches between S3 and S4, a large glitch on 6th September 2004 and a smaller glitch on 22nd November 2004 [24]. The effect of glitches on the relative phase between the electromagnetic pulse and any possible gravitational wave signal is unknown, so there is uncertainty whether the (phase-coherent) combined S3/S4 result is valid. The combined result stands, but the reader should be aware that it includes the assumption of trans-glitch phase coherence.

VII. ASTROPHYSICAL INTERPRETATION

We have produced new, tight, upper limits on gravitational wave signal strength from a large selection of known pulsars and for the Crab pulsar we are very near the fiducial limit set by spin-down arguments.

It can be seen from Table III and Fig. 4 that, for the majority of pulsars, the gravitational wave detector up-

per limits are at least one hundred times above those from the spin-down argument, so is there anything that we can take from the results in terms of astrophysics?

First we should note that spin-down limits on gravitational wave luminosity are plausible, but model dependent. They assume a model for the structure of the neutron star (for instance, that it is not accreting and is rigidly rotating, in addition to assumptions about its equation of state) and they take dispersion measure distance as a consistently good measure of true distance. There is some considerable uncertainty associated with all of these assumptions. In contrast our observations set direct limits on a source’s gravitational wave strain.

Secondly, for globular cluster pulsars the spin-down measured from radio timing observations is a combination of the spin-down intrinsic to the pulsar and acceleration along the line-of-sight $a_{||}$ in the cluster’s gravitational potential (see Eq. (6.4)). In general, the magnitude and sign of the acceleration is unknown but the intrinsic $\dot{P}_{\text{int}} > 0$ of millisecond pulsars is usually small and often smaller than the extrinsic contribution. Only if $\dot{P}_{\text{obs}} < 0$ one can be sure that $a_{||} < 0$. Therefore, the limits derived from our gravitational wave observations provide the only direct limits on \dot{P}_{int} which are independent from biasing kinematic effects. These can be combined with the observed spin-down to provide a limit on the acceleration in the cluster, i.e. $a_{||} \geq c(\dot{P}_{\text{obs}} - \dot{P}_{\text{gw}}^{\text{limit}})/P$.

Finally, it is interesting to note that our ellipticity limits are well into the range permitted by some models of strange quark stars or hybrid stars ($\epsilon \sim$ a few times $10^{-4} - 10^{-5}$) and are reaching into the range permitted by more conventional neutron star EOSs ($\epsilon \sim$ a few times 10^{-7}) [53].

Currently the fifth LSC science run (S5) is underway, and this promises to beat the Crab pulsar spin-down limit within a few months of its start. For many other pulsars we should be able to reach amplitude upper limits of $< 1 \times 10^{-25}$ and ellipticities of $\sim 1 \times 10^{-7}$.

Acknowledgments

The authors gratefully acknowledge the support of the United States National Science Foundation for the construction and operation of the LIGO Laboratory and the Particle Physics and Astronomy Research Council of the United Kingdom, the Max-Planck-Society and the State of Niedersachsen/Germany for support of the construction and operation of the GEO600 detector. The authors also gratefully acknowledge the support of the research by these agencies and by the Australian Research Council, the Natural Sciences and Engineering Research Council of Canada, the Council of Scientific and Industrial Research of India, the Department of Science and Technology of India, the Spanish Ministerio de Ciencia y Tecnología, the John Simon Guggenheim Foundation, the Leverhulme Trust, the David and Lucile Packard Foundation, the Research Corporation, and the Alfred P. Sloan

-
- [1] R. J. Dupuis and G. Woan, Phys. Rev. D **72**, 102002 (2005).
- [2] B. Abbott *et al.* (LIGO Scientific Collaboration), Phys. Rev. D **69** 082004 (2004).
- [3] B. Abbott *et al.* (LIGO Scientific Collaboration), Phys. Rev. Lett. **94** 181103 (2005).
- [4] B. Abbott *et al.* (LIGO Scientific Collaboration), arXiv:gr-qc/0605028 (2006).
- [5] P. Jaranowski, A. Królak, and B. F. Schutz, Phys. Rev. D **58**, 063001 (1998).
- [6] H. Hirakawa, K. Tsubono and M. K. Fujimoto, Phys. Rev. D **17**, 1919 (1978).
- [7] T. Suzuki, First Edoardo Amaldi Conference on Gravitational Wave Experiments, 115 (1995).
- [8] J. Hough *et al.*, Nature **303**, 216 (1983).
- [9] M. Hereld, PhD thesis, California Institute of Technology (1983).
- [10] Australia Telescope National Facility Pulsar Catalogue <http://www.atnf.csiro.au/research/pulsar/psrcat/>.
- [11] R. J. Dupuis, PhD Thesis, University of Glasgow (2004).
- [12] F. Acernese *et al.*, Class. Quant. Grav. **22**, 869 (2005).
- [13] T. Creighton, Class. Quant. Grav. **20**, 853 (2003).
- [14] <http://einstein.phys.uwm.edu/PartialS3Results>
- [15] B. Abbott *et al.* (LIGO Scientific Collaboration), Phys. Rev. D **72** 102004 (2005).
- [16] B. Abbott *et al.* (LIGO Scientific Collaboration) “*All-sky LIGO search for periodic gravitational waves in the S4 data*”, in preparation.
- [17] J. H. Taylor and J. M. Weisberg, Astrophys. J. **345**, 434 (1989).
- [18] J. R. Smith *et al.*, Class. Quant. Grav. **21**, S1737 (2004).
- [19] S. Hild *et al.*, J. Phys.: Conf. Ser. **32**, 66 (2006).
- [20] <http://pulsar.princeton.edu/tempo>.
- [21] M. D. Pitkin and G. Woan, in preparation (2006).
- [22] C. Lange *et al.*, Mon. Not. R. Astron. Soc. **326**, 274 (2001).
- [23] R. N. Manchester *et al.*, Astron. J. **129**, 1993 (2005).
- [24] Jodrell Bank Crab Pulsar Monthly Ephemeris <http://www.jb.man.ac.uk/research/pulsar/crab.html>.
- [25] G. Hobbs *et al.*, Mon. Not. R. Astron. Soc. **353**, 1311 (2004).
- [26] F. E. Marshall *et al.*, Astrophys. J. **603**, 682 (2004).
- [27] S. M. Ransom *et al.*, Science **307**, 892 (2005).
- [28] J. M. Cordes and D. J. Helfand, Astrophys. J. **239**, 640 (1980).
- [29] Z. Arzoumanian *et al.*, Astrophys. J. **422**, 671 (1994).
- [30] G. Hobbs, A. G. Lyne and M. Kramer, in preparation.
- [31] G. Hobbs *et al.*, Mon. Not. R. Astron. Soc. **360**, 974 (2005).
- [32] J. M. Cordes and G. Greenstein, Astrophys. J. **245**, 1060-1079 (1981).
- [33] D. I. Jones, Phys. Rev. D **70**, 042002 (2004).
- [34] M. Pitkin and G. Woan, Class. Quant. Grav., **21**, 843 (2004).
- [35] <http://www.lsc-group.phys.uwm.edu/daswg/projects/lalapps.html>.
- [36] D. A. Frail and J. M. Weisberg, Astron. J. **100**, 743-757 (1990).
- [37] A. Lyne and F. Graham-Smith, Pulsar Astronomy, Cambridge University Press (1998).
- [38] I. S. Shklovskii, Soviet Astronomy **13**, 562 (1970).
- [39] E. S. Phinney, ASP Conf. Ser. 50: Structure and Dynamics of Globular Clusters (1993).
- [40] S. E. Thorsett and D. Chakrabarty, Astrophys. J. **512**, 288 (1999).
- [41] M. Bejger, T. Bulik and P. Haensel, Mon. Not. R. Astron. Soc. **364**, 635 (2005).
- [42] D. J. Nice *et al.*, Astrophys. J. **634**, 1242-1249 (2005).
- [43] B. D. Lackey, Undergraduate honours thesis, The Pennsylvania State University (2006).
- [44] B. D. Lackey, M. Nayyar and B. J. Owen, Phys. Rev. D **73**, 024021 (2006).
- [45] M. Bejger and P. Haensel, Astron. Astrophys. **396**, 917 (2002).
- [46] M. Bejger and P. Haensel, Astron. Astrophys. **405**, 747 (2003).
- [47] I. A. Morrison *et al.*, Astrophys. J. **617**, L135 (2004).
- [48] J. M. Lattimer and B. F. Schutz, Astrophys. J. **629**, 979 (2005).
- [49] M. Kramer *et al.*, Science **314**, 5796, 97 (2006).
- [50] M. Pitkin *et al.* (LIGO Scientific Collaboration), Class. Quant. Grav. **22**, S1277 (2005).
- [51] M. Zimmerman, Nature **271**, 527 (1978).
- [52] C. Palomba, Astron. Astrophys. **354**, 163 (2000).
- [53] B. Owen, Phys. Rev. Lett. **95**, 211101 (2005).
- [54] U. Weiland *et al.*, Class. Quant. Grav. **21**, 861 (2004).

APPENDIX A: INJECTIONS

1. S3 injections

An initial analysis of the S3 pulsar injections is given in Ref. [11]. The data have since been re-analysed with more recent versions of the detector calibrations, the results of which are presented here. For S3 initially 10 pulsars signals were injected, with a further one added at the end of the run to be in coincidence with a single injection into GEO 600 [54]. The majority of injection parameters were decided upon randomly, although pulsar frequencies were chosen to avoid major instrumental or calibration lines, and amplitudes were dependent on the frequency. The injections were split into two groups of five, where values of h_0 were calculated to give two each with signal-to-noise ratios of approximately 3, 9, 27, 81 and 243. The parameter values are shown in Table I. The 10 initial signals were injected into the LIGO detectors for approximately the first half of the run, then turned off for two weeks, to ensure data were present that was not artificially contaminated, and then turned back on with the two loudest signals removed. The simultaneous injection with GEO 600 was switched on near the end of the run.

These signals were extracted from the data using the analysis techniques described in §III and Ref. [1]. The two most important parameters for checking that the calibration of the instruments was correct were the amplitude and initial phase, so in the Bayesian parameter estimation procedure the ι and ψ parameters were held fixed at their known values. This was done because the correlations between h_0 and $\cos \iota$ and ϕ_0 and ψ , respectively could lead to the marginalised posterior probability distributions (pdfs) for each parameter being distorted or spread out (see Ref. [11] for examples of this). The extracted pdfs of h_0 and ϕ_0 for each of the injections, after corrections described below in §A 3, can be seen in Fig. 7.

For the vast majority of signals the extracted pdfs overlap with the injected value. For the strongest injections with the largest signal-to-noise ratios the pdfs are rather narrow, and any uncertainties in the calibration become evident, with a maximum offset of the order of 10–15%. The far wider pdfs associated with the L1 signal injections reflect the lower L1 sensitivity and lower duty factor compared with the H1 and H2 detectors. It can be seen that the injected phases for each detector agree with each other to within a few degrees and are within the uncertainty of the method. This provides some evidence that there is phase coherence between the detectors and that a joint analysis, combining the data from all the detectors, is possible.

Two main discrepancies have been identified as operational mistakes made during the injection procedure: PULSAR7 was injected into H2 with a much lower amplitude than intended, and remained undetected, therefore no joint analysis was performed, and PULSAR0 was injected into H1 with an amplitude 1.6 times larger than

intended.

The injection of the signal into GEO 600 is described in Ref. [54], and its analysis is described in Ref. [11]. It was found that the injection performed during S3 was badly contaminated and could not be used. However, a subsequent injection performed shortly after S3 has verified that the signal parameters were correctly injected and extracted, validating the injection hardware and analysis software.

2. S4 injections

The 10 injections used in S3 were used again for S4 to create artificial signals in the LIGO interferometers. However, their amplitudes were adjusted to give approximately the same signal-to-noise ratios as seen in S3, taking into account the better sensitivity during the S4 run. For all except PULSAR9 the h_0 values were reduced by a \sim half, with PULSAR9 being so strong that its amplitude was reduced by a factor of \sim 20. These signals were injected for the second half of the run from 8th March 2005 onwards. The updated h_0 values are shown in Table I. There were also an additional two signals (PULSAR10 and 11), simulated to be from pulsars in binary systems, injected for the last day of the run. The binary pulsar injections allowed the testing of the binary timing code described in §III B as the injection code and extraction code were written independently. The binary injection signal parameters for PULSAR10 and PULSAR11 were taken from PULSAR3 and 8 respectively, with the frequencies changed, and amplitudes increased to make sure they were visible over the short injection time. The frequency, amplitude and binary system parameters are shown in Table II. The binary system parameters were chosen to have one in a relatively eccentric orbit and one in a circular orbit. We chose fairly short periods, so that they would have completed or nearly completed at least one full orbit during the injection. The T_0 values are given in the pulsar rest frame. For the recovery of the binary system injections the BT model was used, although as no relativistic parameters were included any of the models could have been used.

The extracted amplitude and phase pdfs, after corrections described below in §A 3, are shown in Fig. 8 The observed phase consistency between the detectors, means that joint likelihoods, using all three detectors, can be calculated. In general the values of h_0 are well matched with the injection values. It can again be seen that for the strongest signals the narrow pdfs are offset from the injected value in h_0 reflecting the calibration uncertainties of 5-10%.

The binary pulsar injections show matches to their injected values. This is a good confirmation that the binary timing code can track the phase well and has no significant errors.

TABLE I: The parameter values for the pulsar hardware injections in S3 and S4.

PULSAR	α (rads)	δ (rads)	ν_{gw} (Hz)	$\dot{\nu}_{\text{gw}}$ (Hz/s)	h_0 (S3)	h_0 (S4)	ϕ_0 (rads)	ι (rads)	ψ (rads)
0	1.25	-0.98	265.5	-4.15×10^{-12}	9.38×10^{-25}	4.93×10^{-25}	2.66	0.65	0.77
1	0.65	-0.51	849.1	-3.00×10^{-10}	8.49×10^{-24}	4.24×10^{-24}	1.28	1.09	0.36
2	3.76	0.06	575.2	-1.37×10^{-13}	1.56×10^{-23}	8.04×10^{-24}	4.03	2.76	-0.22
3	3.11	-0.58	108.9	-1.46×10^{-17}	6.16×10^{-23}	3.26×10^{-23}	5.53	1.65	0.44
4	4.89	-0.21	1430.2	-2.54×10^{-8}	1.01×10^{-21}	4.56×10^{-22}	4.83	1.29	-0.65
5	5.28	-1.46	52.8	-4.03×10^{-18}	1.83×10^{-23}	9.70×10^{-24}	2.23	1.09	-0.36
6	6.26	-1.14	148.7	-6.73×10^{-9}	5.24×10^{-24}	2.77×10^{-24}	0.97	1.73	0.47
7	3.90	-0.36	1221.0	-1.12×10^{-9}	2.81×10^{-23}	1.32×10^{-23}	5.24	0.71	0.51
8	6.13	-0.58	194.3	-8.65×10^{-9}	6.02×10^{-23}	3.18×10^{-23}	5.89	1.50	0.17
9	3.47	1.32	763.8	-1.45×10^{-17}	1.61×10^{-22}	8.13×10^{-24}	1.01	2.23	-0.01
GEO	0.78	-0.62	1125.6	-2.87×10^{-11}	7.5×10^{-22}	*	1.99	0.84	0.37

TABLE II: The parameter values for the S4 binary pulsar hardware injections.

PULSAR	ν_{gw} (Hz)	h_0	T_0 (MJD)	P_b (days)	e	ω_0 (degs)	$a \sin i$ (secs)
10	250.6	1.30×10^{-22}	51749.71156482	1.35405939	0.0	0.0	1.65284
11	188.0	5.21×10^{-22}	52812.92041176	0.31963390	0.180567	322.571	2.7564

3. Calibration Issues

A brief note should be made of the effect of calibrations on the above extracted pulsar hardware injections. The injections were for the most part analysed using exactly the same pipeline as applied to the general known pulsar analysis. However due to the nature of the hardware injections some additional post-processing of the results has had to be applied. To calculate the amplitude and phase of the injections, when applying forces to the interferometer end test masses, a reference calibration must be used. These reference calibrations are different for each interferometer. For both S3 and S4

these reference calibrations differed by small, but not insignificant amounts, from the final calibration used when extracting the signals, meaning that upon extracting the signals the amplitude and phase appear offset from the injected values. As the difference between the reference and final calibrations are different for each interferometer there will also be slight offsets between the extracted parameters between detectors. The extracted signals from each interferometer therefore have had to be adjusted to reflect these differences, determined independently of the hardware injections, and correct them so as to give the same input signal. This allows the combined joint upper limits to be produced.




FIG. 7: The pdfs of h_0 and ϕ_0 for 10 isolated pulsar injections into the LIGO detectors during S3. The anomaly seen in PULSAR7 is discussed in the text.




FIG. 8: The pdfs of h_0 and ϕ_0 for 10 isolated and 2 binary pulsar injections into the LIGO detectors during S4.

APPENDIX B: TABLES OF UPPER LIMIT RESULTS

TABLE III: Pulsar upper limits using LIGO data from the S3 and S4 runs. The approximate pulsar spin frequencies and spin-down rates are given. A * denotes globular cluster pulsars for which no spin-down upper limit could be set. The values marked with a † represent pulsars for which the spin-down limit has been corrected for the Shklovskii effect. The ratio column gives the ratio of our experimental upper limits to the spin-down upper limits.

PULSAR	ν (Hz)	$\dot{\nu}$ (Hz s ⁻¹)	S3						S4						S3 and S4					
			log $h_0^{95\%}$				log ε	ratio	log $h_0^{95\%}$				log ε	ratio	log $h_0^{95\%}$				log ε	ratio
			H1	H2	L1	Joint			H1	H2	L1	Joint			H1	H2	L1	Joint		
J0024-7204C	173.71	$+1.50 \times 10^{-15}$	-23.19	-23.42	-22.96	-23.51	-3.94	*	-22.97	-23.49	-23.21	-23.51	-3.94	*	-23.28	-23.62	-23.25	-23.68	-4.10	*
J0024-7204D	186.65	$+1.20 \times 10^{-16}$	-23.70	-23.60	-23.23	-23.83	-4.32	*	-23.93	-23.88	-23.87	-24.12	-4.61	*	-24.09	-23.94	-23.85	-24.28	-4.76	*
J0024-7204E	282.78	-7.88×10^{-15}	-23.82	-23.45	-23.10	-23.84	-4.69	1631 [†]	-23.93	-23.85	-23.74	-24.07	-4.92	961 [†]	-24.00	-23.91	-23.75	-24.11	-4.96	890 [†]
J0024-7204F	381.16	-9.37×10^{-15}	-23.62	-23.37	-22.92	-23.67	-4.78	2576 [†]	-23.82	-23.62	-23.50	-23.93	-5.04	1420 [†]	-23.94	-23.63	-23.49	-24.05	-5.16	1069 [†]
J0024-7204G	247.50	$+2.58 \times 10^{-15}$	-23.92	-23.54	-23.11	-23.98	-4.71	*	-24.00	-23.90	-23.78	-24.18	-4.91	*	-24.08	-23.94	-23.79	-24.25	-4.98	*
J0024-7204I	286.94	$+3.78 \times 10^{-15}$	-23.72	-23.35	-23.07	-23.77	-4.63	*	-23.95	-23.86	-23.53	-24.01	-4.87	*	-23.94	-23.88	-23.57	-24.00	-4.86	*
J0024-7204J	476.05	$+2.22 \times 10^{-15}$	-23.57	-23.10	-22.78	-23.57	-4.87	*	-23.74	-23.66	-23.32	-23.92	-5.22	*	-23.80	-23.67	-23.32	-23.98	-5.28	*
J0024-7204L	230.09	$+6.46 \times 10^{-15}$	-23.89	-23.49	-23.19	-23.93	-4.59	*	-23.92	-23.89	-23.74	-23.98	-4.65	*	-24.06	-23.94	-23.74	-24.07	-4.74	*
J0024-7204M	271.99	$+2.84 \times 10^{-15}$	-23.85	-23.44	-23.01	-23.86	-4.68	*	-23.96	-23.90	-23.76	-24.11	-4.92	*	-24.02	-23.95	-23.73	-24.15	-4.97	*
J0024-7204N	327.44	$+2.34 \times 10^{-15}$	-23.74	-23.37	-23.02	-23.77	-4.74	*	-23.79	-23.72	-23.65	-24.04	-5.01	*	-23.94	-23.83	-23.64	-24.15	-5.12	*
J0024-7204Q	247.94	-2.09×10^{-15}	-23.86	-23.42	-23.19	-23.95	-4.69	2274	-23.98	-23.91	-23.63	-24.06	-4.80	1777	-24.07	-23.95	-23.64	-24.17	-4.90	1382
J0024-7204S	353.31	$+1.50 \times 10^{-14}$	-23.57	-23.31	-23.00	-23.61	-4.65	*	-23.86	-23.79	-23.57	-24.04	-5.08	*	-23.91	-23.84	-23.57	-24.04	-5.08	*
J0024-7204T	131.78	-5.10×10^{-15}	-24.09	-23.62	-23.39	-24.12	-4.31	718	-24.19	-24.08	-24.04	-24.36	-4.55	418	-24.24	-24.10	-24.06	-24.40	-4.59	377
J0024-7204U	230.26	-5.05×10^{-15}	-23.70	-23.53	-23.20	-23.79	-4.46	2079 [†]	-23.96	-23.94	-23.68	-24.18	-4.85	849 [†]	-23.91	-23.99	-23.69	-24.13	-4.80	943 [†]
J0034-0534	532.71	-1.41×10^{-15}	-23.22	-23.12	-22.68	-23.39	-5.48	3017	-23.39	-23.36	-23.41	-23.72	-5.81	1416	-23.48	-23.41	-23.40	-23.79	-5.87	1224
J0218+4232	430.46	-1.43×10^{-14}	-23.53	-23.29	-22.88	-23.61	-4.73	3117	-23.73	-23.48	-23.49	-23.79	-4.92	2024	-23.83	-23.57	-23.52	-23.88	-5.01	1643
J0534+2200	29.80	-3.73×10^{-10}	-23.09	-22.04	-22.37	-23.12	-2.39	5.31 [†]	-23.29	-23.08	-22.83	-23.36	-2.63	3.07 [†]	-23.37	-23.08	-22.85	-23.42	-2.69	2.68 [†]
J0613-0200	326.60	-1.02×10^{-15}	-23.66	-23.26	-23.02	-23.70	-5.01	3205 [†]	-23.67	-23.70	-23.67	-23.93	-5.24	1887 [†]	-23.85	-23.74	-23.68	-24.00	-5.31	1619 [†]
J0621+1002	34.66	-5.68×10^{-17}	-23.50	-22.56	-22.91	-23.54	-2.97	5354 [†]	-23.78	-23.36	-23.81	-24.04	-3.47	1700 [†]	-23.83	-23.36	-23.82	-24.04	-3.47	1680 [†]
J0711-6830	182.11	-4.94×10^{-16}	-23.90	-23.21	-23.19	-23.94	-5.07	1203 [†]	-23.98	-23.92	-23.82	-24.10	-5.23	837 [†]	-24.13	-23.97	-23.81	-24.23	-5.36	626 [†]
J0737-3039A	44.05	-3.38×10^{-15}	-23.96	-23.13	-23.13	-23.95	-4.11	91	-24.08	-23.75	-23.99	-24.22	-4.38	49	-24.20	-23.77	-23.99	-24.27	-4.43	43
J0751+1807	287.46	-6.43×10^{-16}	-23.77	-23.39	-23.14	-23.82	-5.06	2521	-23.74	-23.73	-23.65	-23.91	-5.15	2057	-23.86	-23.79	-23.63	-23.98	-5.22	1747
J1012+5307	190.27	-6.20×10^{-16}	-23.96	-23.61	-23.23	-23.97	-5.44	442 [†]	-24.11	-23.96	-23.85	-24.29	-5.76	212 [†]	-24.20	-24.01	-23.88	-24.35	-5.82	185 [†]
J1022+1001	60.78	-1.60×10^{-16}	-24.06	-22.87	-23.05	-24.07	-4.78	197	-23.96	-23.78	-24.13	-24.24	-4.96	131	-24.13	-23.78	-24.14	-24.31	-5.03	113
J1024-0719	193.72	-6.95×10^{-16}	-23.77	-23.40	-23.34	-23.88	-5.54	302	-23.87	-23.85	-23.87	-24.12	-5.78	174	-23.99	-23.89	-23.89	-24.22	-5.88	139
J1045-4509	133.79	-3.16×10^{-16}	-23.91	-23.52	-23.39	-23.95	-4.31	3311 [†]	-24.08	-23.93	-23.91	-24.21	-4.58	1816 [†]	-24.18	-23.94	-23.95	-24.20	-4.57	1832 [†]
J1300+1240	160.81	-2.95×10^{-15}	-23.90	-23.46	-23.37	-23.93	-5.08	863 [†]	-23.83	-23.79	-24.01	-24.05	-5.21	643 [†]	-23.92	-23.80	-24.00	-24.08	-5.23	606 [†]
J1435-6100	106.98	-2.80×10^{-16}	-24.17	-23.58	-23.46	-24.20	-4.37	1574	-24.16	-24.00	-24.08	-24.39	-4.56	1024	-24.33	-24.09	-24.11	-24.47	-4.64	850
J1455-3330	125.20	-3.81×10^{-16}	-24.05	-23.37	-23.26	-24.07	-5.02	562 [†]	-24.17	-24.01	-23.87	-24.29	-5.24	338 [†]	-24.27	-23.99	-23.88	-24.34	-5.29	299 [†]
J1518+0205A	180.06	-1.34×10^{-15}	-23.64	-22.59	-22.81	-23.67	-3.91	7627	-23.47	-23.69	-23.77	-23.85	-4.10	5007	-23.76	-23.69	-23.77	-23.94	-4.19	4030

TABLE III cont.

			S3					S4					S3 and S4									
			$\log h_0^{95\%}$				$\log \varepsilon$	ratio	$\log h_0^{95\%}$				$\log \varepsilon$	ratio	$\log h_0^{95\%}$				$\log \varepsilon$	ratio		
PULSAR	ν (Hz)	$\dot{\nu}$ (Hz s $^{-1}$)	H1	H2	L1	Joint		H1	H2	L1	Joint		H1	H2	L1	Joint		H1	H2	L1	Joint	
J1537+1155	26.38	-1.69×10^{-15}	-22.79	-22.00	-22.19	-22.79	-2.31	2285 †	-23.35	-22.92	-23.30	-23.57	-3.08	383 †	-23.36	-22.92	-23.31	-23.58	-3.10	369 †		
J1603-7202	67.38	-7.10×10^{-17}	-24.23	-23.50	-23.41	-24.25	-4.32	1318 †	-24.30	-23.87	-24.05	-24.38	-4.45	978 †	-24.42	-23.92	-24.05	-24.50	-4.57	743 †		
J1623-2631	90.29	-5.47×10^{-15}	-23.94	-23.34	-23.43	-23.98	-4.17	383 †	-24.03	-23.73	-24.05	-24.13	-4.32	273 †	-24.16	-23.76	-24.05	-24.36	-4.55	161 †		
J1629-6902	166.65	-2.78×10^{-16}	-24.03	-23.60	-23.29	-24.10	-5.04	1041	-24.07	-24.01	-23.97	-24.26	-5.20	711	-24.24	-24.05	-23.97	-24.36	-5.30	567		
J1640+2224	316.12	-2.83×10^{-16}	-23.58	-23.38	-22.97	-23.59	-5.14	5329 †	-23.74	-23.74	-23.67	-23.96	-5.51	2277 †	-23.81	-23.76	-23.67	-23.99	-5.55	2079 †		
J1643-1224	216.37	-8.66×10^{-16}	-23.69	-23.26	-23.18	-23.77	-4.38	5776 †	-23.88	-23.76	-23.82	-23.95	-4.56	3821 †	-23.93	-23.76	-23.88	-23.97	-4.58	3616 †		
J1701-3006A	190.78	$+4.80 \times 10^{-15}$	-23.96	-23.37	-23.33	-23.93	-4.28	*	-24.01	-23.83	-23.89	-24.17	-4.52	*	-24.15	-23.82	-23.91	-24.29	-4.64	*		
J1701-3006B	278.25	$+2.71 \times 10^{-14}$	-23.84	-23.41	-23.07	-23.85	-4.52	*	-23.92	-23.79	-23.55	-23.99	-4.67	*	-24.03	-23.82	-23.56	-24.09	-4.77	*		
J1701-3006C	262.71	$+2.20 \times 10^{-15}$	-23.82	-23.38	-23.14	-23.80	-4.42	*	-23.86	-23.82	-23.73	-24.04	-4.66	*	-24.03	-23.83	-23.72	-24.12	-4.74	*		
J1713+0747	218.81	-4.08×10^{-16}	-23.62	-23.18	-23.14	-23.60	-4.85	2663 †	-23.90	-23.87	-23.78	-24.05	-5.31	938 †	-23.95	-23.84	-23.77	-24.10	-5.35	838 †		
J1744-1134	245.43	-5.39×10^{-16}	-23.78	-23.39	-23.26	-23.84	-5.69	483 †	-23.72	-23.80	-23.84	-23.97	-5.82	353 †	-23.86	-23.80	-23.86	-24.04	-5.89	301 †		
J1745-0952	51.61	-2.53×10^{-16}	-23.94	-23.29	-23.28	-23.95	-3.63	1482	-24.19	-23.77	-24.09	-24.30	-3.98	662	-24.21	-23.78	-24.09	-24.33	-4.01	623		
J1748-2446A	172.96	$+2.54 \times 10^{-16}$	-24.10	-23.56	-23.50	-24.11	-3.92	*	-	-	-	-	-	-	-	-	-	-	-	-		
J1748-2446C	118.54	$+8.52 \times 10^{-15}$	-24.08	-23.10	-23.46	-24.09	-3.92	*	-24.15	-23.89	-24.11	-24.36	-4.20	*	-24.28	-23.90	-24.12	-24.41	-4.25	*		
J1756-2251	35.14	-1.26×10^{-15}	-23.63	-22.76	-22.91	-23.63	-2.88	1422	-23.79	-23.29	-23.60	-23.78	-3.04	997	-23.81	-23.30	-23.60	-23.79	-3.05	976		
J1757-5322	112.74	-3.34×10^{-16}	-24.10	-23.44	-23.41	-24.14	-4.73	715	-24.12	-24.04	-24.07	-24.35	-4.95	437	-24.22	-24.04	-24.06	-24.38	-4.98	409		
J1801-1417	275.85	-4.02×10^{-16}	-23.66	-23.32	-23.11	-23.76	-5.01	3224	-23.83	-23.74	-23.79	-24.04	-5.29	1686	-23.91	-23.78	-23.78	-24.10	-5.36	1456		
J1802-2124	79.07	-4.50×10^{-16}	-24.02	-23.41	-23.45	-24.01	-3.91	1704	-24.12	-24.00	-24.10	-24.32	-4.22	836	-24.17	-23.99	-24.11	-24.25	-4.15	980		
J1804-0735	43.29	-8.75×10^{-16}	-23.82	-23.13	-23.08	-23.82	-2.79	3532	-24.04	-23.65	-23.99	-24.19	-3.17	1490	-24.15	-23.67	-24.00	-24.24	-3.21	1336		
J1804-2717	107.03	-4.68×10^{-16}	-24.03	-23.46	-23.46	-24.00	-4.62	686	-24.12	-24.03	-24.13	-24.31	-4.93	340	-24.25	-24.05	-24.13	-24.32	-4.94	329		
J1807-2459A	326.86	$+4.87 \times 10^{-16}$	-23.73	-23.30	-23.02	-23.79	-5.01	*	-23.84	-23.70	-23.74	-24.01	-5.23	*	-23.95	-23.75	-23.74	-24.08	-5.31	*		
J1810-2005	30.47	-1.40×10^{-16}	-23.17	-22.15	-22.61	-23.19	-2.18	15134	-23.37	-23.15	-23.34	-23.62	-2.61	5612	-23.50	-23.14	-23.34	-23.64	-2.63	5342		
J1823-3021A	367.65	-1.14×10^{-13}	-23.75	-23.43	-23.31	-23.78	-4.04	648	-	-	-	-	-	-	-	-	-	-	-	-		
J1824-2452	327.41	-1.74×10^{-13}	-23.78	-23.34	-23.01	-23.80	-4.77	416 †	-23.78	-23.70	-23.63	-23.99	-4.96	269 †	-23.93	-23.77	-23.65	-24.08	-5.04	221 †		
J1843-1113	541.81	-2.82×10^{-15}	-23.26	-23.09	-22.71	-23.32	-5.12	5167	-23.51	-23.48	-23.40	-23.75	-5.55	1887	-23.57	-23.52	-23.40	-23.80	-5.60	1690		
J1857+0943	186.49	-6.20×10^{-16}	-23.74	-23.42	-23.32	-23.75	-4.96	1117 †	-23.88	-23.76	-23.94	-24.18	-5.39	412 †	-23.96	-23.78	-23.96	-24.21	-5.42	388 †		
J1905+0400	264.24	-3.39×10^{-16}	-23.76	-23.23	-23.18	-23.76	-5.10	2556	-23.80	-23.68	-23.75	-23.89	-5.24	1877	-23.95	-23.76	-23.77	-24.01	-5.35	1440		
J1909-3744	339.32	-1.61×10^{-15}	-23.74	-23.34	-22.95	-23.81	-5.59	1095 †	-23.78	-23.77	-23.63	-23.95	-5.72	805 †	-23.85	-23.83	-23.67	-24.05	-5.83	628 †		
J1910-5959A	306.17	-2.88×10^{-16}	-23.83	-23.31	-23.04	-23.90	-4.89	6488	-23.78	-23.60	-23.69	-24.01	-5.01	4974	-23.96	-23.77	-23.71	-24.12	-5.12	3863		
J1910-5959B	119.65	$+1.14 \times 10^{-14}$	-24.15	-23.03	-23.26	-24.16	-4.34	*	-23.61	-23.94	-23.99	-24.13	-4.31	*	-24.17	-23.93	-24.00	-24.29	-4.47	*		

TABLE III cont.

			S3						S4						S3 and S4					
			$\log h_0^{95\%}$			$\log \varepsilon$	ratio	$\log h_0^{95\%}$			$\log \varepsilon$	ratio	$\log h_0^{95\%}$			$\log \varepsilon$	ratio			
PULSAR	ν (Hz)	$\dot{\nu}$ (Hz s $^{-1}$)	H1	H2	L1	Joint			H1	H2	L1	Joint			H1	H2	L1	Joint		
J1910-5959C	189.49	-7.90×10^{-17}	-24.15	-23.03	-23.26	-24.16	-4.34	*	-24.08	-23.98	-23.89	-24.18	-4.76	*	-24.20	-24.01	-23.91	-24.30	-4.88	*
J1910-5959D	110.68	-1.18×10^{-14}	-23.95	-23.54	-23.31	-23.97	-4.55	8176	-24.28	-24.00	-24.07	-24.39	-4.50	198	-24.33	-24.02	-24.05	-24.39	-4.50	198
J1910-5959E	218.73	$+2.09 \times 10^{-14}$	-23.92	-23.58	-23.24	-23.97	-4.67	*	-24.01	-23.90	-23.87	-24.22	-4.93	*	-24.09	-23.94	-23.88	-24.25	-4.96	*
J1911+0101A	276.36	$+5.03 \times 10^{-16}$	-23.56	-23.31	-23.07	-23.72	-4.36	*	-23.80	-23.73	-23.75	-24.03	-4.67	*	-23.95	-23.75	-23.74	-24.11	-4.75	*
J1911+0101B	185.72	$+6.90 \times 10^{-17}$	-23.65	-23.37	-23.27	-23.73	-4.03	*	-23.91	-23.89	-23.94	-24.14	-4.44	*	-23.91	-23.94	-23.91	-24.10	-4.39	*
J1911-1114	275.81	-1.08×10^{-15}	-23.75	-23.37	-23.02	-23.76	-5.06	2617 †	-23.87	-23.72	-23.67	-24.00	-5.31	1504 †	-23.94	-23.77	-23.66	-24.06	-5.37	1308 †
J1939+2134	641.93	-4.33×10^{-14}	-23.14	-23.11	-22.49	-23.30	-4.98	2733 †	-23.40	-23.38	-23.29	-23.63	-5.31	1280 †	-23.46	-23.43	-23.27	-23.71	-5.39	1063 †
J1955+2908	163.05	-7.91×10^{-16}	-23.78	-23.51	-23.32	-23.84	-4.16	4497 †	-24.10	-23.87	-23.94	-24.23	-4.55	1837 †	-24.16	-23.89	-23.93	-24.30	-4.62	1558 †
J1959+2048	622.12	-6.52×10^{-15}	-23.21	-23.00	-22.56	-23.28	-5.31	3769 †	-23.52	-23.33	-23.41	-23.68	-5.71	1481 †	-23.56	-23.34	-23.42	-23.72	-5.75	1363 †
J2019+2425	254.16	-4.54×10^{-16}	-23.72	-23.40	-23.16	-23.83	-5.31	2075 †	-23.86	-23.80	-23.74	-24.07	-5.55	1189 †	-23.89	-23.82	-23.72	-24.10	-5.58	1114 †
J2051-0827	221.80	-6.27×10^{-16}	-23.84	-23.40	-23.16	-23.91	-5.12	1187 †	-23.82	-23.74	-23.73	-23.99	-5.20	983 †	-23.96	-23.77	-23.73	-24.07	-5.28	814 †
J2124-3358	202.79	-8.45×10^{-16}	-23.95	-23.50	-23.26	-23.96	-5.81	206 †	-23.90	-23.85	-23.89	-24.10	-5.94	151 †	-24.11	-23.94	-23.89	-24.23	-6.07	112 †
J2129-5721	268.36	-1.49×10^{-15}	-23.78	-23.40	-23.15	-23.84	-4.92	2014 †	-23.94	-23.86	-23.79	-24.11	-5.19	1087 †	-23.98	-23.85	-23.79	-24.13	-5.20	1038 †
J2140-2310A	90.75	$+4.27 \times 10^{-16}$	-24.11	-23.38	-23.45	-24.12	-3.71	*	-24.21	-23.91	-24.01	-24.35	-3.94	*	-24.33	-23.91	-24.04	-24.43	-4.02	*
J2145-0750	62.30	-1.15×10^{-16}	-24.06	-23.08	-23.40	-24.09	-4.60	400 †	-24.17	-23.81	-24.17	-24.37	-4.89	208 †	-24.25	-23.82	-24.18	-24.40	-4.92	195 †
J2229+2643	335.82	-1.65×10^{-16}	-23.57	-23.26	-23.02	-23.61	-5.13	6185	-23.72	-23.49	-23.64	-23.86	-5.38	3497	-23.84	-23.57	-23.64	-24.03	-5.55	2357
J2317+1439	290.25	-2.04×10^{-16}	-23.46	-23.33	-23.10	-23.58	-4.85	9406 †	-23.80	-23.77	-23.71	-24.04	-5.31	3238 †	-23.83	-23.78	-23.73	-24.06	-5.33	3091 †
J2322+2057	207.97	-4.20×10^{-16}	-23.79	-23.41	-23.18	-23.93	-5.30	1229 †	-23.92	-23.83	-23.81	-24.11	-5.48	812 †	-24.00	-23.87	-23.87	-24.17	-5.54	702 †

TABLE IV: Upper limits including GEO 600 for the two fastest pulsars in the analysis.

			S3						S4						S3 and S4								
			$\log h_0^{95\%}$			$\log \varepsilon$	ratio	$\log h_0^{95\%}$			$\log \varepsilon$	ratio	$\log h_0^{95\%}$			$\log \varepsilon$	ratio						
PULSAR	ν (Hz)	$\dot{\nu}$ (Hz s $^{-1}$)	H1	H2	L1	GEO	Joint			H1	H2	L1	GEO	Joint			H1	H2	L1	GEO	Joint		
J1843-1113	541.81	-2.82×10^{-15}	-23.26	-23.09	-22.71	*	-23.32	-5.12	5167	-23.51	-23.48	-23.40	-22.42	-23.75	-5.55	1887	-23.57	-23.52	-23.40	-22.43	-23.80	-5.60	1690
J1939+2134	641.93	-4.33×10^{-14}	-23.14	-23.11	-22.49	-22.07	-23.30	-4.98	2733 †	-23.40	-23.38	-23.29	-22.59	-23.63	-5.31	1280 †	-23.46	-23.43	-23.27	-22.60	-23.71	-5.39	1063 †