



# Equilibrium Solubility In Simulated Intestinal Media

Orbito Team – Strathclyde University

WP 1

September 2018


# Work Package 1

- Task 1.1 Design of Experiment
- Task 1.3 Design and characterisation of a set of SGIM reflecting the compositions of human GI fluids in the fasted and fed states
- Task 1.4 Optimisation and validation of novel small scale models for dissolution rate and apparent solubility in the designed SGIM
- Task 1.11 Determination of excipient effects on API behaviour in developed in vitro models

# Coverage


- Media Parameters
- Basic Experimental Protocol
- Design of Experiment Studies
- Four Component Mixture Design Studies
  - Intestinal Solubility Topography
- Practical Application
  - BCS, Intestinal Solubility Window, PK Prediction
- Experimental Caveats
- Conclusions

# Media Parameters




**Fig. 4.** (a) Bile salt (BS) concentration, (b) phospholipid (PL) concentration, and (c) BS/PL ratio in fasted gastric, duodenal and jejunal fluids. Box-whisker plots show minimum and maximum values, as well as 25, 50 and 75 percentile. The cross indicates the mean value. Each data point represents a group of participants ( $n = 1-10$  colored red,  $n = 11-20$  colored blue and  $n > 20$  colored green) reported in one publication. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Bergstrom et.al., (2014) Eur.J.Pharm.Sci., 57; 173-1999



# Equilibrium Solubility Measurement





Khadra, I., et.al., (2015) Eur.J.Pharm.Sci., 67; 67-75  
 Zhou, Z., et.al., (2017) Eur.J.Pharm.Sci., 99; 95-104

# Media Parameters - Design of Experiment

- Fasted
  - Fractional factorial design (quarter)
  - 7 parameters
  - 66 experiments (including duplicates)
- Fed
  - D-optimal design
  - 8 parameters
  - 94 experiments (including duplicates)



Parameter/ Ingredient	Substance	Fasted		Fed		Full Range	
		Lower Limit	Upper limit	Lower Limit	Upper Limit	Lower Limit	Upper Limit
Bile salt (mM)	Sodium TC	1.5	5.9	3.6	24	1.5	24
Lecithin (mM)	Egg PL	0.2	0.75	0.5	4.8	0.2	4.8
Buffer (mM)	NaH2PO4	15	45	-	-	15	45
	Maleic acid	-	-	28.6	58.09	-	-
Salt (mM)	NaCl	68	106	125	203	68	203
pH	NaOH/HCl	5	7	5	7	5	7
Enzyme (U/ml) <sup>1</sup>	Pancreatin	0.5	1	100	150	-	-
Fatty acid (mM)	Sodium oleate	0.5	10	0.8	52	0.5	52
Monoglyceri de (mM)	Glyceryl Mono- oleate	-	-	0.5	6.5	0.5	6.5

# Fasted - Design of Experiment





Khadra, I., et.al., (2015) Eur.J.Pharm.Sci., 67; 67-75  
See also – Madsen, C., et.al., (2018) Eur.J.Pharm.Sci., 111; 311-319

Unpublished Data



# Multiple – Design of Experiment Systems

- Fasted
- Fed
- Combined
- Dual range
- 9 point Fasted and Fed



Khadra, I., et.al., (2015) Eur.J.Pharm.Sci., 67; 67-75  
 Zhou, Z., et.al., (2017) Eur.J.Pharm.Sci., 99; 95-104  
 Perrier, J., et.al., (2018) Eur.J.Pharm.Sci., 111; 247-256  
 Ainousah, B.E., et.al., (2017) Mol.Pharmaceutics 14; 4170-4180  
 9 Point – unpublished data

# Impact DoE Design on Factor Significance

	Fasted	Fed	Dual Level – Fasted	Dual Level – Fed	9DOE Fasted	9DOE Fed
<b>Indomethacin</b>	pH Bile Salt Buffer Oleate	pH Oleate Bile Salt	pH	pH	pH	pH
<b>Aprepitant</b>	Oleate pH Lecithin	Oleate Bile Salt pH	Oleate Lecithin MG	NS	NS	NS
<b>Tadalafil</b>	Bile Salt pH Buffer Lecithin Oleate Salt	Bile Salt Oleate	NS	NS	pH	NS
<b>Zafirlukast</b>	pH Oleate Bile Salt Lecithin	pH Bile Salt Oleate	pH Cholesterol MG	NS	NS	NS
<b>Carvedilol</b>	Bile Salt Oleate	Bile Salt pH Buffer Oleate	NS	NS	Bile Salt pH	NS
<b>Felodipine</b>	pH Oleate Lecithin Bile Salt	Oleate Bile Salt pH Lecithin	pH Oleate Lecithin MG	Oleate Lecithin Bile Salt MG	Oleate	Bile Salt
<b>Probucol</b>	pH Oleate	Bile Salt MG Oleate Lecithin pH	Oleate BS:PL	Oleate	pH	NS

Smaller Design = Lower  
Factor Resolution

# Four Component Mixture Design (4CMD)

- DoE – indicates drug specific behaviour

Difficult to visualise

- Four components

Bile salt

Sodium oleate

Lecithin



Glyceryl monooleate

- Total amphiphile concentration

11.7 mM


- pH 7.4

Tetrahedron of solubility  
Each Triangle – Ternary Phase Diagram




# 4CMD pH 7.4 TAC 11.7 mM


Aprepitant




Carvedilol




Zafirlukast




Fenofibrate



Felodipine




Spironolactone




Zhou, Z., et.al., (2017) Mol.Pharmaceutics, 14; 4132-4144

# 4CMD Matrix

- Indomethacin




Unpublished Data





# Relationship with DoE

- Indomethacin





# Practical Application - BCS & DCS



# Practical Application - Compartmental Analysis


- 4CMD represents 9 possible intestinal compartments



Unpublished Data


# Practical Application – Prediction

## Solubility



Unpublished Data

## Pharmacokinetics



# Experimental Caveats – Factor Concentrations





Figure 8. Overall variability in the composition of HIF expressed as a function of time from 10 volunteers, presented as individual data points. Mean and median values are displayed using a solid or dashed line, respectively. The exact value for mean (above the solid line) and median (above the dashed line) are displayed below their respective graphs, in the same unit of measurement as used in the graph.

Riethorst, D., et.al., (2015) J.Pharm.Sci., 105;673-681

Cholesterol

Fasted analysis only

X




Khadra, I., et.al., (2015) Eur.J.Pharm.Sci., 67; 67-75

# Experimental Caveats – Factor Ratios

- 4CMD Amphiphile number & amphiphile ratio

Fenofibrate



# Conclusions

- Single media composition solubility measurement (SIF or HIF) – limited value
    - Position within topography unknown, variability unknown
  - Increase number of measurement points – increased information
    - Big DoE – factor and solubility range information
    - Wee DoE – solubility range – with caveats – limited or no factor information
  - Choice of media components – the greater the better – see below
    - Number, Concentration and Ratio – appropriate
  - Beware – drug specific behavior
- 
- Improved statistical information – HIF composition and SIF solubility
  - Structured solubility behavior is present
  - Solubility prediction – looking possible – limited initially

# Acknowledgment

- Strathclyde OrBiTo Team
  - Gavin Halbert, Clive Wilson, Ibrahim Khadra, Claire Dunn, Jeremy Perrier, Bayan Ainousah
- WP1 Team & entire OrBiTo Team – Management and all
- IMI for funding
- Mr Jeremy Perrier and Gavin Halbert
  - Funded by Cancer Research UK
- Ms Bayan Ainousah
  - Funded by Saudi Government

Thank you – for listening