

Alkali Metal Effects in Trans-Metal-Trapping (TMT): Comparing LiTMP with NaTMP in Cooperative MTMP/Ga(CH₂SiMe₃)₃ Metalation Reactions

Ross McLellan

Marina Uzelac

Leonie J. Bole


Jose María Gil-Negrete

David R. Armstrong

Alan R. Kennedy

Robert E. Mulvey*

Eva Hevia*



WestCHEM, Department of Pure and Applied Chemistry, University of Strathclyde, Glasgow G1 1XL, UK
 r.e.mulvey@strath.ac.uk
 eva.hevia@strath.ac.uk

Published as part of the 50 Years SYNTHESIS – Golden Anniversary Issue

Received: 30.11.2018

Accepted: 05.12.2018

Published online: 15.01.2019

DOI: 10.1055/s-0037-1611646; Art ID: ss-2018-z0802-fa

License terms:

Abstract Stepwise metalation and trapping, so called trans-metal-trapping (TMT), of anisole is studied using LiTMP as base and Ga(CH₂SiMe₃)₃ as trap. The isolated 'trapped' intermediate is also assessed in C–C bond forming reactions, highlighting the inherent advantages and remaining challenges of this system. The same base trap mixture is found to metallate N–Me bonds of the diamines TMEDA and PMDETA. Comparative studies replacing LiTMP by NaTMP have found significant alkali metal effects on the extent of both base-trap cocomplexation and onward reactivities of TMT products.

Key words metalation, gallium trans-metal-trapping, carbanions, cooperative effects, lithium, sodium

Deprotonative metalation (C–H to C–M exchange) of aryl and heteroaryl substrates is a widely successful tool, utilised in the construction of important organic molecules, usually producing organometallic intermediates primed for onward reactivity. Longstanding reagents of choice in this context are alkyllithiums and lithium amides.¹ Synthetic drawbacks to these metalation transformations are commonly, poor functional group tolerance (at convenient temperatures) and selectivity, necessitating cryogenic reaction conditions to prevent unwanted side reactions or decomposition of the lithiated species. One solution to these limitations is to employ metal salts such as MgCl₂ or ZnCl₂·2LiCl as in situ trapping agents during LiTMP metalations of arenes and N-heterocycles in THF, in reactions that have been performed under continuous flow conditions (TMP = 2,2,6,6-tetramethylpiperidine).² A second approach is to

use bimetallic systems containing two metals of different polarity within the same molecule. These systems can be shown to behave in concert, resulting in higher selectivity and in many cases using ambient reaction conditions during metalation. Important examples of bimetallic formulations used in deprotonation reactions are the classic Lochmann–Schlosser superbases,³ Knochel's turbo Hauser reagents,⁴ and Uchiyama and Mongin's TMP zincates.⁵ This bimetallic approach, combining an alkali metal with a less electropositive metal (e.g., Mg or Zn) into an 'ate' complex has been termed alkali-metal-mediated metalation and, a series of reactivity and structural studies have demonstrated that the high reactivity of the alkali metal can be harnessed by the less polar metal, while retaining the advantages of selectivity that the non-alkali metal provides, thus representing a 'best of both worlds' scenario.⁶ Recently we have shown in our Strathclyde research groups that such prodigious metal...metal cooperative effects can, in certain cases work in sequence in a process that has been coined trans-metal-trapping (TMT).⁷ A general depiction of TMT is given in Scheme 1, and describes a process reliant upon the stepwise reactivity of the two metal organic reagents with the aromatic substrate. Deprotonation of the substrate with a lithium amide base exists in a pK_a dependent equilibrium that typically lies towards the lithium amide and unreacted aromatic substrate, meaning that when used in isolation, a stoichiometric amount of lithium amide only delivers ca. <10% of product, when intercepted with an electrophile such as iodine.

However, in the presence of the Lewis acidic carbophilic trap, the lithiated intermediate can be readily intercepted, resulting in a so called crossover bimetallic complex that stops short of full transmetalation, where separated lithium

Biographical Sketches



Ross McLellan received his M.Chem. degree in chemistry from Heriot-Watt University in 2007. He obtained his Ph.D. degree from the same institution in 2011 under the supervision of Professor

Alan Welch. His first postdoctoral position with Dr Scott Dalgarno focussed on the controlled formation and magnetism of 3d/4f polynuclear metal clusters in collaboration with Professor Euan

Brechin. He is currently a postdoctoral research fellow at the University of Strathclyde, where he is interested in developing main group systems and investigating their use in catalysis.



Marina Uzelac obtained her B.Sc. and M.Sci. in chemistry from the University of Zagreb, Croatia. She then moved to the UK, to carry out her Ph.D. studies at the University of Strathclyde under the supervision of Professor Eva Hevia developing new gallium-mediated transfor-

mations for C–H functionalisation and small molecule activation processes. She graduated in 2016 and then took up a postdoctoral position working in a collaborative project with Professors Robert Mulvey and Eva Hevia exploiting multi-component s-block metal reagents in

deprotonative metalation and C–C bond forming processes. In 2018 Marina joined the group of Professor Michael J. Ingleson working on the development of new organozinc catalysts with application in bond forming processes.



Leonie J. Bole obtained her undergraduate degree in 2016 from the University of Strathclyde completing her M.Sc. on

inverse crown chemistry under the supervision of Dr Charles T. O'Hara. She is currently working towards her Ph.D. in

main group organometallic chemistry with Prof. Eva Hevia and Dr Charles T. O'Hara at the University of Strathclyde.



Jose Maria Gil-Negrete obtained both his B.Sc. and M.Sc. from the University of A Coruña, Spain. Since 2014 he is pursuing a Ph.D. under the direction of Professors Luis Sarandeses and Jose Pérez

Setstelo, also at the University of A Coruña. His research is focussed on the development of new applications for indium organometallics in organic synthesis. During the summer of 2017, Jose

Maria was a visiting researcher in the University of Strathclyde, Glasgow, within the group of Professor Eva Hevia.



David Armstrong was born in the North-East of England in County Durham and educated at Newcastle University for both his B.Sc. and Ph.D. degrees. The latter was carried out under the supervision of Peter Perkins and was a theoretical in-

vestigation of Group III compounds. Appointed at the University of Strathclyde in 1970 as a lecturer and subsequently promoted to a senior lectureship, he currently holds an emeritus position. David has been performing theoretical calcula-

tions for about 50 years, publishing over two hundred research papers with his most recent work focussing on different aspects of polar organometallic chemistry.



Alan Kennedy was awarded both B.Sc. and Ph.D. (K. Muir and R. Cross) degrees in chemistry by the University of Glasgow (1986–1993). He has worked at

the University of Strathclyde since 1994 and is a crystallographer specialising in structure-property relationships in small molecules, notably pharmaceuticals, co-

lourants, and metal complexes. He has published over 550 career papers.



Robert Mulvey was born in Glasgow (Scotland) in 1959 and did his B.Sc. and Ph.D. (R. Snaith) at the University of Strathclyde (1977–1984). Following two years postdoctoral research at the University of Durham (K. Wade), he returned to Strathclyde in 1986 as a Royal Society Research Fellow, and was pro-

moted to Professor in 1995. His research has mainly focused on alkali, alkali-earth, and early p-block metals with emphasis on synergistic effects in bimetallic systems. Prof. Mulvey has received many distinguished prizes including the RSC Meldola Medal, RSC Main Group Element Award, Royal Society Wolfson Research


Merit Award, Royal Society Leverhulme Trust Senior Research Fellowship, GDCh Arfvedson Schlenk Prize, and a Humboldt Research Award. A Fellow of both the Royal Society of Edinburgh and Royal Society of Chemistry, he has published approaching 300 papers.



Eva Hevia received her M.Sci. degree in chemistry from the Universidad de Oviedo (Spain) in 1998. She obtained her Ph.D. degree from the same institution in 2002 under the supervision of Professor Victor Riera and Dr Julio Perez. Next she was awarded a Marie Curie Fellowship held at the University of Strathclyde under the direction of Prof. Robert Mul-

vey. In 2006 she took up a Royal Society University Research Fellowship at the University of Strathclyde where she is currently a Professor of Inorganic Chemistry. Her research interests include s-block metal-mediated transformations with a particular emphasis on green chemistry and catalysis. She has published over 120 peer-reviewed journal ar-


ticles and her research has been recognised with several awards including the Royal Society of Chemistry (RSC) Harrison-Meldola Prize (2009), SRUK Emerging Investigator Awards (2016), and, most recently, the RSC Corday-Morgan Prize (2017). In 2018 Eva was elected as Fellow of the Royal Society of Edinburgh.




and non-lithium metal products would form. A key aspect of this chemistry proven with lithium is that the base and the trap do not combine to form a bimetallic complex that would be inert towards the substrate. Functioning only on emergence of the deprotonated substrate (carbanion), the trap inserts into the Li–C bond and drives the equilibrium towards the desired metallated substrate.

In this article we present a deeper understanding of key mechanistic insights of trans-metal-trapping (TMT), by combining structural, reactivity, and theoretical studies, using anisole, a classical substrate in directed *ortho*-metalation, as a case study. A key feature of our exploratory studies of TMT is the stepwise reactivity exhibited by the two organometallic constituents. In this respect the bulky tris(trimethylsilylmethyl)gallium has been demonstrated as an excellent trap for diazine and fluoroaromatic anions, that when metallated by conventional bases are prone to rapid decomposition.⁸ Therefore this study aims to highlight the benefits of LiTMP/Ga(CH₂SiMe₃)₃ in metalation. From a synthetic perspective, liquid Ga(CH₂SiMe₃)₃ (hereafter GaR₃) possesses good hydrocarbon solubility (similar to *i*Bu₂AlTMP another successful trapping agent)⁷ giving it a decided advantage over salt traps (e.g., MgCl₂, ZnCl₂),⁹ which generally need the use of ethereal solvents (usually THF) and often require low temperatures to avoid competing salt metathesis reactions. More importantly, gallium is characterised by strong carbophilicity and is therefore well equipped to sedate ultra-sensitive anions post metalation with LiTMP. This fact is well illustrated in our comparative studies into the metalation of fluoroarenes, where it was shown that the resultant gallium TMT product had far superior stability against decomposition (via benzyne formation) than the aluminium counterpart.^{8b} The poor stability may also be attributed to the fluorophilicity of aluminium meaning that aluminium traps may in general be incompatible with fluorinated substrates. Apart from these intrinsic properties of GaR₃, a key specification of the trapping reagent, as demonstrated with *i*Bu₂AlTMP previously,⁷ is its

inertness towards co-complexation with LiTMP, and therefore our first step was to establish whether GaR₃ is bulky enough to compromise its ability to form a weakly basic, coordinatively saturated ate complex with LiTMP. Such separation of the organometallic reagents appears a prerequisite for an effective TMT process, and this is underlined by the fact that LiTMP/*i*Bu₂AlTMP mixtures are better TMT agents than LiTMP/*i*Bu₃Al mixtures, which form the aluminate LiAl(TMP)(*i*Bu)₃ in a complicated equilibrium mixture of five compounds.^{7a} Thus a comparison of the ¹H NMR spectra (Figure 1) of an equimolar mixture of LiTMP and GaR₃ and those of the individual components reveals that the two TMT reagents remain separate in benzene-*d*₆ solution. The lack of co-complexation is best deduced by the informative singlet at $\delta = 0.13$ which corresponds to coincidentally overlapping CH₂ and CH₃ resonances of the CH₂-SiMe₃ group on gallium.¹⁰ Furthermore, resonances for both tetrameric and trimeric forms of LiTMP are present and identical to those previously reported.¹¹



To further understand the ability (or lack thereof) to co-complexation we elected to perform some DFT studies, in order to determine energetics of such a process, forming a hypothetical lithium gallate **I**. In the optimised geometry of **I** the metals are connected by a TMP bridge and an alkyl bridge with another two monosilyl groups terminally bonded to the gallium atom. GaR₃ was modelled as a monomer according to its known structure in the solid state,¹² while LiTMP was modelled as a monomer, and a trimer and tetramer which are the two known aggregates of this compound in non-polar solvents.¹¹ Co-complexation between monomeric LiTMP is energetically favoured (–17.1 kcal mol^{–1}), which is unsurprising given the high energy of monomeric LiTMP (Table 1). Crucially, co-complexation of both trimeric and tetrameric LiTMP with GaR₃ are energetically disfavoured (+9.0 kcal mol^{–1}), and in line with the finding from the solution studies.



Table 1 Comparison of DFT-Computed Relative Energies of Co-complexation of LiTMP and GaR₃


n	ΔE Co-complexation (kcal mol ⁻¹)
1	-17.1
3	+9.0
4	+9.0

Next we determined to ascertain the effect of Lewis donor ligands on the TMT process. In particular PMDETA (*N,N,N',N''*-pentamethyldiethylenetriamine) has been crucial in facilitating the crystallisation of TMT products,⁸ which is important since structural data of these complexes provide valuable information on the *modus operandi* of metalation. Reaction between PMDETA, LiTMP, and GaR₃ at room temperature in hexane for one hour lead to the isolation and structural characterization of [Li{Me₂NCH₂CH₂N(Me)CH₂CH₂N(Me)CH₂}(TMP)GaR₂] (**1**). The molecular structure of **1** (Figure 2, top) reveals that PMDETA has been metallated at a terminal NCH₃ group. The lithium atom reveals a contacted ion pair arrangement where the two metal centres connect through two anions, namely a TMP bridge and an ambidentate NCH₂ fragment of the metallated PMDETA.


Interestingly, that a TMP anion is retained in the structure might suggest that it is GaR₃ that deprotonated the substrate as the structure incorporates a TMP anion and only two monosilyl groups on gallium. However GaR₃ on its own is not a sufficiently strong base to metallate PMDETA, therefore the α -deprotonation is based on the stepwise cooperation between LiTMP and GaR₃. LiTMP deprotonates the triamine followed by the fast trans-metal-trapping process with GaR₃ yielding a proposed intermediate **A** (Scheme 2) where concomitantly produced TMPH helps to fill the coordination sphere of lithium and is thus in close proximity to GaR₃. The sterically encumbered intermediate and enhanced acidity of coordinated TMPH makes it possible for the otherwise inert gallium alkyl to react affording **1** and Me₄Si.

Although we have no direct evidence for the proposed mechanism, it is supported indirectly by the notion that triorganogallium reagents cannot deprotonate coordinating additives such as PMDETA or TMEDA as C–H bonds adjacent to N centres in tertiary amines are only weakly acidic. In

**Figure 2** Molecular structure of **1** with 30% probability displacement ellipsoids (top); all hydrogen atoms except those on metallated CH₂ group of PMDETA have been omitted for clarity. Molecular structure of **2** with 30% probability displacement ellipsoids (bottom); all hydrogen atoms except those on metallated CH₂ group of TMEDA have been omitted for clarity.**Scheme 2** Proposed reaction sequence for the surprising formation of **1**

addition, it was found that the bulkiness of a reagent such as GaR₃ precludes chelation and instead, it acts as a bridging ligand as observed in the crystal structure of R₃Ga–TMEDA–GaR₃,¹³ Further, to an extent, experimental support for the proposed pathway came from the addition of a similar, but smaller Lewis donor TMEDA to the mixture of LiTMP and GaR₃ from which we were able to isolate crystals of [(TMEDA)Li{Me₂NCH₂CH₂N(Me)CH₂}(GaR₃)] (**2**) (Figure 2, bottom). The molecular structure of **2** has a contacted ion pair structure with three alkyl groups on gallium and no TMP anion incorporation, reminiscent of the proposed intermediate **A**. Here, due to the smaller size of the diamine, N₄-tetracoordinated lithium is capped with two molecules of TMEDA one of which is metallated and the other one is neutral, completing the coordination sphere and avoiding the close

proximity of TMPH and GaR_3 , which would induce alkyl basicity. Confident that LiTMP and GaR_3 indeed remain separate in non-coordinating solvent, yet cooperate in metalation reactions we next tested this mixture as a TMT reagent using anisole as a benchmark molecule in directed *ortho*-metalation. Thus, to a hexane suspension of equimolar amounts of GaR_3 and LiTMP at room temperature, a molar equivalent of anisole was added to give a light yellow suspension. After stirring the mixture for one hour, an equivalent of PMDETA was added and the solution placed at -33°C affording a crop of colourless crystals of $[(\text{PMDETA})\text{Li}(o\text{-C}_6\text{H}_4\text{OMe})\text{GaR}_3]$ (**3**) in 55% isolated yield (Scheme 3).



Scheme 3 Synthesis of $[(\text{PMDETA})\text{Li}(o\text{-C}_6\text{H}_4\text{OMe})\text{Ga}(\text{CH}_2\text{SiMe}_3)_3]$ (**3**)

The structure of **3** (Figure 3) revealed the formation of a mixed-metal lithium gallate with the metal centres connected by an ambidentate *ortho*-metallated anisole fragment giving rise to a contacted ion-pair structure. The *ortho* carbon of the metallated anisole fragment bonds to gallium forming a new Ga–C bond (Ga–C13 2.0501(15) Å). The distorted tetrahedral lithium is fully solvated by the tridentate PMDETA as well as from the oxygen atom of anisole. In these studies we have demonstrated that LiTMP and GaR_3 are a highly efficient combination of base and trap, to selectively metallate useful aromatic molecules, illustrated by the potency of LiTMP as a base and of GaR_3 in being able to stabilise the newly formed organic carbanions. Probing the synthetic utility of **3**, I_2 was added as an electrophilic source in an effort to prepare 2-iodoanisole. Interestingly, only a trace amount of ca. 1% of the quenched product was obtained, presumably reflecting the high stability of the metallated C–Ga bond. Thus, this TMT system is excellent in the stabilisation of incipient carbanions albeit the stability likely inhibits the downstream utility in simple electrophilic quenching studies.

Underscoring the utility of this metal pairing, we next investigated an analogous system using NaTMP in place of LiTMP , rationalising that the larger alkali metal may be better equipped to form a complex with GaR_3 , hence limiting the ability of the system to promote TMT. Firstly a control reaction between NaTMP and anisole in hexane at room temperature, followed by a standard iodine quench in THF afforded only ca. 20% of 2-iodoanisole. A second control reaction of the in situ $\text{GaR}_3/\text{NaTMP}$ mixture with PMDETA reveals metalation of a methyl group of the Lewis donor after structural characterisation. The structure of this product




Figure 3 Molecular structure of **3** with 30% probability displacement ellipsoids; all hydrogen atoms have been omitted for clarity

$[\text{Na}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}(\text{Me})\text{CH}_2\text{CH}_2\text{N}(\text{Me})\text{CH}_2\}(\text{TMP})\text{GaR}_2]$ (**4**) (Figure 4, top) is directly analogous to that of the lithium gallate **1**. Anticipating that since metalation occurs in this instance, then GaR_3 and NaTMP remain as separate entities in solution, GaR_3 was added to NaTMP in a J. Young's NMR tube in C_6D_6 , causing dissolution of the normally insoluble NaTMP . The ^1H NMR spectrum after 15 minutes revealed, in contrast to the case with LiTMP , that the characteristic resonance of GaR_3 was absent, indicating the surprising formation of a co-complex, albeit the resonances are broad and indicative of either a mixture of compounds or a system undergoing a degree of exchange. This reaction mixture was probed further by adding anisole directly and reaction monitoring revealed a small amount of metalation alongside the signals of coordinated anisole (Figure 5). Further monitoring by ^1H NMR details that additional metalation does not occur over an 18 hour window at room temperature. Repeating the reaction in a Schlenk flask at room temperature in hexane followed by addition of TMEDA after two hours stirring, and placing at 4°C , resulted in a crop of crystals that were characterised X-ray diffraction as $[(\text{TMEDA})\text{Na}(o\text{-C}_6\text{H}_4\text{OMe})\text{GaR}_3]$ (**5**, 17%) (Figure 4, bottom), a sodium analogue of **3**. This result is surprising since reaction of the suggested sodium gallate with anisole would likely possess insufficient basicity to promote metalation, comparable with the $\text{LiTMP}/i\text{BuAl}_3$ mixture that forms a lithium aluminate that lacks the intrinsic basicity to promote C–H bond cleavage.^{7a} Thus we attribute the reactivity as the result of unrestricted NaTMP that had not yet formed a complex with GaR_3 , in accordance with the busier ^1H NMR spectrum recorded after 15 minutes. Probing the co-complex formation further, an equimolar mixture of NaTMP and GaR_3 in C_6D_6 was monitored by ^1H NMR until the spectra ceased evolving. At this point (3 days) anisole was added and the resulting spectrum detailed that metalation does not occur, giving further evidence to our original hypothesis and emphasizing the importance of the lithium reagent in preventing co-complexation and therefore subdued reactivity (Figure 5).




Figure 4 Molecular structure of **4** with 30% probability displacement ellipsoids (top); all hydrogen atoms have been omitted for clarity except for those on the metallated CH_2 group of PMDETA. Molecular structure of **5** with 30% probability displacement ellipsoids (bottom); all hydrogen atoms have been omitted for clarity.





Figure 5 Comparative NMR spectra illustrating that as co-complex formation progresses in time the extent of metalation decreases

Lastly we sought to investigate whether the order of addition would prejudice the reaction in favour of the metallated (trapped) product, rationalising that pre-complexation of anisole with Lewis acidic GaR_3 would compromise the moderately slow complexation of GaR_3 with NaTMP and hence result in enhanced yields of metallated products. In this case reaction between GaR_3 and anisole in a J. Young's

NMR tube in C_6D_6 for 15 minutes was followed by addition of NaTMP. At this point trans-metal-trapped anisole [(TMP(H)Na(*o*- $\text{C}_6\text{H}_4\text{OMe}$) GaR_3)] (**6**) was observed in the ^1H NMR spectrum in 74% yield against hexamethylbenzene (C_6Me_6) as internal standard (Scheme 4). Addition of TME-DA to this mixture affords a spectrum reminiscent of **5**.



Scheme 4 The effect of addition order in TMT using $\text{GaR}_3/\text{NaTMP}$ system. GaR_3 and NaTMP form complex limiting TMT to give **5** (left). Pre-reaction of GaR_3 with anisole enhances yield of **6** (right).

Repeating this reaction in a Schlenk flask and conducting an electrophilic quench with iodine resulted in formation of 2-iodoanisole in 65% yield. That sodium gallate **5** affords 2-iodoanisole in reasonable yields after an electrophilic quench is itself surprising since the analogous reaction with **3** only affords trace amounts of products. This clear alkali metal effect, with the sodium gallate exhibiting far superior onward reactivity than the lithium counterpart suggests that when paired with the appropriate secondary metal, gallium–carbon bond functionalisation can be facile.

Finally, while we have demonstrated that the GaR_3 trans-metal-trapping system is highly efficient in stabilising emergent carbanions, post metalation with LiTMP, taking this trans-metal-trapping to the next level, that is, using the complexes in further C–element bond formation, remains to be realised as a routine procedure. The challenge is that since the trap must be a strong Lewis acidic carbophilic metal complex, logically that will produce a strong metal–carbon bond that may not be easily broken by an electrophile. Standard electrophilic quenching strategies using either iodine, or *N*-bromosuccinimide proved to be unsuccessful, resulting in essentially hydrolysed material after reaction and aqueous workup. This is rather unsurprising given the robust nature of the Ga–C bonds formed using the LiTMP GaR_3 system. Thus, we turned attention to C–C bond formation via palladium-catalysed cross-coupling protocols. A reaction between **3**, 4-bromobenzonitrile, and $\text{Pd}(\text{PPh}_3)_4$ in THF at 80 °C for 16 hours was performed. After a standard organic workup, the cross-coupled product 2'-methoxybiphenyl-4-carbonitrile (**7**) was isolated in 73% yield. However the reaction yield was not reproducible (typically yields varied in the range 50–70% on repeated runs), and variable amounts of 4-[(trimethylsilyl)methyl]benzonitrile were obtained, indicating that under these conditions the reaction evidenced poor selectivity. Using a different organic electrophile, benzoyl chloride, under the same conditions resulted in a moderate 62% yield of 2-benzoylanisole (**8**) (Scheme 5).



Scheme 5 Palladium-mediated C–C bond formation from **3**, using 4-bromobenzonitrile or benzoyl chloride, affording 2'-methoxybiphenyl-4-carbonitrile or 2-benzoylanisole, respectively

Rationalising that under the reasonably harsh reaction conditions required to promote cross-coupling, competing reaction of a CH_2SiMe_3 occurs, we decided to perform one more reaction between the homoleptic lithium gallate LiGaR_4 , and 4-iodoacetophenone. Once more the cross-coupled product was formed in a moderate 62% yield against a ferrocene internal standard. Thus while we demonstrate a rare example of gallium-based cross-coupling,^{8b,14} conclusions can be drawn. A major facet of the trans-metal-trapping strategy is the stabilisation of incipient carbanions by forming strong C–Ga bonds. This however, as one would expect, appears to be problematic, from the standpoint of facile onward reactivity. Thus, this case study clearly suggests a way forward for trans-metal-trapping. Finding new pairings of base and trapping agent that do not co-complex in the mild conditions used in reaction is the first goal since the lack of co-complexation promotes higher yielding processes. The choice of trap should also fulfil two requirements: (i) the atom should be able to form strong enough M–C bonds to stabilise the frequently sensitive metalation anions; (ii) the resulting M–C bonds should also be labile enough to promote straightforward reactivity into more complicated and synthetically useful bis-aryl or heteroaryl molecules. Moreover, despite the fact that the $\text{LiTMP}/\text{GaR}_3$ system is a superior base/trap pairing, the $\text{NaTMP}/\text{GaR}_3$ system is more predisposed to favour onward Ga–C functionalisation. Replacing lithium by sodium as the pre-eminent metallating agent would be attractive from a sustainability viewpoint given their comparative earth abundance. Future work will determine whether this alkali metal effect is due primarily to sodium versus lithium or whether the different coordination spheres involved (e.g., TMEDA versus PMDETA) influences the remarkable difference in iodination efficiency. Current work in our laboratories is focussed on meeting these goals by incorporating the crucial elements into new trans-metal-trapping systems.

All reactions and manipulations were conducted under a protective argon atmosphere using either standard Schlenk techniques or an MBraun glove box fitted with a gas purification and recirculation unit. NMR experiments were conducted in J. Young's tubes oven dried and flushed with argon prior to use. Hexane, toluene, and THF were dried by heating to reflux over sodium benzophenone ketyl and then distilled under N_2 prior to use. All other reagents were purchased commercially from Sigma-Aldrich and dried via distillation (where appro-

priate) from the appropriate drying agent prior to use. LiTMP^{11} and $\text{Ga}(\text{CH}_2\text{SiMe}_3)_3^{15}$ were prepared as previously described or by slight variations thereof. NMR Spectroscopy: NMR spectra were recorded on a Bruker AV3 or AV 400 MHz spectrometer operating at 400.13 MHz for ^1H , 376.46 MHz for ^7Li and 100.62 MHz for ^{13}C . All ^{13}C spectra were proton decoupled. ^1H and ^{13}C NMR spectra were referenced against the appropriate solvent signal. ^7Li NMR spectra were referenced against LiCl in D_2O at $\delta = 0.00$. X-ray Crystallography: Crystallographic data were collected on Oxford Diffraction instruments with Mo $\text{K}\alpha$ radiation ($\lambda = 0.71073 \text{ \AA}$) or Cu $\text{K}\alpha$ radiation ($\lambda = 1.54184 \text{ \AA}$). Structures were solved using SHELXS^{16a} or OLEX2,^{16b} while refinement was carried out on F^2 against all independent reflections by the full matrix least-squares method using the SHELXL programs or by the Gauss-Newton algorithm using OLEX2. All non-hydrogen atoms were refined using anisotropic thermal parameters. Selected crystallographic details and refinement details are provided in Table S1. CCDC 1880970–1880974 contains the supplementary crystallographic data for these structures. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/getstructures. DFT computational studies¹⁷ employing the B3LYP method^{18,19} and the 6-311G(d,p) basis set.²⁰

[(PMDETA)Li(o-C₆H₄OMe)Ga(CH₂SiMe₃)₃] (**3**)

To a suspension of LiTMP (0.074 g, 0.5 mmol) and $\text{Ga}(\text{CH}_2\text{SiMe}_3)_3$ (0.165 g, 0.5 mmol) in hexane (10 mL), an equivalent of anisole (0.054 g, 54 μL , 0.5 mmol) was added at r.t. As soon as anisole was added, a yellow fine suspension was formed which persisted during stirring for 1 hour at r.t. PMDETA (0.11 mL, 0.5 mmol) was added, the solvent was exchanged in vacuo for toluene and the yellow solution placed in freezer to obtain **3** (0.17 g, 55%) as colourless crystals suitable for X-ray diffraction analysis.

^1H NMR (400.13 MHz, 298 K, $\text{THF}-d_6$): $\delta = 7.44$ (d, 1 H, Ar-H), 6.78 (t, 1 H, Ar-H), 6.53 (t, 1 H, *p*-CH), 6.40 (d, 1 H, Ar-H), 6.40 (d, 1 H, Ar-H), 3.59 (s, 3 H, OCH_3), 2.47 (m, 4 H, $\text{NCH}_2\text{CH}_2\text{N}$), 2.37 (m, 4 H, $\text{NCH}_2\text{CH}_2\text{N}$), 2.26 (s, 3 H, NCH_3), 2.20 [s, 12 H, $\text{N}(\text{CH}_3)_2$], -0.19 [s, 27 H, $\text{Si}(\text{CH}_3)_3$], -0.85 (s, 6 H, CH_2SiMe_3).

^{13}C NMR (100.62 MHz, 298 K, $\text{THF}-d_6$): $\delta = 166.7$ (Ar-C), 155.2 (C-Ga), 138.8 (Ar-C), 125.0 (Ar-C), 120.9 (Ar-C), 119.4 (Ar-C), 107.5 (Ar-C), 58.5 (PMDETA), 56.2 (PMDETA), 54.3 (OCH_3), 46.0 (PMDETA), 43.7 (PMDETA), 3.9 [$\text{Si}(\text{CH}_3)_3$], 0.5 (CH_2SiMe_3).

^7Li NMR (376.46 MHz, 298 K, $\text{THF}-d_6$): $\delta = 0.14$.

Anal. Calcd for $\text{C}_{28}\text{H}_{63}\text{GaLiN}_3\text{OSi}_3$: C, 54.35; H, 10.26; N, 6.79. Found: C, 54.95; H, 9.94; N, 7.55.

[(TMEDA)Na(o-C₆H₄OMe)Ga(CH₂SiMe₃)₃] (**5**)

In a Schlenk flask, $\text{Ga}(\text{CH}_2\text{SiMe}_3)_3$ (0.33 g, 1 mmol) and NaTMP (0.163 g, 1 mmol) were suspended in hexane and stirred at r.t. for 1 h. To this mixture, one equivalent of anisole (0.11 mL, 1 mmol) was added, retaining the white suspension, and stirred at r.t. for a further 2 h. Upon

addition of TMEDA (0.21 mL, 1 mmol), a colourless solution was obtained which, with slow cooling to 4 °C, produced a crop of colourless crystals of **5** (98 mg, 17%).

¹H NMR (400.13 MHz, 298 K, C₆D₆): δ = 8.00 (dd, 1 H, Ar-*H*), 7.09 (td, 1 H, Ar-*H*), 7.04 (td, 1 H, Ar-*H*), 6.56 (dd, 1 H, Ar-*H*), 6.40, 3.46 (s, 3 H, OCH₃), 1.59 [s, 12 H, N(CH₃)₂], 1.54 (s, 4 H, NCH₂CH₂N), 0.33 [s, 27 H, Si(CH₃)₃], -0.51 (s, 6 H, CH₂SiMe₃).

¹³C NMR (100.62 MHz, 298 K, C₆D₆): δ = 165.1 (Ar-C), 149.7 (C-Ga), 141.4 (Ar-C), 127.9 (Ar-C), 123.0 (Ar-C), 112.2 (Ar-C), 56.9 (OCH₃), 56.5 (TMEDA), 45.1 (TMEDA), 3.6 [Si(CH₃)₃], -0.7 (CH₂SiMe₃).

2'-Methoxybiphenyl-4-carbonitrile (**7**)

To a solution of **3** (200 mg, 0.323 mmol) in THF (8 mL) were added 4-bromobenzonitrile (59 mg, 0.323 mmol) and Pd(PPh₃)₄ (18 mg 5 mol%). The mixture was then stirred at reflux temperature for 16 h. After removal of all volatiles in vacuo, the residue was extracted with Et₂O (20 mL), then washed with H₂O (2 × 10 mL) and brine (2 × 10 mL). The organic phase was then dried (MgSO₄) and concentrated. Column chromatography (silica gel, hexane/EtOAc 95:5–90:10) afforded **7** (49 mg, 0.24 mmol, 73%) as a colourless oil; *R*_f = 0.45. ¹H and ¹³C NMR spectra are in agreement with those previously published.²¹

2-Benzoylanisole (**8**)

To a solution of **3** (200 mg, 0.323 mmol) in THF (8 mL) were added Pd(PPh₃)₄ (18 mg, 5 mol%) followed by benzoyl chloride (0.037 mL, 0.323 mmol) via syringe. The mixture was then stirred at reflux temperature for 16 h. After removal of all volatiles in vacuo, the residue was extracted with Et₂O (20 mL), then washed with H₂O (2 × 10 mL) and brine (2 × 10 mL). The organic phase was then dried (MgSO₄) and concentrated. Column chromatography (silica gel, hexane/EtOAc 95:5–90:10) afforded **8** (43 mg, 0.20 mmol, 62%) as a colourless solid; *R*_f = 0.35. ¹H and ¹³C NMR spectra are in agreement with those previously published.²²

NMR Study: Synthesis of [(TMP(H)Na(o-C₆H₄OMe)GaR₃)] (**6**)

In a J. Young's NMR tube, Ga(CH₂SiMe₃)₃ (83 mg, 0.25 mmol) and anisole (27 μL, 0.25 mmol) were combined together in C₆D₆ solvent affording a colourless solution. ¹H NMR analysis confirmed no reactivity had occurred between these two species. To this mixture, NaTMP (41 mg, 0.25 mmol) was added and upon dissolution gave a colourless solution. Analysis by ¹H NMR spectroscopy indicated metalation of anisole had occurred in 74% yield (using C₆Me₆ as an internal standard) to give **6**. [Note: TMP(H) resonances are lower frequency by ~0.3 ppm than 'free' TMP(H); we propose this is due to coordination of TMP(H) to Na in the absence of any other donor molecules.] Finally, addition of TMEDA (38 μL, 0.25 mmol) resulted in its coordination to the sodium centre and release of TMP(H) (confirmed by ¹H NMR).

[(TMP(H)Na(o-C₆H₄OMe)GaR₃)] (**6**)

¹H NMR (400.13 MHz, 298 K, C₆D₆): δ = 7.97 (dd, 1 H, Ar-*H*), 7.09 (m, 1 H, Ar-*H*), 7.02 (td, 1 H, *p*-CH), 6.61 (d, 1 H, Ar-*H*), 3.46 (s, 3 H, OCH₃), 2.08 (s, C₆Me₆), 1.28 [m, 2 H, γ-TMP(H)], 0.92 [t, 4 H, β-TMP(H)], 0.70 [s, 12 H, CH₃ of TMP(H)], 0.26 [s, 28 H, Si(CH₃)₃ + TMP(H)], -0.57 (s, 6 H, CH₂SiMe₃).

Post-TMEDA Addition:

¹H NMR (400.13 MHz, 298 K, C₆D₆): δ = 7.90 (dd, 1 H, Ar-*H*), 7.06 (m, 1 H, Ar-*H*), 6.98 (td, 1 H, *p*-CH), 6.58 (d, 1 H, Ar-*H*), 3.48 (s, 3 H, OCH₃), 2.08 (s, C₆Me₆), 1.65 (m, 16 H, TMEDA), 1.53 [m, 2 H, γ-TMP(H)], 1.24

[t, 4 H, β-TMP(H)], 1.05 [s, 12 H, CH₃ of TMP(H)], 0.25 [s, 28 H, Si(CH₃)₃ + TMP(H)], -0.59 (s, 6 H, CH₂SiMe₃).

NMR Study: Synthesis of [(PMDETA)Li(o-C₆H₄OMe)Ga(CH₂SiMe₃)₃] (**3**)

In a J. Young's NMR tube, equimolar amounts of Ga(CH₂SiMe₃)₃ (83 mg, 0.25 mmol) and LiTMP (36 mg, 0.25 mmol) were combined in C₆D₆ (0.5 mL) resulting in a white suspension. Addition of anisole (27 μL, 0.25 mmol) was then performed and allowed to react for 1 h at r.t. with a white suspension persisting. Dissolution was achieved by the addition of PMDETA (53 μL, 0.25 mmol) to afford a yellow solution; compound **3** was obtained in 75% yield (¹H NMR with C₆Me₆ as internal standard).

¹H NMR (400.13 MHz, 298 K, C₆D₆): δ = 8.19 (d, 1 H, Ar-*H*), 7.06 (t, 1 H, Ar-*H*), 7.02 (dt, 1 H, *p*-CH), 6.31 (d, 1 H, Ar-*H*), 3.44 (s, 3 H, OCH₃), 2.03 (s, 3 H, NCH₃), 1.83 (m, 4 H, NCH₂CH₂N), 1.75 (m, 4 H, NCH₂CH₂N), 1.57 [s, 12 H, N(CH₃)₂], 0.32 [s, 27 H, Si(CH₃)₃], -0.34 (s, 6 H, CH₂SiMe₃).

¹³C NMR (100.62 MHz, 298 K, C₆D₆): δ = 164.6 (Ar-C), 142.6 (Ar-C), 131.7 (C-Ga), 124.3 (Ar-C), 125.9 (Ar-C), 116.5 (Ar-C), 62.7 (OCH₃), 57.2 (PMDETA), 53.5 (PMDETA), 45.6 (PMDETA), 44.5 (PMDETA), 4.1 [Si(CH₃)₃], 1.5 (CH₂SiMe₃).

⁷Li NMR (376.46 MHz, 298 K, C₆D₆): δ = 0.10.

Metalation of Anisole followed by Electrophilic Quenching Using I₂: NaTMP + Anisole

In a Schlenk flask, NaTMP (0.163 g, 1 mmol) was suspended in hexane (10 mL) at r.t. To this, anisole (0.11 mL, 1 mmol) was added affording a white suspension, and the mixture was stirred at r.t. for 2 h. In a separate Schlenk tube, a solution of I₂ (6 mmol) was prepared in THF (10 mL). Both Schlenk tubes were then cooled to -78 °C in an acetone/dry ice bath and stirred for 20 min until completely cooled. Then, the I₂/THF solution was syphoned into the mixture at -78 °C and allowed to slowly warm up to r.t. over 16 h. Workup of the mixture was achieved by addition of sat. aq Na₂S₂O₃ solution until bleaching occurred (approx. 20 mL), followed by addition of sat. aq NH₄Cl (10 mL). The mixture was then extracted with EtOAc (3 × 10 mL) and the combined organic layers were washed with brine (10 mL), dried (MgSO₄), and filtered. All organic solvents were then removed under reduced pressure to give a brown solid; 2-iodoanisole was obtained in 20% yield (NMR using C₆Me₆ a internal standard in CDCl₃). Spectroscopic values obtained are in good agreement with those previously reported in the literature.²³

Metalation of Anisole followed by Electrophilic Quenching Using I₂: LiTMP (or NaTMP) + GaR₃ + Anisole

In a Schlenk flask, LiTMP (0.146 g, 1 mmol) and Ga(CH₂SiMe₃)₃ (0.33 g, 1 mmol) were suspended in hexane (10 mL) at r.t. To this, anisole (0.11 mL, 1 mmol) was added retaining a white suspension, and the mixture was stirred at r.t. for 2 h [for NaTMP (0.163 g, 1 mmol) reaction, the order of addition was reversed so that NaTMP was added last]. In a separate Schlenk tube, a solution of I₂ (6 mmol) was prepared in THF (10 mL). Both flasks were then cooled to -78 °C in an acetone/dry ice bath and stirred for 20 min until completely cooled. Then, the I₂/THF solution was syphoned into the mixture at -78 °C and allowed to slowly warm up to r.t. overnight (approx. 16 h in total). Workup of the mixture was achieved by addition of sat. aq Na₂S₂O₃ solution until bleaching occurred (approx. 20 mL), followed by addition of sat. aq NH₄Cl (10 mL). The mixture was then extracted with EtOAc (3 × 10 mL) and the combined organic layers were washed with brine (10 mL), dried (MgSO₄), and filtered. All organic solvents

were then removed under reduced pressure to give a brown solid. The yield of 2-iodoanisole was 65% (using NaTMP) and 1.5% (LiTMP) (NMR using C₆Me₆ as internal standard in CDCl₃). NMR data are in agreement with those previously published for 2-iodoanisole.²³

Funding Information

We thank the European Research Council (ERC StG, MixMetApps) and the EPSRC (EP/N011384/1) for their generous sponsorship of this research.

Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/s-0037-1611646>.


References

- (1) (a) Clayden, J. *In Organolithiums: Selectivity for Synthesis* 2002. (b) Snieckus, V. *Chem. Rev.* **1990**, *90*, 879. (c) Mongin, F.; Schlosser, M. *Tetrahedron Lett.* **1996**, *37*, 6551. (d) Schlosser, M. *Angew. Chem. Int. Ed.* **2005**, *44*, 376. (e) Werner, V.; Klatt, T.; Fujii, M.; Markiewicz, J.; Apeloig, Y.; Knochel, P. *Chem. Eur. J.* **2014**, *20*, 8338. (f) Mulvey, R. E.; Robertson, S. D. *Angew. Chem. Int. Ed.* **2013**, *52*, 11470.
- (2) Becker, M. R.; Knochel, P. *Angew. Chem. Int. Ed.* **2015**, *54*, 12501.
- (3) (a) Schlosser, M. *J. Organomet. Chem.* **1967**, *8*, 9. (b) Lochmann, L.; Pospisil, J.; Lim, D. *Tetrahedron Lett.* **1966**, *7*, 257.
- (4) (a) Wunderlich, S. H.; Rohbogner, C. J.; Unsinn, A.; Knochel, P. *Org. Process Res. Dev.* **2010**, *14*, 339. (b) Haag, B.; Mosrin, M.; Ila, H.; Malakhov, V.; Knochel, P. *Angew. Chem. Int. Ed.* **2011**, *50*, 9794.
- (5) (a) Uchiyama, M.; Kameda, M.; Mishima, O.; Yokoyama, N.; Koike, M.; Kondo, Y.; Sakamoto, T. *J. Am. Chem. Soc.* **1998**, *120*, 4934. (b) Kondo, Y.; Shilai, M.; Uchiyama, M.; Sakamoto, T. *J. Am. Chem. Soc.* **1999**, *121*, 3539. (c) Snégaroff, K.; Komagawa, S.; Chevallier, F.; Gros, P. C.; Golhen, S.; Roisnel, T.; Uchiyama, M.; Mongin, F. *Chem. Eur. J.* **2010**, *16*, 8191. (d) Akimoto, G.; Otsuka, M.; Takita, R.; Uchiyama, M.; Hedidi, M.; Bentabed-Ababsa, G.; Lassagne, F.; Erb, W.; Mogin, F. *J. Org. Chem.* **2018**, *83*, 13498.
- (6) (a) Mulvey, R. E.; Mongin, F.; Uchiyama, M.; Kondo, Y. *Angew. Chem. Int. Ed.* **2007**, *46*, 3802. (b) Andrikopoulos, P. C.; Armstrong, D. R.; Graham, D. V.; Hevia, E.; Kennedy, A. R.; Mulvey, R. E.; O'Hara, C. T.; Talmard, C. *Angew. Chem. Int. Ed.* **2005**, *44*, 3459. (c) Armstrong, D. R.; Clegg, W.; Dale, S. H.; Graham, D. V.; Hevia, E.; Hogg, L. M.; Honeyman, G. W.; Kennedy, A. R.; Mulvey, R. E. *Chem. Commun.* **2007**, 598. (d) Clegg, W.; Conway, B.; Graham, D. V.; Hevia, E.; Kennedy, A. R.; Mulvey, R. E.; Russo, L.; Wright, D. S. *Chem. Eur. J.* **2009**, *15*, 7074. (e) Armstrong, D. R.; Blair, V. L.; Clegg, W.; Dale, S. H.; Garcia-Alvarez, J.; Honeyman, G. W.; Hevia, E.; Mulvey, R. E.; Russo, L. *J. Am. Chem. Soc.* **2010**, *132*, 9480. (f) Wunderlich, S. H.; Kienle, M.; Knochel, P. *Angew. Chem. Int. Ed.* **2009**, *48*, 7256. (g) Garcia-Álvarez, J.; Kennedy, A. R.; Klett, J.; Mulvey, R. E. *Angew. Chem. Int. Ed.* **2007**, *46*, 1105. (h) Blair, V. L.; Clegg, W.; Conway, B.; Hevia, E.; Kennedy, A. R.; Klett, J.; Mulvey, R. E.; Russo, L. *Chem. Eur. J.* **2008**, *14*, 65. (i) Wunderlich, S. H.; Knochel, P. *Angew. Chem. Int. Ed.* **2009**, *48*, 9717. (j) Alborés, P.; Carrella, L. M.; Clegg, W.; García-Álvarez, P.; Kennedy, A. R.; Klett, J.; Mulvey, R. E.; Rentschler, E.; Russo, L. *Angew. Chem. Int. Ed.* **2009**, *48*, 3317. (k) Nagaradja, E.; Chevallier, F.; Roisnel, T.; Jouikov, V.; Mongin, F. *Tetrahedron* **2012**, *68*, 3063. (l) Bedford, R. B.; Brenner, P. B.; Carter, E.; Cogswell, P. M.; Haddow, M. F.; Harvey, J. N.; Murphy, D. M.; Nunn, J.; Woodall, C. H. *Angew. Chem. Int. Ed.* **2014**, *53*, 1804. (m) Martínez-Martínez, A. J.; Kennedy, A. R.; Mulvey, R. E.; O'Hara, C. T. *Science (Washington, D. C.)* **2014**, *346*, 834.
- (7) (a) Armstrong, D. R.; Crosbie, E.; Hevia, E.; Mulvey, R. E.; Ramsay, D. L.; Robertson, S. D. *Chem. Sci.* **2014**, *5*, 3031. (b) Uzelac, M.; Kennedy, A. R.; Hevia, E. *Inorg. Chem.* **2017**, *56*, 8615. (c) Uzelac, M.; Mulvey, R. E. *Chem. Eur. J.* **2018**, *24*, 7786.
- (8) (a) Uzelac, M.; Kennedy, A. R.; Hevia, E.; Mulvey, R. E. *Angew. Chem. Int. Ed.* **2016**, *55*, 13147. (b) McLellan, R.; Uzelac, M.; Kennedy, A. R.; Hevia, E.; Mulvey, R. E. *Angew. Chem. Int. Ed.* **2017**, *56*, 9566.
- (9) Frischmuth, A.; Fernández, M.; Barl, N. M.; Achraimer, F.; Zipse, H.; Berionni, G.; Mayr, H.; Karaghiosoff, K.; Knochel, P. *Angew. Chem. Int. Ed.* **2014**, *53*, 7928.
- (10) Armstrong, D. R.; Brammer, E.; Cadenbach, T.; Hevia, E.; Kennedy, A. R. *Organometallics* **2013**, *32*, 480.
- (11) Hevia, E.; Kennedy, A. R.; Mulvey, R. E.; Ramsay, D. L.; Robertson, S. D. *Chem. Eur. J.* **2013**, *19*, 14069.
- (12) Kramer, M. U.; Robert, D.; Nakajima, Y.; Englert, U.; Spaniol, T. P.; Okuda, J. *Eur. J. Inorg. Chem.* **2007**, 665.
- (13) Hallock, R. B.; Hunter, W. E.; Atwood, J. L.; Beachley, O. T. Jr. *Organometallics* **1985**, *4*, 547.
- (14) (a) Han, Y.; Fang, L.; Tao, W.-T.; Huang, Y.-Z. *Tetrahedron Lett.* **1995**, *36*, 1287. (b) Blum, J.; Gelman, D.; Baidossi, W.; Shakh, E.; Rosenfeld, A.; Aizenshtat, Z.; Wassermann, B. C.; Frick, M.; Heymer, B.; Schutte, S.; Wernik, S.; Schumann, H. *J. Org. Chem.* **1997**, *62*, 8681. (c) Gelman, D.; Schumann, H.; Blum, J. *Tetrahedron Lett.* **2000**, *41*, 7555. (d) Mikami, S.; Yorimitsu, H.; Oshima, K. *Synlett* **2002**, 1137.
- (15) Dennis, L. M.; Patnode, W. *J. Am. Chem. Soc.* **1932**, *54*, 182.
- (16) (a) Sheldrick, G. M. *Acta Crystallogr., Sect. C* **2015**, *71*, 3. (b) Dolomanov, O. V.; Bourhis, L. J.; Gildea, R. J.; Howard, J. A. K.; Puschmann, H. *J. Appl. Crystallogr.* **2009**, *42*, 339.
- (17) Kohn, W.; Becke, A. D.; Parr, R. G. *J. Phys. Chem.* **1996**, *100*, 12974.
- (18) Becke, A. D. *Phys. Rev. A* **1988**, *38*, 3098.
- (19) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785.
- (20) McLean, A. D.; Chandler, G. S. *J. Chem. Phys.* **1980**, *72*, 5639.
- (21) Alam, N.; Amatore, C.; Combellas, C.; Pinson, J.; Savéant, J.-M.; Thiébaud, A.; Verpeaux, J.-N. *J. Org. Chem.* **1988**, *53*, 1496.
- (22) Jafarpour, F.; Rashidi-Ranjbar, P.; Kashani, A. O. *Eur. J. Org. Chem.* **2011**, 2128.
- (23) Kulbitski, K.; Nisnevich, G.; Gandelman, M. *Adv. Synth. Catal.* **2011**, *353*, 1438.

Supporting Information
for DOI: 10.1055/s-0037-1611646
© Georg Thieme Verlag KG Stuttgart · New York 2019

Table S1 Crystallographic parameters for compounds 1 - 5

	1	2	3	4	5
Empirical formula	C ₂₆ H ₆₂ Ga ₁ Li ₁ N ₄ Si ₂	C ₂₄ H ₆₄ Ga ₁ Li ₁ N ₄ Si ₃	C ₂₈ H ₆₃ Ga ₁ Li ₁ N ₃ O ₁ Si ₃	C ₂₆ H ₆₂ Ga ₁ N ₄ Na ₁ Si ₂	C ₂₅ H ₅₆ Ga ₁ N ₂ Na ₁ O ₁ Si ₃
Mol. Mass	563.63	569.72	618.74	579.68	577.69
Crystal system	orthorhombic	monoclinic	monoclinic	orthorhombic	triclinic
a/ Å	18.6987(10)	10.4261(3)	11.7056(6)	18.5579(11)	10.0679(4)
b/ Å	10.8705(5)	23.2204(7)	18.4572(8)	11.1293(5)	11.0101(4)
c/ Å	16.4967(9)	14.7638(4)	18.2034(10)	16.5363(10)	15.6691(6)
α / °	90	90	90	90	83.626(3)
β / °	90	94.048(3)	107.720(6)	90	88.933(3)
γ / °	90	90	90	90	83.389(3)
V/ Å ³	3353.2(3)	3565.37(18)	3746.3(3)	3415.3(3)	1714.65(11)
Z	4	4	4	4	2
λ / Å	0.71073	0.71073	0.71073	1.54184	1.54184
Measured reflections	24436	36068	47025	16366	11249
Unique reflections	8796	9726	10393	5258	6837
R_{int}	0.0448	0.0541	0.0525	0.0497	0.0380
Observed rflns [$I > 2\sigma(I)$]	7573	6921	8291	4675	6270
Goof	1.111	1.034	1.049	1.071	1.065
R [on F , obs rflns only]	0.0488	0.0438	0.0355	0.0473	0.0751
ωR [on F^2 , all data]	0.1316	0.0934	0.0837	0.1199	0.2183
Largest diff. Peak/hole. e/ Å ⁻³	0.795 / -0.571	0.591 / -0.300	0.415 / -0.310	1.006 / -0.311	1.720 / -1.239



NMR Characterisation of **3**




Figure S1 ^1H NMR in d_8 -THF




Figure S2 ^{13}C NMR in $\text{d}_8\text{-THF}$




Figure S3 ^7Li NMR in $\text{d}_8\text{-THF}$

Reaction monitoring of formation of **3** from reaction between LiTMP, GaR₃, Anisole then PMDETA. **3** forms in 74% against hexamethylbenzene as an internal standard.




Figure S4 ¹H NMR spectrum in C₆D₆




Figure S5 ⁷Li NMR spectrum in C₆D₆





Figure S6 ^{13}C NMR in C_6D_6

Iodine quench of **3**



Labelled peaks correspond to the low yield of 4 iodo anisole *ca.* 1% - Calculated against hexamethylbenzene internal standard (also labelled).




Figure S7 1H NMR in $CDCl_3$

NaTMP + anisole - Iodine quench – This reaction results in only 20% of 2-I-anisole.






Figure S8 ¹H NMR spectrum in CDCl₃



NMR Characterisation of isolated crystals of **5** – isolated in 17% yield




Figure S9 ^1H NMR spectrum in C_6D_6





Figure S10 ^{13}C NMR spectrum in C_6D_6



Changing the order of addition – Characterisation of **6** – conversion to 74% against hexamethylbenzene as an internal standard

GaR_3 + anisole then NaTMP





Figure S11 ^1H NMR spectrum in C_6D_6



GaR_3 + anisole then NaTMP followed by an iodine quench, affording 2-iodo-anisole in 63% against hexamethylbenzene as an internal standard





Figure S12 ^1H NMR spectrum in CDCl_3



NMR characterisation of **7**




Figure S13 ^1H NMR spectrum in CDCl_3





Figure S14 ^{13}C NMR spectrum in CDCl_3



NMR characterisation of **8**




Figure S15 ^1H NMR spectrum in CDCl_3




Figure S16 ^{13}C NMR spectrum in CDCl_3

checkCIF (basic structural check) running

Checking for embedded fcf data in CIF ...

Found embedded fcf data in CIF. Extracting fcf data from uploaded CIF, please wait
.....

checkCIF/PLATON (basic structural check)

Structure factors have been supplied for datablock(s) lb-11-257c, mu547b, mu687c, mu826c, rmlb1

THIS REPORT IS FOR GUIDANCE ONLY. IF USED AS PART OF A REVIEW PROCEDURE FOR PUBLICATION, IT SHOULD NOT REPLACE THE EXPERTISE OF AN EXPERIENCED CRYSTALLOGRAPHIC REFEREE.

No syntax errors found.

Please wait while processing

[Structure factor report](#)

[CIF dictionary](#)

[Interpreting this report](#)

Datablock: mu826c

Bond precision: C-C = 0.0086 A Wavelength=0.71073

Cell: a=18.6987(10) b=10.8705(5) c=16.4967(9)

alpha=90 beta=90 gamma=90

Temperature: 130 K

	Calculated	Reported
Volume	3353.2(3)	3353.2(3)
Space group	P n a 21	P n a 21
Hall group	P 2c -2n	P 2c -2n
Moiety formula	C26 H62 Ga Li N4 Si2	C26 H62 Ga Li N4 Si2
Sum formula	C26 H62 Ga Li N4 Si2	C26 H62 Ga Li N4 Si2
Mr	563.64	563.63
Dx, g cm ⁻³	1.117	1.116
Z	4	4
Mu (mm ⁻¹)	0.911	0.911
F000	1232.0	1232.0
F000'	1233.74	
h, k, lmax	26, 15, 23	26, 15, 22
Nref	9795[5056]	8796
Tmin, Tmax	0.804, 0.913	0.927, 1.000
Tmin'	0.695	

Correction method= # Reported T Limits: Tmin=0.927

Tmax=1.000 AbsCorr = MULTI-SCAN

Data completeness= 1.74/0.90 Theta(max)= 29.997

R(reflections)= 0.0488(7573) wR2(reflections)= 0.1316(8796)

S = 1.111

Npar= 322

The following ALERTS were generated. Each ALERT has the format

test-name_ALERT_alert-type_alert-level.

Click on the hyperlinks for more details of the test.

●Alert level B

PLAT934_ALERT_3_B Number of (Iobs-Icalc)/SigmaW > 10 Outliers 2 Check

●Alert level C

PLAT220_ALERT_2_C Non-Solvent Resd 1 C Ueq(max)/Ueq(min) Range 3.2 Ratio

PLAT241_ALERT_2_C High 'MainMol' Ueq as Compared to Neighbors of C15 Check

PLAT242_ALERT_2_C Low 'MainMol' Ueq as Compared to Neighbors of N4 Check

PLAT341_ALERT_3_C Low Bond Precision on C-C Bonds 0.0086 Ang.

PLAT412_ALERT_2_C Short Intra XH3 .. XHn H1A ..H23C . 1.84 Ang.

x,y,z = 1_555 Check

PLAT910_ALERT_3_C Missing # of FCF Reflection(s) Below Theta(Min). 7 Note

PLAT911_ALERT_3_C Missing FCF Refl Between Thmin & STh/L= 0.600 2 Report

PLAT915_ALERT_3_C No Flack x Check Done: Low Friedel Pair Coverage 86 %

PLAT977_ALERT_2_C Check Negative Difference Density on H14A -0.50 eA-3

PLAT978_ALERT_2_C Number C-C Bonds with Positive Residual Density. 0 Info

●Alert level G

PLAT343_ALERT_2_G Unusual sp3 Angle Range in Main Residue for C1 Check

PLAT792_ALERT_1_G Model has Chirality at N2 (Polar SPGR) R Verify

PLAT792_ALERT_1_G Model has Chirality at N3 (Polar SPGR) S Verify

PLAT912_ALERT_4_G Missing # of FCF Reflections Above STh/L= 0.600 310 Note

PLAT913_ALERT_3_G Missing # of Very Strong Reflections in FCF 1 Note

PLAT933_ALERT_2_G Number of OMIT Records in Embedded .res File ... 3 Note

0 **ALERT level A** = Most likely a serious problem - resolve or explain

1 **ALERT level B** = A potentially serious problem, consider carefully

10 **ALERT level C** = Check. Ensure it is not caused by an omission or oversight

6 **ALERT level G** = General information/check it is not something unexpected

2 ALERT type 1 CIF construction/syntax error, inconsistent or missing data

8 ALERT type 2 Indicator that the structure model may be wrong or deficient

6 ALERT type 3 Indicator that the structure quality may be low

1 ALERT type 4 Improvement, methodology, query or suggestion

0 ALERT type 5 Informative message, check

Datablock: mu547b

Bond precision: C-C = 0.0035 A Wavelength=0.71073

Cell: a=10.4261(3) b=23.2204(7) c=14.7638(4)

alpha=90 beta=94.048(3) gamma=90

Temperature: 123 K

	Calculated	Reported
Volume	3565.37(18)	3565.37(18)
Space group	P 21/c	P 21/c
Hall group	-P 2ybc	-P 2ybc
Moiety formula	C24 H64 Ga Li N4 Si3	C24 H64 Ga Li N4 Si3
Sum formula	C24 H64 Ga Li N4 Si3	C24 H64 Ga Li N4 Si3
Mr	569.72	569.72

Dx, g cm ⁻³	1.061	1.061
Z	4	4
Mu (mm ⁻¹)	0.889	0.889
F000	1248.0	1248.0
F000'	1250.04	
h, k, lmax	14, 32, 20	14, 30, 20
Nref	10382	9726
Tmin, Tmax	0.659, 0.915	0.782, 1.000
Tmin'	0.581	

Correction method= # Reported T Limits: Tmin=0.782
Tmax=1.000 AbsCorr = MULTI-SCAN

Data completeness= 0.937 Theta(max)= 29.977

R(reflections)= 0.0438(6921) wR2(reflections)= 0.0934(9726)

S = 1.034 Npar= 314

The following ALERTS were generated. Each ALERT has the format

test-name_ALERT_alert-type_alert-level.

Click on the hyperlinks for more details of the test.

●Alert level B

[PLAT910_ALERT_3_B](#) Missing # of FCF Reflection(s) Below Theta(Min). 11 Note

●Alert level C

[PLAT220_ALERT_2_C](#) Non-Solvent Resd 1 C Ueq(max)/Ueq(min) Range 4.5 Ratio

[PLAT222_ALERT_3_C](#) Non-Solv. Resd 1 H Uiso(max)/Uiso(min) Range 5.7 Ratio

[PLAT906_ALERT_3_C](#) Large K Value in the Analysis of Variance 3.315 Check

●Alert level G

[PLAT343_ALERT_2_G](#) Unusual sp3 Angle Range in Main Residue for C17 Check

[PLAT793_ALERT_4_G](#) Model has Chirality at N1 (Centro SPGR) R Verify

[PLAT912_ALERT_4_G](#) Missing # of FCF Reflections Above STh/L= 0.600 580 Note

[PLAT913_ALERT_3_G](#) Missing # of Very Strong Reflections in FCF 1 Note

[PLAT951_ALERT_5_G](#) Calculated (ThMax) and CIF-Reported Kmax Differ 2 Units

[PLAT957_ALERT_1_G](#) Calculated (ThMax) and Actual (FCF) Kmax Differ 2 Units

[PLAT978_ALERT_2_G](#) Number C-C Bonds with Positive Residual Density. 1 Info

0 **ALERT level A** = Most likely a serious problem - resolve or explain

1 **ALERT level B** = A potentially serious problem, consider carefully

3 **ALERT level C** = Check. Ensure it is not caused by an omission or oversight

7 **ALERT level G** = General information/check it is not something unexpected

1 ALERT type 1 CIF construction/syntax error, inconsistent or missing data

3 ALERT type 2 Indicator that the structure model may be wrong or deficient

4 ALERT type 3 Indicator that the structure quality may be low

2 ALERT type 4 Improvement, methodology, query or suggestion

1 ALERT type 5 Informative message, check

Datablock: mu687c

Bond precision: C-C = 0.0025 A Wavelength=0.71073

Cell: a=11.7056(6) b=18.4572(8) c=18.2034(10)

 alpha=90 beta=107.720(6) gamma=90

Temperature: 123 K

	Calculated	Reported
Volume	3746.3(3)	3746.3(3)
Space group	P 21/n	P 21/n
Hall group	-P 2yn	-P 2yn
Moiety formula	C28 H63 Ga Li N3 O Si3	C28 H63 Ga Li N3 O Si3
Sum formula	C28 H63 Ga Li N3 O Si3	C28 H63 Ga Li N3 O Si3
Mr	618.74	618.74
Dx, g cm ⁻³	1.097	1.097
Z	4	4
Mu (mm ⁻¹)	0.853	0.853
F000	1344.0	1344.0
F000'	1346.11	
h, k, lmax	16, 25, 25	16, 25, 25
Nref	10921	10393
Tmin, Tmax	0.843, 0.958	0.861, 1.000
Tmin'	0.843	
Correction method= # Reported T Limits: Tmin=0.861 Tmax=1.000 AbsCorr = MULTI-SCAN		
Data completeness= 0.952 Theta(max)= 29.997		
R(reflections)= 0.0355(8291) wR2(reflections)= 0.0837(10393)		
S = 1.049 Npar= 349		

The following ALERTS were generated. Each ALERT has the format
test-name_ALERT_alert-type_alert-level.
Click on the hyperlinks for more details of the test.

● Alert level B

[PLAT910_ALERT_3_B](#) Missing # of FCF Reflection(s) Below Theta(Min). 13 Note

● Alert level C

[PLAT220_ALERT_2_C](#) Non-Solvent Resd 1 C Ueq(max)/Ueq(min) Range 3.7 Ratio
[PLAT222_ALERT_3_C](#) Non-Solv. Resd 1 H Uiso(max)/Uiso(min) Range 4.2 Ratio

● Alert level G

[PLAT912_ALERT_4_G](#) Missing # of FCF Reflections Above STh/L= 0.600 515 Note
[PLAT978_ALERT_2_G](#) Number C-C Bonds with Positive Residual Density. 4 Info

- 0 **ALERT level A** = Most likely a serious problem - resolve or explain
- 1 **ALERT level B** = A potentially serious problem, consider carefully
- 2 **ALERT level C** = Check. Ensure it is not caused by an omission or oversight
- 2 **ALERT level G** = General information/check it is not something unexpected

- 0 ALERT type 1 CIF construction/syntax error, inconsistent or missing data
- 2 ALERT type 2 Indicator that the structure model may be wrong or deficient
- 2 ALERT type 3 Indicator that the structure quality may be low
- 1 ALERT type 4 Improvement, methodology, query or suggestion
- 0 ALERT type 5 Informative message, check

Datablock: rmlb1

Bond precision: C-C = 0.0092 A Wavelength=1.54184
Cell: a=18.5579(11) b=11.1293(5) c=16.5363(10)

4 ALERT type 2 Indicator that the structure model may be wrong or deficient
4 ALERT type 3 Indicator that the structure quality may be low
3 ALERT type 4 Improvement, methodology, query or suggestion
0 ALERT type 5 Informative message, check

Datablock: Ib-11-257c

Bond precision: C-C = 0.0041 Å Wavelength=1.54184
Cell: a=10.0679(4) b=11.0101(4) c=15.6691(6)
alpha=83.626(3) beta=88.933(3) gamma=83.389(3)
Temperature: 123 K

	Calculated	Reported
Volume	1714.65(11)	1714.65(11)
Space group	P -1	P -1
Hall group	-P 1	-P 1
Moiety formula	C25 H56 Ga N2 Na O Si3	C25 H56 Ga N2 Na O Si3
Sum formula	C25 H56 Ga N2 Na O Si3	C25 H56 Ga N2 Na O Si3
Mr	577.70	577.69
Dx, g cm ⁻³	1.119	1.119
Z	2	2
Mu (mm ⁻¹)	2.353	2.353
F000	624.0	624.0
F000'	624.22	
h, k, lmax	12, 13, 19	0, 0, 0
Nref	6910	6837
Tmin, Tmax	0.305, 0.625	0.232, 1.000
Tmin'	0.195	

Correction method= # Reported T Limits: Tmin=0.232
Tmax=1.000 AbsCorr = MULTI-SCAN
Data completeness= 0.989 Theta(max)= 73.400
R(reflections)= 0.0751(6270) wR2(reflections)= 0.2183(6837)
S = 1.065 Npar= 312

The following ALERTS were generated. Each ALERT has the format
[test-name_ALERT_alert-type_alert-level](#).
Click on the hyperlinks for more details of the test.

● Alert level C

[PLAT242_ALERT_2_C](#) Low 'MainMol' Ueq as Compared to Neighbors of Si2 Check
[PLAT971_ALERT_2_C](#) Check Calcd Resid. Dens. 0.96A From Ga1 1.74 eA-3

And 3 other PLAT971 Alerts

More ...

[PLAT973_ALERT_2_C](#) Check Calcd Positive Resid. Density on Ga1 1.49 eA-3
[PLAT977_ALERT_2_C](#) Check Negative Difference Density on H16A -0.54 eA-3
[PLAT977_ALERT_2_C](#) Check Negative Difference Density on H22B -0.38 eA-3
[PLAT978_ALERT_2_C](#) Number C-C Bonds with Positive Residual Density. 0 Info

● Alert level G

[PLAT072_ALERT_2_G](#) SHELXL First Parameter in WGHT Unusually Large 0.18 Report
[PLAT154_ALERT_1_G](#) The s.u.'s on the Cell Angles are Equal ..(Note) 0.003 Degree
[PLAT910_ALERT_3_G](#) Missing # of FCF Reflection(s) Below Theta(Min). 1 Note
[PLAT912_ALERT_4_G](#) Missing # of FCF Reflections Above STh/L= 0.600 72 Note

0 **ALERT level A** = Most likely a serious problem - resolve or explain
0 **ALERT level B** = A potentially serious problem, consider carefully
9 **ALERT level C** = Check. Ensure it is not caused by an omission or oversight
4 **ALERT level G** = General information/check it is not something unexpected

1 ALERT type 1 CIF construction/syntax error, inconsistent or missing data
10 ALERT type 2 Indicator that the structure model may be wrong or deficient
1 ALERT type 3 Indicator that the structure quality may be low
1 ALERT type 4 Improvement, methodology, query or suggestion
0 ALERT type 5 Informative message, check