

Neuromechanical differences between successful and failed sit-to-stand movements and response to rehabilitation early after stroke.

Andy Kerr PhD <sup>1</sup>, Allan Clark PhD<sup>2</sup>, Valerie M. Pomeroy PhD <sup>2</sup>

<sup>1</sup>University of Strathclyde, Glasgow, United Kingdom

<sup>2</sup>University of East Anglia, Norwich, United Kingdom

Corresponding author

Dr Andy Kerr, Dept. of Biomedical Engineering, University of Strathclyde,

Email [a.kerr@strath.ac.uk](mailto:a.kerr@strath.ac.uk)

Tel +44 (0)141 548 2855

Word count 3,634

Tables 3

Figures 0

### Abstract

#### Background

Recovery of the sit-to-stand (StS) movement early after stroke could be improved by targeting physical therapy at the underlying movement deficits in those people likely to respond.

#### Aim

To compare the movement characteristics of successful and failed StS movements in people early after stroke and identify which characteristics change in people recovering their ability to perform this movement independently following rehabilitation.

#### Methods

Muscle activity and kinematic (including centre of mass, CoM) data were recorded from 91 participants (mean 35 days after stroke) performing the StS movement before (baseline), immediately after (outcome) and three months after (follow-up) rehabilitation. Three sub-groups (never-able (n=19), always-able (n=51) and able-after-baseline, n=21) were compared at baseline with the able-after-baseline sub-group compared before and after rehabilitation.

#### Results

The sub-groups differed at baseline for quadriceps onset time ( $p=0.009$ ) and forward body position when quadriceps peaked ( $p=0.038$ ). Following rehabilitation the able-after-baseline sub-group increased their forward position ( $p<0.001$ ), decreased the time difference between bilateral quadriceps peaks ( $p<0.001$ ) and between

quadriceps and hamstrings peaks on the non-hemiplegic side ( $p=0.007$ ). An improved performance in the always-able sub-group was associated with a number of baseline factors including forward positioning ( $p=0.002$ ) and time difference between peak activity of bilateral quadriceps ( $p=0.001$ ).

### Conclusions

This neuromechanical study of StS before and after rehabilitation in a sample of people early after stroke identified the importance of temporal coupling between forward trunk movement and quadriceps and hamstrings' activity. These findings advance the science of stroke rehabilitation by providing evidence based therapy targets to promote recovery of the StS movement.

Keywords: Stroke rehabilitation; physical therapy; sit-to-stand; kinematics; EMG; therapy targets.

### Introduction

Rehabilitation is known to improve physical function after a stroke [1]. Movement repetition is the key principle for this rehabilitation [2] with the quality of movement considered important in traditional theories [3]. Clarifying the characteristics of a movement that should be emphasised during practice could optimise outcomes through tailoring rehabilitation activities (targeted therapy). This could be achieved by distinguishing the kinematic and muscle activity variations between successful and failed attempts at functional tasks performed by impaired populations as well identifying the variables associated with improved ability after rehabilitation.

Considered to be one of the most physically challenging functional movements [4], the sit-to-stand (StS) task is nevertheless performed frequently as part of everyday activities [5]. An impaired ability will inevitably threaten an individual's ability to live independently [6]. Recovering this movement is therefore included in stroke rehabilitation guidelines with a recommendation for frequent practice [7].

Laboratory based biomechanical studies of healthy people have identified two key features of a successful StS movement. First a forward trunk lean while seated measured between  $22.2 \pm 7.8^\circ$  [8] and  $38.8 \pm 8.1^\circ$  [4], to the vertical. This movement is typically performed at velocities around  $87^\circ/\text{s}$  [9]. Although this relatively high velocity is considered desirable in terms of efficiency it is not essential to success [10]. Second, the generation of a downward force on the ground that exceeds body weight during the rising phase [11-13]. Studies of failed StS attempts have complemented these findings. Reduced forward displacement while seated and vertical forces less than body weight have both been described during failed attempts

by older adults (n=13) affected, predominantly, by vestibular hypofunction and in a single case of traumatic brain injury [14, 15]. While contributing to our understanding of StS these findings have a limited influence on stroke rehabilitation practice.

A biomechanical approach to analysing movement is clearly informative but provides only indirect evidence of the underlying muscle activity. This is particularly limiting if the population of interest has a neurological impairment.

Electromyographical (EMG) studies of the StS movement in healthy participants point to a consistent temporal pattern of muscle activation, in particular, the agonist muscle activity (hip and knee extensors) needed to create the vertical force to lift the body up during the rising phase. Quadriceps and hamstring muscle groups are observed to reach simultaneous levels of peak activity around the commencement of the rising phase [16-19]. Generating the force necessary to lift the body is clearly easier if both legs contribute equally [20]. Symmetrical activation of the hip and knee extensors to produce this net moment, while not an absolute determinant of success, is likely to be an important factor [11] for safe and frequent movement. EMG studies of the StS movement among people after a stroke are inconsistent in their findings but generally report altered muscle activation times on the hemiplegic side [17, 21, 22]. The value of these studies to understanding recovery of this movement after stroke is limited by the recruitment of participants long after their stroke (e.g. 3.5 years [21]), lack of data before and after rehabilitation and not including failed StS attempts in the analysis.

This study aimed to provide therapy targets for StS training early after stroke by comparing the neuromechanical characteristics of successful and failed StS movements and then identifying the movement characteristics associated with a

change in ability (i.e. changing from failed to successful movement) and performance immediately following a six-week rehabilitation intervention and three months thereafter.

### **Methods**

The study reported here was embedded in an observer-blind, multi-centre, randomized controlled early phase trial. Full protocol details, including the intervention, are provided in an earlier publication [23].

Briefly, baseline measures were conducted with participants who were subsequently randomised to one of three sub-groups: conventional physiotherapy (CPT); CPT+ movement performance therapy (MPT); or CPT+ functional strength training (FST). The intervention period lasted six weeks. Outcome measures were recorded immediately after the end of the intervention period and 12 weeks thereafter. The local Research Ethics Committees covering the clinical centres granted ethical approval (Merton and Sutton Ethics Committee, 205cl01). The trial was registered on a clinical trials database (NCT00322192).

Our previous publications reported no statistical difference in StS performance across the three intervention sub-groups [23, 24]. The analysis reported here therefore considers the sample as a single group.

### Participants

All participants provided written informed consent and met the following criteria:

- In-patients aged 18+ years who were between 1 and 13 weeks of a stroke (haemorrhage or infarction) in the territory of the anterior circulation.

- Able to produce some voluntary contraction of their paretic lower limb muscle (i.e. scoring at least 28/100 on lower limb section of the Motricity Index [25]).
- Able to follow one-step verbal commands and were independently mobile, with or without aids, prior to the index stroke with no lower limb orthopaedic surgery or trauma in the lower limb in the previous 8 weeks or any previous history of neurological disease other than stroke.

### Interventions

The CPT group received 9.2 (SD 6.9) hours of rehabilitative exercises/activities as determined by the clinical physiotherapist which included, but was not limited to: soft tissue mobilisation, facilitation of muscle activity, facilitation of coordinated multi-joint movement, tactile and proprioceptive input, resistive exercise, and functional retraining [23].

The CPT+MPT group received 23.0 (SD 10.4) hours of an intervention that combined CPT (as described above) and MPT, which emphasised the recovery of movement quality through movement repetition with facilitation and feedback provided by a therapist [23].

The CPT + FST group received 23.5 (SD 10.0) hours of an intervention that combined CPT (as described above) and FST, which focused on repetitive, progressive resistive exercise during goal-directed functional activities. The differential for this treatment group was the use of graded resistance by manipulating the gravitational moment acting on the whole body and limbs, increasing the range of movement or distance over which bodyweight was transported, and simply changing the weight of external objects [23].

### The StS movement task

Participants were instructed to stand up five times from a plinth adjusted to the height of their knee. The starting position for all participants was sitting on the plinth with both thighs parallel to the ground, both ankles directly below the knees and the trunk perpendicular to the thighs. The upper limbs were positioned so that they hung freely alongside the thighs. Participants were asked not to use their hands for support or to push up from the plinth during the movement attempt. Participants wore tight fitting clothing to improve the quality of the motion capture and footwear usually worn for therapy sessions. To ensure the activity did not result in a fall, a physiotherapist supervised all attempts but did not provide any physical assistance. An audio trigger instructed the participant to commence the task, this trigger was also used to synchronise the measurement instruments.

Only movements that met the task success criteria (no hands, aids or support used) and had every segment tracked throughout the movement with complete data, were used for analysis. In cases where the movement was not successfully completed (i.e. a failed attempt) the first clear purposeful attempt (where the body moves forward with a clear attempt to lift the body) was used for analysis.

### Data collection and analysis

#### *Kinematic data*

Eight Vicon motion cameras (Oxford Metrics, Oxford, UK) captured the 3D trajectories of retro-reflective markers attached to the body at anatomical locations at a sampling rate of 120Hz. The trajectories were then processed by filling any small gaps (<10 frames) with a spline fill function and filtered with a low pass (cut off



frequency 6Hz) sixth order Butterworth filter to reduce the noise content of the signal. Once processed a geometric model, consisting of 15 body segments (head, trunk, upper arms, forearms, hands, pelvis, thighs, lower legs and feet), was constructed using proprietary software (Nexus, Oxford Metrics, Oxford, UK) which allowed anatomical angles and the overall centre of mass (CoM) to be calculated from the weighted average of the 15 tracked segments. These data were used to derive the following temporal events commonly used in sit to stand analysis.

- 1) Onset event: First occurrence (time point) of a forward movement of the CoM that lasted at least 10 frames.
- 2) Seat-off event: First occurrence (time point) of increasing upwards CoM velocity that lasted at least 10 frames.
- 3) End event: First occurrence (time point) of the CoM reaching its maximal vertical displacement.

These events allowed the movement to be separated into two phases; Phase one: onset to seat-off, Phase two: seat-off to end, as well as a value for total movement duration (onset to end).

### *Muscle activity*

A surface telemetric electromyography (EMG) system (MT8, MIE Medical Research Ltd.) was used to record the electrical activity of two agonist/antagonist muscle pairs: quadriceps/hamstrings and tibialis anterior/gastrocnemius on both-sides during the task. Data from eight muscle groups were therefore available for analysis, however, only the hamstrings and quadriceps muscles on both sides were analysed. The

consistent pattern of their activity, close temporal coupling and function as prime movers to lift the body distinguished these muscles as being fundamental to movement success compared to the tibialis anterior/gastrocnemius pair which are more inconsistent in their activity and considered to act, primarily, as stabilisers and not prime movers [18]. The system sampled the data at 1080 Hz and was synchronised with the kinematic capture software (Nexus, Oxford Metrics, UK) using a synchronisation pulse. The preparation and positioning of the EMG electrodes followed SENIAM guidelines [26]. The raw EMG signals were processed into a linear envelope (full wave rectification low pass filter (cut off frequency 6Hz) and integration) using a custom-made Matlab (Mathworks Inc. Massachusetts, USA) program. Muscle onset times were identified from a threshold detection algorithm of the signal exceeding a baseline mean (+ 3 standard deviations) [18] which was calculated from 1000 frames preceding the start signal. The time of peak muscle activity was identified with a peak detection function (Matlab, Mathworks Inc. Massachusetts, USA). Muscle onsets and peaks were timed relative to the movement onset event. Temporal relationships between muscle pairs (hamstrings and quadriceps on each side and quadriceps on opposite sides) were calculated as an absolute time difference.

Demographic (age and gender) and stroke specific information (time since stroke, and hemiplegic side) were recorded from the medical notes. The presence of neglect was assessed with the star cancellation test [27] with a score between 0 and 49 indicating neglect. The Modified Rivermead Mobility Index [28] and gait speed (measured over a flat 10 m walkway) were recorded by the research team.

### Statistical analysis

The demographic, stroke, clinical and neuromechanical variables at baseline were summarised for the three separate sub-groups: never-able; always-able; and able-after-baseline. The neuromechanical variables were selected *a priori* based on the available literature. Chi-squared and T-tests were used to compare the able-after-baseline and never-able sub-groups. A one-factor anova tested for statistically significant changes in the neuromechanical variables of the able-after baseline sub-group between baseline, outcome and follow-up.

Finally a statistical analysis was undertaken to identify factors associated with improved StS performance after rehabilitation in the always-able sub-group (n=51), this was the only sub-group with successful attempts at each time point. A linear regression model was fitted using the outcome of movement time separately for each variable and then a variable selection technique was used to identify which factors were independently associated with improvement.

All analyses were conducted using standard statistical software packages (Minitab, version 17 & Stata version 14).

### **Results**

Full data sets were available from 91 participants at three time points (baseline, outcome and follow up, a consort diagram is available in the supplementary information.

#### Baseline differences

The baseline characteristics were described for the three sub-groups and statistically tested between the never-able and able-after baseline sub-groups (see table 1).

CoM position at the time of peak quadriceps activity (hemiplegic side) was

statistically different ( $p=0.038$ ), with the never-able sub-group less far forward ( $68.86, \pm 82.70\text{mm}$ ) than the able-after-baseline sub-group ( $128.87 \pm 73.55\text{mm}$ ). Quadriceps onset time (hemiplegic side) was also statistically different ( $p=0.009$ ), occurring much earlier in the never-able sub-group ( $0.91 \pm 0.63\text{s}$ ) compared to the able-after-baseline sub-group ( $1.73 \pm 1.08\text{s}$ ).

### Response to rehabilitation

As the only participants that recovered their StS ability (i.e. from being unable to being able), the movement characteristics of the able-after-baseline sub-group were described at each time point (see table 2). Four statistically significant differences were found using a one factor (time), ANOVA:

1. Peak forward position of the CoM increased from  $173.97\text{mm}$  ( $69.17$ ) at baseline to  $273.20\text{mm}$  ( $57.60$ ) at outcome and  $279.30\text{mm}$  ( $44.30$ ) by follow-up ( $p<0.001$ ).
2. CoM forward position at the time of peak quadriceps on both the hemiplegic ( $p<0.001$ ) and non-hemiplegic sides ( $p<0.001$ ). At the time of peak quadriceps activity on the hemiplegic side, the CoM forward position increased from  $128.87$  ( $73.55$ ) mm at baseline, to  $226.30$  ( $71.40$ ) mm at outcome and was unchanged at follow-up. On the non-hemiplegic side, the CoM forward position, at the time of peak quadriceps, increased from  $114.00\text{mm}$  ( $164.90$ ) at baseline to  $220.30\text{mm}$  ( $78.40$ ) at outcome and then  $231.90$  ( $43.60$ ) mm by follow-up.
3. The time difference between peak quadriceps activity on the hemiplegic and non-hemiplegic sides changed significantly over time ( $p=0.000$ ). The time difference decreased from  $1.38\text{s}$  ( $1.25$ ) at baseline to  $0.45\text{s}$  ( $0.59$ ) at outcome and then slightly increased to  $0.52\text{s}$  ( $0.83$ ) at follow-up.

4. The time difference between hamstrings and quadriceps peaks on the non-hemiplegic side changed significantly over time ( $p=0.007$ ), whereas a significant difference was not found for the hemiplegic side ( $p=0.058$ ). The time difference between hamstring and quadriceps peaks on the non-hemiplegic side decreased from 1.22s (SD1.00) at baseline to 0.41s (0.52) at outcome and then increased to 0.67s (0.80) at follow-up.

Insert table 1 here

Insert table 2 here

#### Baseline characteristics associated with improved performance

At baseline the movement duration (onset until end) for the always-able sub-group was 3.51s (2.01), this decreased to 2.56s (1.38) at outcome ( $p=0.001$ ) and to 2.34s (1.39) by follow up ( $p=0.000$ ). The linear regression analysis found 11 neuromechanical and one stroke related baseline characteristic (neglect) to be significantly associated with this improvement by follow up (Table 3). The neuromechanical characteristics related to the temporal synergy across the muscles involved in raising the body (quadriceps and hamstrings on the hemiplegic and non-hemiplegic sides) and ability to bring the body forward. Participants with neglect showed less improvement.

Insert table 3 here

#### **Discussion**

This study is the first study to describe the neuromechanical differences between successful and failed StS movement attempts in people early after a stroke and then track these movement variables in response to rehabilitation. With an aim of identifying therapy targets, analysis focussed on the sub-group of people who recovered StS independence following rehabilitation i.e. participants who changed from being unable to being able. In sum, the able-after-baseline sub-group differed statistically at baseline from the never-able sub-group in their ability to time peak quadriceps activity when the body was much further forward. Following rehabilitation, this sub-group increased the forward movement of their body, reduced the time difference between peak quadriceps' bilaterally and between quadriceps and hamstrings on the non-hemiplegic side.

Unsurprisingly the sub-groups, never-able (n=19), always-able (n=51) and able-after-baseline (n=21) differed statistically at baseline. These differences related to forward movement of the body (CoM) and timing quadriceps' activity to when a greater amount of this forward movement had occurred. This confirms previous reports on the importance of bringing the body forward before generating peak extensor moments at the hip and knee, to lift the body mass [29]. Generating peak knee extensor activity when the body mass has not been brought sufficiently far forward (e.g. only 6cm in the never-able sub-group) is likely to result in a fall back [30]. Elements of the basic movement pattern appear to be retained in the able-after-baseline sub-group with forward displacement and quadriceps timings similar to the always-able sub-group.

Peak activity of the hamstring muscle group was timed closely with the quadriceps peak in the always-able sub-group (hemiplegic side 0.61s, non-

hemiplegic side 0.51s). This co-contraction of agonist/antagonist pairs might seem counterintuitive but has been observed before during StS movements [31] probably due to the need to generate large hip and knee extensor moments, simultaneously, around seat-off.

While age and time since stroke did not vary significantly between the never-able and able-after-baseline sub-groups at baseline it is notable that the never-able sub-group were both the oldest and had the longest delay before rehabilitation. Age is consistently cited as a factor in functional recovery [32] and a delay in rehabilitation raises the possibility of confounding factors such as co-morbidities. The lack of statistical variance across the sub-groups for mobility (Modified Rivermead Mobility Index, MRMI) and stroke specific factors, such as neglect and hemiplegic side, were surprising but may relate to the small sub-group sizes. Using a baseline MRMI score of 18.5 Shum et al. [33] predicted 88% of patients being able to walk one month after stroke, interestingly, in our study, this cut-off was exceeded by the able-after-baseline sub-group (24.20) but very similar to the never-able sub-group (19.00).

Two sub-groups improved their StS ability following rehabilitation, 19/38 (50%) of individuals unable to perform the movement at baseline, changed to being independent in the movement at follow-up, a success rate similar to previous studies [34]. The movement characteristics that changed in these individuals is arguably the most interesting part of this study as it provides potential therapy targets. Greater forward movement and closer timing between bilateral quadriceps and unilateral hamstrings/quadriceps point to a development in movement skill following rehabilitation, with greater co-ordination between muscles to generate vertical lift at the optimal time [17]. Close timing between quadriceps and hamstrings has been

reported previously with Khemlani et al. observing virtually synchronous onsets of the rectus and biceps femoris muscles in healthy adults [18]. Finding closer muscle timings between the hemiplegic and non-hemiplegic sides, builds on previous reports of improved symmetry following rehabilitation in this population [35] [36].

Statistically faster performance times after rehabilitation are common among stroke populations already independent in the StS movement [1, 34]. In our study this improvement was not only associated with earlier muscle timings relative to movement onset, as might be expected in a faster movement, but also the retention after stroke of a close synergy between the forward positioning of the body and peak activity of bilateral quadriceps and ipsilateral hamstrings and quadriceps on the non-hemiplegic side. These factors, it would appear, are not only crucial for recovering movement independence, as discussed earlier, but also for decreasing movement time which is a commonly used outcome measure in rehabilitation trials [1].

### Limitations

A strength of the present study was the pre-selection of a relatively small number of variables clustered around muscle timing (onset and peak) and position of the centre of mass which were informed by previous findings [17, 21, 34, 35, 37]. Consequently the study was hypothesis driven. However, this pre-selection does raise the possibility that we missed an important factor. Knee extension velocity, for example, has been identified as a sensitive measurement for StS [38]. A subsequent study could explore a more extensive range of variables.

The decision to remove tibialis anterior and gastrocnemius muscles from the analysis was intended to reduce the data by focussing on the muscles generating the extension moments for the lifting phase, considered the most demanding part of the



StS movement [6, 31]. Tibialis anterior and gastrocnemius activity has been observed before during the StS movement, particularly during the preparatory period to stabilise the foot and lower leg and may have been useful in understanding motor control in this sub-group. Future studies might explore the role of other muscles during successful and failed attempts before and after rehabilitation.

Another potential limitation is the sample size. Although larger than most studies of this nature, it is possible that some of the associations found in this study might not be representative of a larger population. We recommend these findings are tested in future larger studies.

Despite these limitations, this was a robustly executed study of the StS movement before and after rehabilitation in a large sample of people early after stroke. The findings are an important step towards clarifying the important motor features of a successful StS and identifying potential factors for predicting responders to rehabilitation so that interventions can become more tailored and effective [39] .

### Practice implications

Based on our findings, a forward displacement of 20cm while seated would seem a reasonable therapy target. This can be created by a forward trunk lean of approximately 25-30 degrees to the vertical, which is a more clinically accessible metric. The other key targets for rehabilitation are the timings of peak activity in the quadriceps and hamstrings muscle groups relative to each other and the forward position of the body. Although this information is well known amongst physiotherapists, [40], the present findings confirm this clinical knowledge and provide numerical references. Incorporating this knowledge into the provision of

feedback for individuals after stroke would provide a practical way to translate this knowledge into clinical practice. For example, EMG biofeedback systems could be used to provide muscle timing information, perhaps combined with simple motion tracking systems to promote the synchronisation between the forward lean and the quadriceps/hamstrings synergy. These findings will help design future prospective trials targeted at recovering independence and improving performance in this important everyday movement.

### **Conclusion**

This large study of failed and successful StS movements before and after rehabilitation in people early after stroke provides therapy targets for regaining independence and improving movement performance. The temporal coupling of bilateral quadriceps with hamstrings activity to coincide with the forward positioning of the body differed significantly between failed and successful attempts before rehabilitation and then changed significantly in the sub-group of participants who regained independence in this movement. This study provides the evidence to support therapy targeting these movement characteristics and lays the foundation to develop feedback methods, including the use of technology, to promote the recovery of this important everyday movement.

### **Acknowledgements**

We gratefully acknowledge funding provided by The Healthcare Foundation for the trial into which this experimental study was embedded. We also thank Dr Emma Cooke and Professor Raymond Tallis for their contributions to the concept, design

and conduction of the previously published clinical efficacy trial. In addition we thank Ashraf Sameja for assistance with data collection.

The authors declare that there are no conflicts of interest.

## References

- [1] Pollock A, Baer G, Campbell P, Choo PL, Forster A, Morris J, et al. Physical rehabilitation approaches for the recovery of function and mobility following stroke. *Cochrane Database of Systematic Reviews*. 2014(4).
- [2] Lohse KR, Lang CE, Boyd LA. Is more better? Using metadata to explore dose-response relationships in stroke rehabilitation. *Stroke*. 2014 Jul;45(7):2053-8.
- [3] Langhorne P, Bernhardt J, Kwakkel G. Stroke rehabilitation. *The Lancet*. 2011;377(9778):1693-702.
- [4] Dubost V, Beauchet O, Manckoundia P, Herrmann F, Mourey F. Decreased trunk angular displacement during sitting down: an early feature of aging. *Physical Therapy*. 2005;85(5):404-12.
- [5] Grant PM, Dall PM, Kerr A. Daily and hourly frequency of the sit to stand movement in older adults: a comparison of day hospital, rehabilitation ward and community living groups. *Aging Clinical and Experimental Research*. 2011;23(5-6):437-44.
- [6] Yoshioka S, Nagano A, Hay DC, Fukashiro S. The minimum required muscle force for a sit-to-stand task. *Journal of Biomechanics*. 2012 2012/02/23;45(4):699-705.
- [7] SIGN. Management of patients with stroke: rehabilitation, prevention and management of complications, and discharge planning. In: Scotland NQI, ed. Edinburgh: SIGN 2010.
- [8] Ganea R, Paraschiv-Ionescu A, Büla C, Rochat S, Aminian K. Multi-parametric evaluation of sit-to-stand and stand-to-sit transitions in elderly people. *Medical Engineering & Physics*. 2011 2011/11/01;33(9):1086-93.
- [9] Ikeda ER, Schenkman ML, Riley PO, Hodge WA. Influence of Age on Dynamics of Rising from a Chair. *Physical Therapy*. 1991;71(6):473-81.
- [10] Mazzà C, Stanhope SJ, Taviani A, Cappozzo A. Biomechanic Modeling of Sit-to-Stand to Upright Posture for Mobility Assessment of Persons With Chronic Stroke. *Archives of Physical Medicine and Rehabilitation*. 2006 2006/05/01;87(5):635-41.
- [11] Burnett DR, Campbell-Kyureghyan NH, Cerrito PB, Quesada PM. Symmetry of ground reaction forces and muscle activity in asymptomatic subjects during walking, sit-to-stand, and stand-to-sit tasks. *Journal of Electromyography and Kinesiology*. 2011 2011/08/01;21(4):610-5.
- [12] Houck J, Kneiss J, Bukata SV, Puzas JE. Analysis of vertical ground reaction force variables during a Sit to Stand task in participants recovering from a hip fracture. *Clinical Biomechanics*. 2011 2011/06/01;26(5):470-6.

- [13] Vander Linden DW, Brunt D, McCulloch MU. Variant and invariant characteristics of the sit-to-stand task in healthy elderly adults. *Archives of Physical Medicine and Rehabilitation*. 75(6):653-60.
- [14] Riley PO, Krebs DE, Popat RA. Biomechanical analysis of failed sit-to-stand. *IEEE Transactions on Rehabilitation Engineering*. 1997;5(4):353-9.
- [15] Zablony CM, Nawoczenski DA, Yu B. Comparison between successful and failed sit-to-stand trials of a patient after traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*. 2003 2003/11/01;84(11):1721-5.
- [16] Rodrigues-de-Paula Goulart F, Valls-Solé J. Patterned electromyographic activity in the sit-to-stand movement. *Clinical Neurophysiology*. 1999 1999/09/01;110(9):1634-40.
- [17] Cheng P-T, Chen C-L, Wang C-M, Hong W-H. Leg Muscle Activation Patterns of Sit-to-Stand Movement in Stroke Patients. *American Journal of Physical Medicine & Rehabilitation*. 2004;83(1):10-6.
- [18] Khemlani MM, Carr JH, Crosbie WJ. Muscle synergies and joint linkages in sit-to-stand under two initial foot positions. *Clinical Biomechanics*. 1999;14:236-46.
- [19] Boukadida A, Piotte F, Dehail P, Nadeau S. Determinants of sit-to-stand tasks in individuals with hemiparesis post stroke: a review. *Annals of physical and rehabilitation medicine*. 2015;58(3):167-72.
- [20] Hirschfeld H, Thorsteinsdottir M, Olsson E. Coordinated ground forces exerted by buttocks and feet are adequately programmed for weight transfer during sit-to-stand. *Journal of Neurophysiology*. 1999;82(6):3021-9.
- [21] Prudente C, Rodrigues-de-Paula F, Faria CD. Lower limb muscle activation during the sit-to-stand task in subjects who have had a stroke. *American journal of physical medicine & rehabilitation*. 2013;92(8):666-75.
- [22] Cheng PT, Chen CL, Wang CM, Hong WH. Leg muscle activation patterns of sit-to-stand movement in stroke patients. *American Journal of Physical Medicine & Rehabilitation*. 2004;83(1):10-6.
- [23] Cooke EV, Tallis RC, Clark A, Pomeroy VM. Efficacy of functional strength training on restoration of lower-limb motor function early after stroke: phase I randomized controlled trial. *Neurorehabilitation & Neural Repair*. 2010 Jan;24(1):88-96.
- [24] Kerr A, Clark A, Cooke EV, Rowe P, Pomeroy VM. Functional strength training and movement performance therapy produce analogous improvement in sit-to-stand early after stroke: early-phase randomised controlled trial. *Physiotherapy*. 2017.
- [25] Collin C, Wade D. Assessing motor impairment after stroke: a pilot reliability study. *Journal of Neurology, Neurosurgery & Psychiatry*. 1990;53(7):576-9.
- [26] Hermens HJ, Freriks B, Merletti R, Stegeman D, Blok J, Rau G, et al. European recommendations for surface electromyography. *Roessingh research and development*. 1999;8(2):13-54.
- [27] Friedman PJ. The star cancellation test in acute stroke. *Clinical Rehabilitation*. 1992;6(1):23-30.
- [28] Lennon S, Johnson L. The modified rivermead mobility index: validity and reliability. *Disability and rehabilitation*. 2000;22(18):833-9.
- [29] Janssen WGH, Bussmann HBJ, Stam HJ. Determinants of the sit-to-stand movement: A review. *Physical Therapy*. 2002;82(9):866-79.
- [30] Riley PO, Krebs DE, Popat RA. Biomechanical analysis of failed sit to stand. *IEEE Transactions on Rehabilitation Engineering*. 1997 359;5(4):353.

- [31] Roebroeck ME, Doorenbosch CAM, Harlaar J, Jacobs R, Lankhorst GJ. Biomechanics and muscular activity during sit-to-stand transfer. *Clinical Biomechanics*. 1994;9:235-44.
- [32] Mutai H, Furukawa T, Araki K, Misawa K, Hanihara T. Factors associated with functional recovery and home discharge in stroke patients admitted to a convalescent rehabilitation ward. *Geriatrics & gerontology international*. 2012;12(2):215-22.
- [33] Shum S-T, Chiu JK-W, Tsang CP-L, Wong CH-P, Tsang RC-C, Ma S-L, et al. Predicting walking function of patients one month poststroke using modified Rivermead mobility index on admission. *Journal of Stroke and Cerebrovascular Diseases*. 2014;23(8):2117-21.
- [34] Janssen W, Bussmann J, Selles R, Koudstaal P, Ribbers G, Stam H. Recovery of the sit-to-stand movement after stroke: a longitudinal cohort study. *Neurorehabilitation and neural repair*. 2010;24(8):763-9.
- [35] Kerr A, Clark A, Cooke EV, Rowe P, Pomeroy VM. Functional strength training and movement performance therapy produce analogous improvement in sit-to-stand early after stroke: early-phase randomised controlled trial. *Physiotherapy*. 2016 2016/02/11/.
- [36] Liu M, Chen J, Fan W, Mu J, Zhang J, Wang L, et al. Effects of modified sit-to-stand training on balance control in hemiplegic stroke patients: A randomized controlled trial. *Clinical Rehabilitation*. 2015 August 27, 2015.
- [37] Linden DWV, Brunt D, McCulloch MU. Variant and invariant characteristics of the sit-to-stand task in healthy elderly adults. *Archives of Physical Medicine and Rehabilitation*. 1994;75:653-60.
- [38] Richards JD, Pramanik A, Sykes L, Pomeroy VM. A comparison of knee kinematic characteristics of stroke patients and age-matched healthy volunteers. *Clinical Rehabilitation*. 2003 Aug;17(5):565-71.
- [39] Jeffers MS, Karthikeyan S, Gomez-Smith M, Gasinzigwa S, Achenbach J, Feiten A, et al. Does Stroke Rehabilitation Really Matter? Part B: An Algorithm for Prescribing an Effective Intensity of Rehabilitation. *Neurorehabilitation and Neural Repair*. 2018;32(1):73-83.
- [40] Carr JH. *Neurological Rehabilitation*, 2/e: Elsevier India 2011.

Table 1: Summary of participants and their sit to stand neurobiomechanical characteristics, mean (SD) and count (percentage) at baseline, separated into the three sit-to-stand (StS) ability sub-groups.

<b>Characteristic</b>	<b>Never-able (n=19)</b>		<b>Always-able (n=51)</b>		<b>Able-after-baseline (n=21)</b>	
<b>Gender (male)</b>	10 (52.6%)		32 (62.8%)		11 (52.4%)	
<b>Age (years)</b>	74.58 (7.34)		65.78 (13.74)		70.43 (9.46)	
<b>Time since stroke (days)</b>	43.74 (21.87)		27.9 (17.94)		33.05 (18.81)	
<b>Neglect present</b>	5 (26.3%)		6 (11.8%)		7 (33.3%)	
<b>Walking Speed</b>	0.00 (0.00)		0.37 (0.33)		0.05 (0.14)	
<b>Modified Rivermead Mobility Index</b>	19.00 (6.06)		34.28 (8.63)		24.20 (7.95)	
<b>Hemiplegic side (right)</b>	7 (36.8%)		22 (43.1%)		3 (14.3%)	
<b>Peak forward position of CoM (mm)</b>	136.23 (87.61)		259.11 (52.87)		173.97 (69.17)	
<b>Time of peak trunk flexion after onset (s)</b>	5.06 (1.91)		2.67 (1.28)		2.98 (1.52)	
<b>Quadriceps onset time(s) after movement onset</b>	H-side	Non- side	H-side	Non- side	H-side	Non- side
	0.91 (0.63)	2.09 (2.28)	1.33 (1.06)	1.44 (1.26)	1.73 (1.08)	1.69 (1.00)
<b>Difference between quadriceps peaks (s)</b>	1.65 (1.32)		0.9 (1.16)		1.38 (1.25)	
<b>*Quadriceps peak time (s) after movement onset</b>	H-side	Non- side	H-side	Non- side	H-side	Non- side
	3.38 (2.23)	3.76 (2.29)	2.67 (1.79)	2.80 (1.86)	3.00 (1.38)	2.92 (1.55)
<b>Hamstrings peak time (s) after movement onset</b>	H-side	Non- side	H-side	Non- side	H-side	Non- side
	3.6 (1.85)	3.42 (2.08)	2.56 (1.61)	2.58 (1.5)	4.23 (5.58)	3.00 (1.21)
<b>Absolute time difference between quadriceps and hamstrings peaks (s)</b>	H-side	Non- side	H-side	Non- side	H-side	Non- side
	1.67 (1.21)	1.23 (1.30)	0.62 (1.03)	0.51 (0.76)	2.51 (5.36)	1.22 (1.00)
<b>*CoM forward position at time of peak quadriceps (mm)</b>	H-side	Non- side	H-side	Non- side	H-side	Non- side
	68.86 (82.70)	98.92 (56.75)	202.31 (73.42)	211.37 (68.26)	128.87 (73.55)	113.97 (64.89)

H-side = hemiplegic side, Non-side = Non hemiplegic side, \*indicates a characteristic that was statistically significantly ( $p < 0.05$ ) between the never-able and able-after-baseline sub-groups.

Table 2: Neuromechanical characteristics of the able-after-baseline sub-group at baseline, outcome and follow-up

<b>Characteristic</b>	<b>Baseline (n=21)</b>		<b>Outcome (n=21)</b>		<b>Follow up (n=19)</b>	
<b>Peak forward position of CoM (mm)</b>	174.00 (69.17)		273.20 (57.60)		279.30 (44.30)	
<b>Time of peak trunk flexion after onset (s)</b>	2.98 (1.52)		3.00 (1.15)		3.32 (1.75)	
<b>Walking speed (m/s)</b>	0.05 (0.13)		0.60 (1.55)		0.38 (0.40)	
<b>Rivermead</b>	24.2 (7.95)		33.95 (8.78)		36.95 (7.06)	
<b>Quadriceps peak time (s) after movement onset</b>	H-side 3.00 (1.38)	Non-side 2.92 (1.55)	H-side 2.92 (1.20)	Non-side 3.17 (1.51)	H-side 3.65 (2.05)	Non-side 4.04 (2.16)
<b>Hamstring peak time (s) after movement onset</b>	H-side 4.23 (5.58)	Non-side 3.00 (1.21)	H-side 2.76 (1.20)	Non-side 3.03 (1.42)	H-side 3.53 (2.04)	Non-side 3.90 (2.34)
<b>Quadriceps onset time (s) after movement onset</b>	H-side 1.73 (1.08)	Non-side 1.69 (0.99)	H-side 1.61 (1.22)	Non-side 2.01 (1.45)	H-side 1.52 (0.88)	Non-side 2.04 (1.24)
<b>Time difference between quadriceps peaks (s)</b>	1.38 (1.25)		0.45 (0.59)		0.52 (0.83)	
<b>Time difference between quadriceps and hamstring peaks (s)</b>	H-side 2.51 (5.36)	Non-side 1.22 (1.00)	H-side 0.33 (0.51)	Non-side 0.41 (0.52)	H-side 0.50 (0.84)	Non-side 0.67 (0.80)
<b>CoM forward position at time of peak quadriceps (mm)</b>	H-side 128.87 (73.55)	Non-side 114.0 (164.90)	H-side 226.3 (71.40)	Non-side 220.3 (78.40)	H-side 222.7 (49.1)	Non-side 231.9 (43.6)

Table 3: Association of baseline characteristics with an improved StS movement time at follow-up for the always-able sub-group.

<b>Characteristic</b>	<b>Baseline characteristics</b>	<b>Regression coefficient (95% CI)</b>	<b>p-value</b>
<b>Gender (male)</b>	32 (62.8%)	0.58 (-0.72,1.90)	0.370
<b>Age (years)</b>	65.78 (13.74)	-0.01 (-0.06,0.03)	0.534
<b>Time since stroke (days)</b>	27.9 (17.94)	0.02 (-0.01,0.05)	0.188
<b>Neglect</b>	6 (11.8%)	2.40 (0.62,4.18)	0.010*
<b>Hemiplegic side (right)</b>	22 (43.1%)	-0.06 (-1.26,1.14)	0.917
<b>Walking speed</b>	0.37 (0.33)	1.22 (-0.49,2.93)	0.156
<b>Modified Rivermead Mobility Index</b>	34.28 (8.63)	0.03 (-0.04,0.10)	0.442
<b>Forward position of centre of mass</b>	259.11 (52.87)	-0.01 (-0.02,-0.00)	0.041*
<b>Quads peak time (hemiplegic)</b>	2.67 (1.79)	-0.39 (-0.71,-0.08)	0.015*
<b>Quads peak time (non-hemiplegic)</b>	2.80 (1.86)	-0.43(-0.72,-0.14)	0.004*
<b>Hamstring peak time (hemiplegic)</b>	2.56 (1.61)	-0.42 (-0.79,-0.04)	0.030*
<b>Hamstring peak time (non-hemiplegic)</b>	2.58 (1.5)	-0.61 (-0.99,-0.23)	0.003*
<b>Quads onset (hemiplegic)</b>	1.33 (1.06)	-0.18 (-0.73,0.37)	0.510
<b>Quads onset (non-hemiplegic)</b>	1.44 (1.26)	0.20 (-0.35,0.73)	0.444
<b>Difference between quads peak</b>	0.9 (1.16)	-0.75 (-1.15,-0.35)	0.001*
<b>Difference between quads and ham peak times (hemiplegic)</b>	0.62 (1.03)	-0.75 (-1.24,-0.27)	0.003*
<b>Difference between quads and ham peak times (non-hemiplegic)</b>	0.51 (0.76)	-0.75 (-1.24,-0.27)	0.003*
<b>Time peak quads from seat-off (hemiplegic)</b>	0.14 (1.57)	-0.43 (-1.16,0.30)	0.243
<b>Time peak quads from seat-off (non-hemiplegic)</b>	0.264 (1.384)	-0.46 (-0.9,-0.02)	0.042*
<b>CoM at peak quads (hemiplegic)</b>	202.31 (73.42)	0.00 (-0.01,0.00)	0.350
<b>CoM at peak quads (non-hemiplegic)</b>	211.37 (68.26)	-0.01 (-0.02,0.00)	0.164
<b>Difference between quads onset and peak (hemiplegic)</b>	1.39 (1.50)	0.46 (-0.08,1.01)	0.091
<b>Difference between muscle onset and peak (non-hemiplegic)</b>	1.37 (1.26)	0.77 (0.31,1.23)	0.002*
<b>Time of peak trunk flexion (s)</b>	2.67 (1.28)	-0.67 (-1.08,-0.26)	0.002*

\*= significant at 0.05 level or below