

1           **Distribution of Potentially Toxic Soil Elements along a Transect across Kazakhstan**

2  
3   Mert Guney,<sup>1,5</sup> Almira Yagofarova,<sup>2,5</sup> Vadim Yapiyev,<sup>1,3</sup> Christian Schönbach,<sup>4</sup> Jong R. Kim,<sup>1</sup>  
4   Vassilis J. Inglezakis<sup>2,5,\*</sup>

5  
6   <sup>1</sup> Department of Civil & Environmental Engineering, School of Engineering, Nazarbayev  
7   University, Nur-Sultan, Kazakhstan

8   <sup>2</sup> Department of Chemical & Materials Engineering, Environmental Science & Technology Group  
9   (ESTg), School of Engineering, Nazarbayev University, Nur-Sultan, Kazakhstan

10   <sup>3</sup> Department of Geography and Environmental Science, The University of Reading, Reading,  
11   United Kingdom

12   <sup>4</sup> Department of Biology, School of Science and Technology, Nazarbayev University, Nur-Sultan,  
13   Kazakhstan

14   <sup>5</sup> Environment & Resource Efficiency Cluster (EREC), Nazarbayev University, Nur-Sultan,  
15   Kazakhstan

16  
17       \* Corresponding author: [vasileios.inglezakis@nu.edu.kz](mailto:vasileios.inglezakis@nu.edu.kz), +7 (7172) 70 65 34

18

19

20           **Distribution of Potentially Toxic Soil Elements along a Transect across Kazakhstan**

21

22   **Abstract**

23   The present study aims to investigate the distribution of selected potentially toxic elements (PTEs)  
24   in Kazakhstan's topsoils. Soil samples collected across a north-south gradient (n=40) near main  
25   highways connecting major residential/industrial areas were characterized for their As, Cd, Co, Cr,  
26   Cu, Mn, Ni, Pb, and Zn content as well as for soil physio-chemical properties. The majority of the  
27   soils had neutral pH (no significant relationship between pH and PTE concentrations). The soil  
28   organic carbon was higher at the northern and farther southern parts of the transect (along with  
29   higher concentrations of PTEs in soils). As, Mn, and Ni concentrations in soils were elevated in  
30   comparison to relevant background concentrations. Critical concentrations of As, Cd, Co, Mn, and  
31   Ni (with respect to regulatory limits) were found at multiple locations, with As being particularly  
32   of potential concern (range: 8.7-42 mg × kg<sup>-1</sup>). The distance from the nearest settlement seems to  
33   influence the soil PTE concentrations, however the relationship is not statistically significant. In  
34   total, eight locations had statistically outlier PTE concentrations for Cd, Mn, Ni, and Zn. The  
35   overall results were comparable to similar studies across the world except that the Pb content of  
36   the study soils was less elevated. Studies on site characterization and human health risk assessment  
37   covering identified hotspots and PTEs are recommended.

38

39   **Key words:** arsenic, heavy metals, Kazakhstan environment, site characterization, soil pollution

40

41        **1. Introduction**

42        The Republic of Kazakhstan is situated in Central Asia and is the ninth largest country in the  
43 world in terms of its land area, covering more than 2.7 million km<sup>2</sup>. Kazakhstan has vast quantities  
44 of land resources (mainly minerals, oil, and gas). As a result, since gaining its independency, its  
45 government has mainly depended on the extraction of oil, gas, and minerals as well as the  
46 operation of related industries for the country's economic growth. As a result, Kazakhstan is the  
47 political leader among Central Asian countries with an economy showing the strongest  
48 performance with a continuous growth momentum (Makhmutova, 2018).

49        The current economic growth in Kazakhstan is mainly fueled by the exploitation of its land  
50 resources and accompanying booming sectors of construction and government services, and this  
51 may have negative effects on the environment such as problems related to contamination including  
52 of water and soil (Kuroda *et al.* 2005). Kazakhstan's soils not only show a tendency of being  
53 affected from land degradation, desertification, and salinization due to natural and anthropogenic  
54 factors but also may be under environmental strain due to contamination by heavy metals. This  
55 could negatively affect public health as well as economy via a reduction in land productivity,  
56 stability, functions, and services depending on the exploitation of its natural systems.

57        A limited number of localized studies have investigated environmental problems in different  
58 regions of the country, some also focusing on the potentially toxic elements (PTEs) in soils:  
59 Mueller *et al.* (2014) investigated the level of heavy metals in oil-polluted soils in Atyrau region,  
60 the reported concentrations of measured Pb, Co, Ni, Cd, Cu, Zn, V, and Mo were 2-12 times  
61 higher than the maximum permissible concentrations (MPCs) stated in legislation. The researchers  
62 state that the oil-polluted soil became an accumulator and a storage for hazardous forms of the  
63 heavy metals which could migrate to the Caspian Sea and thus could cause problems of larger

64 scale. Also, they investigated hazardous mobile forms of Cu, Pb, Ni, Zn, and Cd in large rice  
65 irrigation farms in southern Kazakhstan, which showed that heavy metal concentrations impacted  
66 soil fertility in agricultural areas where mobile forms of measured heavy metals accumulated in  
67 soil profile, making hazardous metals potentially available to crop plants and in drinking water.  
68 Ramazanova (2016) investigated Pb, Zn, Cd, Cu, and Fe in soils near central heating power plants  
69 at the different distances (50-5000 m) in Almaty: The levels of Fe and Zn in soils complied with  
70 the MPCs; however, the Cd and Pb concentrations were five to ten times higher than the MPCs.  
71 Bayandinova *et al.* (2018) investigated heavy metals, hydrocarbons, nitrogen and carbon oxides,  
72 and organic compounds in air, water resources, and soil at East Kazakhstan region. Specifically,  
73 Cd, Cu, Pb, and Zn were investigated in soils near smelters at distances from <1 km to 3 km near  
74 Ust-Kamenogorsk, Semey, and Ridder. Elemental concentrations were higher than the MPCs in all  
75 investigated districts of Ust-Kamenogorsk in spring and autumn campaigns, the same was also  
76 observed in Ridder and Semey. Specifically, the concentrations of Pb, Cd, and Zn were noted as  
77 very high one km near the large power plants. Muzychenko *et al.* (2017) studied the Pb pollution  
78 in the roads of Almaty and found high concentrations attributed to traffic as well as the industrial  
79 and geological history of the region. Finally, Mukasheva *et al.* (2013a, 2013b) monitored Cr, Cu,  
80 Cd, Pb, Zn, Ni, V, and Mn pollution near Temirtau, Balkhash, and Karaganda towns. All these  
81 cities are industrial centers with large smelters and plants near or inside the city boundaries such as  
82 Karaganda Steel Mill and chemical plants “Jambul Cement” and “ArcelorMittal” in Temirtau.  
83 Their comparative data analysis showed that the concentration of heavy metals in all soil samples  
84 exceed the MPCs and heavy metals could accumulate in soil and crop plants.

85 Although a number of localized studies already indicate environmental adverse impact from  
86 anthropogenic activities, a systematic study that is reporting soil quality conducted on a larger

87 scale is missing in the literature. Although Kazakhstan is the economical leader of Central Asia  
88 and one of the major developing countries around the world, the environmental consequences of  
89 this expansion has not yet been well studied. As a part of a large study that aims to investigate  
90 Kazakhstan's soils in terms of its physical and chemical properties (Yapiyev *et al.* 2018),  
91 contamination status, and nutrient potential, the present study aims to investigate the distribution of  
92 selected PTEs (As, Cd, Co, Cr, Cu, Mn, Ni, Pb, and Zn) in topsoils of Kazakhstan collected  
93 through a north-south transect (n=40, sampling sites approx. 50 km apart) near highways  
94 connecting numerous major residential/industrial landmarks of the country. We aim to identify  
95 hotspots as well as relationships between individual PTE concentrations and site/soil properties.  
96 Establishing a baseline about the current status is important as this not only help identify potential  
97 problem areas but also could be used to compare with the future monitoring and later to predict the  
98 dynamics of the change.

99

## 100 **2. Materials & Methods**

101

### 102 **2.1. Sampling Area, Sampling and Sample Handling**

103 Soil samples (n = 40) have been collected in September 2016 (Fig. 1) from sampling sites  
104 located a minimum of 50 m away from the nearest road. The sample from the northernmost point  
105 was collected near Petropavlovsk city (5454'N, 6906'E) and the southernmost sample was  
106 collected near Chu river (4255'28''N, 7311'6''E). During sampling, following the removal of the  
107 litter layer if present (approximately the top 2 cm), drilled samples were cored from approximately  
108 0-15 cm. The samples, then, were air-dried in the laboratory, sieved to < 2 mm, and stored at room  
109 temperature until analyzed (see also Methods section in Yapiyev *et al.* (2018)).

110

111 **2.2. Laboratory Analyses**

112 Soil pH was measured in soil suspension using 8107UWMD Ross Ultra pH/ATC triode  
113 electrode (Thermo Fisher Scientific, USA) and an Orion 013010MD conductivity cell electrode  
114 (Thermo Fisher Scientific, USA). Soil organic carbon (SOC) was measured using a C/N dry  
115 combustion elemental analyzer (Multi N/C 3100, Analytic Jena). Samples were pre-treated on the  
116 combustion boat by adding 100  $\mu\text{L}$  of  $\text{H}_3\text{PO}_4$  (30-40% w/w) to 100 mg milled soil to dissolve  
117 carbonates first. The samples were then dried overnight at 70  $^\circ\text{C}$  and finally subjected to  
118 combustion at 950  $^\circ\text{C}$  under  $14 \text{ L} \times \text{min}^{-1}$  oxygen flow (see also Methods section in Yapiyev *et al.*  
119 (2018)).

120 The selected PTEs were measured using ICP-MS iCAP RQ. Sample digestion was first  
121 conducted via a microwave system (Multiwave PRO by Anton Paar). Soil samples were dried for  
122 48 h, crushed, and sifted through 150  $\mu\text{m}$  sieve. Then, 1 g of soil was weighed accurately at 0.1 mg  
123 (Radwag XA52/Y) and then was transferred into a vessel. 9 mL HCl (37% w/w) and 3 mL of  
124  $\text{HNO}_3$  (68-70% w/w) were added separately to the vessel, and samples were digested at 140  $^\circ\text{C}$  for  
125 40 min. After the digestion has been complete, all vessels were cooled, then the digestates were  
126 filtered through Whatman No. 1 or equivalent filter paper (11  $\mu\text{m}$  pore size). After cooling and  
127 filtering, solutions were collected in a volumetric flask and then diluted to 50 mL. The reagent  
128 blanks were carried out in parallel with each experimental analysis, All analyses were conducted in  
129 duplicates. Furthermore, for quality control purposes the reference material BGS 102 was digested  
130 and the concentrations of PTEs were measured using ICP-MS iCAP RQ. The concentrations of  
131 measured Cr, Cu, Ni Pb, Zn were close to certified values (i.e. within 20%), except for Cd where

132 the difference between measured and certificated ( $0.275 \pm 0.182 \text{ mg} \times \text{kg}^{-1}$ ) concentration was  
133 about 40%.

134

### 135 ***2.3. Data Processing and Statistical Analyses***

136 Apart from calculations performed for determining selected descriptive statistics (average,  
137 minimum, maximum, standard deviation, coefficient of variation, skewness, kurtosis), The Pearson  
138 correlation tests have been conducted to check the relationship between (a) individual PTE  
139 concentrations, (b) PTE concentrations and soil physio-chemical properties, and (c) PTE  
140 concentrations and distance of sampling sites from nearest settlement (a settlement defined as a  
141 residential zone with population > 1,000). Sampling points were also categorized into two groups  
142 according to their distance from the nearest settlement; the first including points relatively close to  
143 the nearest settlement (distance < 10 km, n = 20) and the second consisting of points relatively far  
144 from the nearest settlement (distance > 10 km, n = 20) where exact distances are presented in the  
145 Supplementary material (Supplementary Table 1). These groups were compared via two sample *t*-  
146 test in order to see whether there are differences between PTE concentrations. Shapiro-Wilk test  
147 along with Q-Q plots have been employed to check the distribution of data with the specific  
148 purpose of identifying outlier PTE concentrations. Furthermore, a box plot visually representing  
149 the data for potentially toxic elements has been prepared (Fig. 3) via software (IBM SPSS 25).  
150 Outliers have also been determined. which consist of points with concentrations of PTEs larger  
151 than  $1.5 \times \text{IQR}$  (interquartile range, the difference between the third and the first quartiles).

152

## 153 **3. Results and Discussion**

154

155        **3.1. Soil Properties**

156        The pH of the samples (Supplementary Table 1) ranged from 5.32 to 9.18. The majority of the  
157 samples had neutral pH (between 6 and 8) except two samples that can be categorized as acidic  
158 (with pH < 6, sampling locations: S7, S10) and 11 samples that could be classified as alkaline  
159 (with pH > 8: S25-S32, S35, S37-S38). The pH has a significant effect on the mobility of trace  
160 metals in soil such that at a lower pH the sorption capacity of soil is lower than that at neutral or  
161 alkaline pH, Consequently, the mobility of most part of metals decreases from acids soils to  
162 alkaline. Soil samples from the present research are mostly neutral therefore the expected mobility  
163 of metal forms is generally at the lower end. There was also a trend that through the north-south  
164 transect (Fig. 2, based on data from Yapiyev *et al.* (2018)), the soils samples become more alkaline  
165 from north to south which could be linked to more arid climate conditions and dominance of  
166 calcareous soils (Slessarev *et al.* 2016) .

167        The amount of SOC in soil (Supplementary Table 1) influences, among many other things, the  
168 mobility and bioavailability of various elements by binding them into its complex structure and  
169 thus limiting their mobility at higher SOC values. The SOC content of study soils ranged of 0.38%  
170 to 4.09% showing that the study samples collected through the transect were mainly mineral  
171 (inorganic) soils. Through the north-south transect, higher SOC values (> 2%) identifying  
172 potentially organic soils have been observed specifically in the north (S3, S5-S9, S12, S14, S16)  
173 and farther south (S34, S36-S40). These zones are also inside the country's two important arable  
174 regions, therefore, such an observed specific elevated profile of SOC would be expected.

175

176        **3.2. Concentrations of PTEs in Soils**



177 The levels of potentially toxic elements (PTEs) were first compared with the background  
178 concentrations of Canadian and Chinese soils (Tables 1 and 2). The background levels may be  
179 useful in order to estimate the differences between natural content of metals including their  
180 geochemical composition as source and the fraction coming from contamination sources (Santos-  
181 Frances *et al.* 2017). The background concentrations of Canadian (MDDELCC 2019) and Chinese  
182 (Huamain *et al.* 1999) soils were used since in certain parts of these countries, the climate as well  
183 as the conditions that the surface soils had been developed were comparable to those in  
184 Kazakhstan.

185 The comparison with the Canadian soil background levels showed that particularly As  
186 concentrations were above background levels such that all of the samples from the present study  
187 had As concentrations larger than  $6 \text{ mg.kg}^{-1}$ , corresponding to an average As concentration in the  
188 study soils was  $20.8 \text{ mg.kg}^{-1}$ . Considering the other PTEs, Mn and Ni average concentrations were  
189 also higher than corresponding background values (34 of 40 points for Mn and 29 of 40 points for  
190 Ni exceeded the background value). The levels of Co, Cr, Cu, and Zn were comparable to the  
191 Canadian background levels (i.e. with comparable average vales) whereas Pb concentrations were  
192 lower (only five points exceeding the background level) and Cd concentrations were much lower  
193 (all points below the background level) than the Canadian background concentrations.

194 The comparison of PTE concentrations from the present study with Chinese soil background  
195 values (which reported values are lower than the Canadian background values except for As)  
196 showed that the found concentrations were higher for As, Cd, Cr, Cu, Ni, and Zn; and were  
197 comparable for Pb. No element had systematically lower concentrations than the background  
198 values (no background values were present for Co or Mn therefore no comparison could be made).

199 Overall, the comparison of the PTE concentrations in the study soils with the background  
200 concentrations of soils of China and Canada indicated high concentrations of especially As and of  
201 Mn and Ni. The evidence regarding the elevated Cd, Cr, Cu, and Zn concentrations was less  
202 clearly cut.

203 The results from the soils of the present study were also compared with regulatory standards  
204 of Kazakhstan and Russia (which share the same regulatory limits), China, and Canada for soils.  
205 Regarding the regulatory standards of Kazakhstan (MEPRK 2004) and Russia (RPR 2006), for As,  
206 Co, Cr, Ni, and Zn, the measured concentrations exceeded the maximum permissible  
207 concentrations (MPCs) in every single sample. The majority of the samples also exceeded the  
208 criteria for Cu (35 of 40 samples) and Pb (22 of 40 samples) whereas for Mn and Cd, less samples  
209 had concentrations higher than the MPC (19 and 9, respectively, of 40 samples). That being said,  
210 many MPCs presented in Kazakh and Russian regulations seem low or very low even in  
211 comparison to natural background concentrations of these elements as in the cases of: As (MPC:  
212  $2\text{mg} \times \text{kg}^{-1}$ ), Cd ( $0.5\text{mg} \times \text{kg}^{-1}$ ), Co ( $5\text{mg} \times \text{kg}^{-1}$ ), Cr ( $6\text{mg} \times \text{kg}^{-1}$ ), Cu ( $33\text{mg} \times \text{kg}^{-1}$ ), Ni  
213 ( $4\text{mg} \times \text{kg}^{-1}$ ), and Zn ( $23\text{mg} \times \text{kg}^{-1}$ ). Since we could not find any supporting documents and  
214 references that provide a scientific rationale for these regulations it is not possible to evaluate the  
215 scientific soundness of these limits. Consequently, it may be recommended to not to depend on  
216 these comparisons alone for evaluating the levels of PTEs of the study soils.

217 When the results from the present study have been compared to the regulatory standards in  
218 Canada for residential soils (MDDELCC 2019), the only element with systematically high  
219 concentrations in topsoils was Mn (concentrations exceeding the limit at 34 of 40 points). Also,  
220 As, Co, and Ni concentrations were high in five, four, and six points, respectively. This  
221 comparison identified four critical sampling locations with elevated PTE concentrations indicating

222 possible contamination with three or more elements: S8 (with As, Co, Mn, and Ni), S35 (with Co,  
223 Mn, and Ni), S36 (with As, Co, Mn, and Ni), and S39 (with As, Mn, and Ni).

224 A similar comparison with the Chinese environmental quality standards (MEPPRC 1995)  
225 showed that Cd and Ni concentrations were higher than the stated norms at the majority of the  
226 points (24 and 29 points, respectively, out of 40). This is mainly because the Chinese regulation  
227 has stricter limits for Ni and Cd in comparison to the Canadian regulation. Parallel to the above  
228 comparison, the As concentrations were higher than the norm at five locations. This comparison  
229 identified seven sampling locations with elevated PTE concentrations indicating possible  
230 contamination with three or more elements: S7 (with Cd, Cr, and Ni), S8 (with As, Cd, and Ni), S9  
231 (with As, Cd, and Ni), S19 (with As, Cd, Ni, and Zn), S28 (with Cd, Cu, and Ni), S36 (with As,  
232 Cd, and Ni), and S39 (with As, Cd, and Ni). The common locations with the comparison to  
233 Canadian standards were S8, S36, and S39.

234

### 235 ***3.3. Analysis of Concentrations and Potential Hotspots***

236 The descriptive statistics for PTEs in the study soils (Table 2) provided no average values or  
237 ranges at the extreme ends for the selected PTEs. As discussed previously, a comparison of the  
238 elemental averages to Canadian and Chinese environmental norms indicated As and Ni (by both  
239 regulations) as well as Cd, Co, and Ni (according to one of the regulations) as PTEs exceeding  
240 norms at some locations. The coefficient of variation values fell in a normal range except that it  
241 was higher for Pb, possibly due one outlier point with high concentration (S11). The skewness of  
242 the data was to the left for all PTEs, indicating a pooling of values at the lower range while a few  
243 points with higher concentrations extended the distribution tail to the right end, which could be  
244 reasonably expected for the soil PTE concentration data from this site. Finally, the data were

245 leptokurtic for the majority of the elements as expected, with the exception of Cr and Cu having  
246 flatter distribution than the other elements which may indicate anthropogenic disturbances.

247 Inter-element relationships may provide information on the sources and pathways of PTEs.  
248 According to the Pearson correlation coefficients (Table 3), the concentrations in eight of the nine  
249 selected PTEs (with the exception of Pb) are significantly correlated with each other at  $p < 0.01$   
250 level. It is known that the behavior of metals in soils are controlled by a number of key parameters  
251 (USEPA 1992) including the soil's pH and cation exchange capacity (which is mainly defined by  
252 soil organic matter and clay content of the soil) when cations are of concern. As a result, it may be  
253 expected to observe such a correlation between elemental concentrations in a set of soil samples as  
254 certain soil physio-chemical properties control the mobility and retention of a variety of PTEs in a  
255 similar fashion. The fact that Pb has a set of correlations with other PTEs with lower and/or with  
256 less/not significant values may be due to different reasons. In terms of its affinity to soils and soil  
257 constituents, Pb has a high relative order of sorption (USEPA 1996) and consequently it is one of  
258 the least mobile heavy metals in soils. As a result, other PTEs in soil might have exhibited  
259 relatively higher mobilization rates in the past which leading to a reduction in their soil  
260 concentrations with time. Another possibility is that Pb might be introduced to the environment  
261 from overland traffic activities (the samples locations have been accessible sites from nearby  
262 highways) therefore shows a different concentration profile than the other PTEs. This, however, is  
263 unlikely as Pb tends to accumulate mostly within the nearby soils, particularly within the first 10 m  
264 whereas the sampling locations were located a minimum of 50 m away from the nearest road. That  
265 being said, the overland traffic activities introduce to the environment not only Pb but also Cu and  
266 Zn (Guney *et al.* 2010), and in the present study Pb is also correlated with these PTEs at 0.01 level,  
267 therefore, this may require further investigation in the future.

268 The correlation between PTEs concentrations and soil pH & SOC (Table 4) showed that  
269 although the concentrations of the majority of PTEs showed an inverse correlation with pH, the  
270 values were not statistically significant. Therefore, the pH of the soils was not a major parameter  
271 controlling the PTE concentrations in soil. On the contrary, the SOC values were positively and  
272 significantly correlated with PTE concentrations. The soil organic matter significantly contributes  
273 to cationic and anionic exchange capacities, therefore may be expected to increase the retention of  
274 PTEs in soils.

275 For the two categories of points according to their distance from the nearest settlement  
276 (population > 1,000), the points that are relatively closer to the nearest settlement (< 10 km, n =  
277 20) had their average concentrations of the majority of the PTEs (Table 5) higher than that of the  
278 points that are relatively farther (>10.0 km, n=20). However, the two sample *t*-test indicated that  
279 the difference between the concentrations of PTEs was only significant for Cr. Although it first  
280 seems that the settlements have an impact on the topsoil concentrations of PTEs, this impact is not  
281 statistically significant. An analysis of Pearson correlations between the concentrations of  
282 individual PTEs and the distance of the sampling location from the nearest settlement (Table 6)  
283 also indicated negative correlations for the majority of the elements: however, the values were low  
284 and were significant at 0.05 level only for three elements. Therefore, it is not possible to talk about  
285 a significant impact of settlements on the PTE concentrations of topsoils. Such a significant impact  
286 may not have been observed due to the fact that some of the settlements are not large enough to  
287 have a major impact on nearby soils via anthropogenic activities. It is also possible that the impact  
288 of anthropogenic activities may not be significant due to relatively large distances between  
289 settlements and sampling locations.

290 Shapiro-Wilk's normality test has been employed (Table 7) together with Q-Q plots and  
291 histograms with the specific purpose of identifying outliers having excess concentrations of PTEs.  
292 The data showed normal distribution only for As, Cr, Mn, and Zn. An examination of test results  
293 together with Q-Q plots and histograms revealed eight locations with outliers: one location with  
294 three outliers (S36 for Cd, Mn, and Ni), two locations with two outliers (S10 for Cd and Mn, S32  
295 for Cd and Zn), and five locations with one outlier concentration (S7, S9, S11, S19, and S28). A  
296 more detailed discussion is presented below regarding the identification of specific sampling  
297 locations of concern in terms of elevated PTE concentrations. Finally, the box plot analysis  
298 identified similar locations with outlier concentrations: at S10 (Pb), S11 (Pb), S32 (Cd), S35 (Co),  
299 and S36 (Co, Mn, Ni).

300 Based on the previously presented comparison with the background concentrations and the  
301 regulatory limits from selected countries, the PTEs of concern for the present study could be  
302 identified as As, and to a lower extent Cd, Mn, and Ni. In terms of locations, a comparison to  
303 regulatory limits and the statistical analyses yielded varying results (Canada: S8, S35, S36, S39;  
304 China: S7, S8, S9, S19, S28, S36, S39; statistical analyses: S36, S10, S32, S7, S9, S11, S19, S28).  
305 At the end, it is possible to identify five locations as potential hotspots: The location S36 has been  
306 unanimously identified as a potential hotspot. Similarly, S8 and S39 are two other potential  
307 hotspots identified by the comparison to regulatory limits. Finally, it is worth to consider S10 and  
308 S32 as potential hotspots as they are pointed by the statistical analyses having outlier  
309 concentrations at each location. It is interesting to note that none of these three points are near  
310 large settlements or under the direct nearby influence of industrial activities. The elevated  
311 concentrations of some PTEs at these locations might be due to natural sources or might result  
312 from long-distance impact of anthropogenic activities such as smelting. It should be noted that the

313 sampling locations have been selected on a random basis to represent the topsoil concentrations of  
314 selected PTEs and a direct investigation of contamination sources for PTEs was not the major goal.  
315 Therefore, at these five locations (S36, S8, S39, S10, and S32), further investigation and site  
316 characterization involving multiple sampling locations and taking potential human exposure into  
317 account is recommended.

318         Among five points identified above as potential hot spots for their elevated PTE  
319 concentrations by a comparison to regulatory limits and/or via statistical means, their soil types  
320 included chernozem (S8, S10), arenosol (S32), regosol (S36), and umbrisol (S39). Chernozems are  
321 soils rich in organic matter commonly found in the steppes of Eurasia up to Siberia. The SOC for  
322 S8 and S10 were 2.41% and 1.63%, respectively. The elevated concentrations of PTEs in S8 (As,  
323 Co, Cd, Mn, and Ni; according to regulations) and S10 (Cd, Mn, Pb; identified as outliers via  
324 statistical analyses) may be attributed to cation retention mechanisms from organic matter limiting  
325 the mobility of elements. Umbrisols are also steppe environment soils, characterized by a surface  
326 layer that is rich in humus. Similar to S8 and S10, this may explain high concentrations of PTEs in  
327 S39 (As, Cd, Mn, and Ni; according to regulations). Regosols are weakly developed soils, a  
328 characteristic of eroding landscape, and it not easy to comment on their properties due to  
329 unconsolidated material they are formed and a lack of soil horizons. However, regarding the  
330 presence of PTEs, S36 is close to other sampling locations with umbrisol soil type (S37 to S39),  
331 therefore, high concentrations of PTEs in S36 (As, Co, Cd, Mn, and Ni; according to regulations)  
332 would be partly explained by the similarities to the concentrations in neighboring sampling points  
333 (e.g. S39). Arenosols, on the contrary, may be expected to have generally low PTEs due to their  
334 sandy nature accompanied by low humus and clay content. Therefore, outlier Cd and Zn  
335 concentrations at S32 warrant further investigation where possible explanations include

336 contaminated dust fallout from industrial facilities or of geogenic origin e.g. relatively common  
337 sulfide minerals greenockite (CdS) and wurtzite (ZnS).

338

### 339 ***3.4. Comparison to Studies Investigating Similar Sites***

340 Anthropogenic sources affect the PTE concentrations in soils. A study on the general patterns  
341 of element abundances in urban soils pointed out that the overall pattern of elements in Earth's  
342 soils tends to mainly repeat those in the Earth's crust (Alekseenko *et al.* 2014). It also indicated  
343 that certain elements tend to be more abundant in urban soils than the Earth's soils in general,  
344 including some of the PTEs (As, Cd, Co, Cu, Pb, and Zn) subjected in the present work.

345 A comparison of the results from the present study to studies in China (where rural areas with  
346 similar climate to the present study sampling locations and under influence of nearby  
347 anthropogenic activities are common) indicated similar findings. Cheng (2003) reviewed the  
348 contamination by heavy metals (focus on Cd, Cu, Pb, and Zn) in different areas of China including  
349 rural and agricultural areas. The background levels of heavy metals in soils were low, but human  
350 activities have been shown to pollute soil, water, and air where metals may be transferred to plants  
351 and food; three identified main sources of contamination being industrial emissions, wastewater,  
352 and solid waste. The reported heavy metal concentrations were within the range in the present  
353 study, with the exception of Cd being higher and Zn and Cr being also slightly more elevated.  
354 Another review of heavy metal contaminations in urban soils, road dusts, and agricultural soils  
355 from China (Wei *et al.* 2010) showed that among selected elements (Cr, Ni, Cu, Pb, Zn, As, Hg,  
356 and Cd), the geoaccumulation index of urban soils showed Cu, Pb, Zn and Cd enrichment and of  
357 agricultural soils had elevated concentrations of Cd, Hg, and Pb due to anthropogenic activities  
358 with various sources (mining, sewage sludge, pesticides, fertilizers, traffic emission, electroplating



359 plant, spring factory, band steel factory, leather factory, petrochemical complex, etc.) noted as the  
360 main sources of trace metals contamination. The contents of PTEs were in general comparable to  
361 the present research. Liao *et al.* (2007) has performed a geochemical survey of soils of Jiangsu  
362 Province, China for 54 elements in 103,000 top-soil samples and presented similar results  
363 regarding PTE concentrations. They reported that for eight PTEs (As, Cd, Hg, Pb, Ni, Cr, Cu, and  
364 Zn), the measured concentrations of these PTEs were higher than their natural background at 40%  
365 of the samples (with over 10% of the land being classified as contaminated). Finally, Zhao *et al.*  
366 (2010) considered anthropogenic activities as the reason of PTE contamination of Yangtze River  
367 Delta region. They analyzed selected PTEs (Cu, Zn, Pb, Cr, Ni, Cd, and Hg); among these, the  
368 concentrations of Cu, Zn, and Pb significantly exceeded background levels.

369 Other studies of comparable nature to the present study have reported similar results to the  
370 present study with a few caveats. The levels of some PTEs (As, Cd, Co, Cu, Fe, Ni, Pb, Mn, Mg,  
371 and Zn) in soil near smelters in the Sudbury region in Ontario, Canada (a region with comparable  
372 climate and soils to the present study) were monitored (Nkongolo *et al.* 2008). The  
373 concentrations of PTEs were within the limits set by Ontario Ministry of Environment and  
374 Energy (OMEE) guidelines (which are similar to MDDELCC guidelines from Quebec, Canada  
375 used in the present study), even in sites within the vicinity of the Falconbridge Smelters. A  
376 comparison to the concentrations in the present study showed similar PTE levels except for  
377 higher Co, Mn, Ni, and Zn content in the present research, where presumably a strict  
378 enforcement of guidelines in Ontario may have led to such a difference. The concentrations of  
379 PTEs reported by Nezat *et al.* (2017) in urban soils of Spokane, Washington also reported similar  
380 results to the present study with the exception that they reported elevated Pb concentrations which  
381 has been likely due to lead-based paint or vehicle emissions in urban zones. A study conducted on

382 urban soils of Yerevan city, Armenia (Tepanosyan *et al.* 2017) reported similar concentrations of  
383 PTEs and some enrichment in urban zones with the exception of clearly elevated Pb concentrations  
384 attributed to the historical pollution in the study zone. These findings on Pb are also parallel to the  
385 findings of Turner *et al.* (2018) in urban soils of Plymouth, UK who reported elevated Pb due to  
386 contamination by paint. Also, Pb contamination tends to decrease to the background value at about  
387 70 m or more roadside distance (Yan *et al.* 2013). This is in agreement with our findings as the  
388 soils of the present study did not exhibit an extensive contamination by Pb because all samples  
389 were taken from locations with relatively limited impact of transportation (sampling locations at  
390 least 50 m away from the roads) or urban activities (many sampling locations distant from urban  
391 settlements). Still, such an impact might be important for other PTEs as different studies indicate  
392 that both local and distant sources of anthropogenic emissions may still lead to enrichment of PTEs  
393 in non-urban soils e.g. in national park soils in Poland by Pb, Zn, Cu and Mn (Mazurek *et al.* 2017)  
394 and in soils from forests and protected areas in Romania by Pb, Zn, Cu, Cr, and Cd (Ungureanu *et*  
395 *al.* 2017). Finally, a very large soil survey covering the entire Europe and analyzing more than  
396 15,000 surface soil samples have proposed 28.3% of the total surface area of the EU for further  
397 assessment because one or more of the elements (among As, Cd, Cr, Cu, Hg, Pb, Zn, Sb, Co and  
398 Ni) exceed the applied threshold concentration, identifying both natural background and  
399 historical/recent industrial and mining areas as potential sources. Similarly, the present study  
400 identified eight locations with at least outlier elemental concentration as pointed out by the  
401 statistical analyses (corresponding to 20% of 40 sampling locations).

402

#### 403 **4. Conclusions**

404 As a rapidly developing country, Kazakhstan relies on its vast land resources of minerals, oil,  
405 and gas together with related industries, which may create a major burden on its environment. Soil  
406 is an important and now scarce resource and its contamination by potentially toxic elements  
407 (PTEs) may limit its usability for agricultural and urban purposes. The present study investigated  
408 the distribution of selected PTEs (As, Cd, Co, Cr, Cu, Mn, Ni, Pb, and Zn) in topsoils of  
409 Kazakhstan collected through a north-south transect at 40 locations. The analyses have shown that  
410 the soils from the study area were mostly neutral in terms of pH, with a tendency of having more  
411 alkaline samples as moved from north to south. The statistical analyses didn't indicate a significant  
412 relationship between pH and the PTE concentrations. The soil organic carbon was higher at the  
413 northern and farther southern parts of the sampling transect and these locations also had higher  
414 concentrations of PTEs in soils. A comparison of PTEs to background concentrations of soils in  
415 two countries with similar climate and soil properties (China and Canada) indicated elevated  
416 concentrations of As, Mn, and Ni in soils. A comparison to regulatory limits of China, and Canada  
417 also indicated critical concentrations of As, Cd, Co, Mn, and Ni in multiple locations (Some of the  
418 regulatory limits of Kazakhstan and of Russia were very low and the scientific basis for  
419 establishing these limits was not disclosed; thus, conclusions based on these limits are not  
420 emphasized here). Although the distance of the sampling locations from the nearest settlement  
421 (population > 1,000) seemed to influence the PTE concentrations, the relationship was not  
422 statistically significant. Further statistical analyses identified eight locations with outlier PTE  
423 concentrations: S36 for Cd, Mn, and Ni, S10 for Cd and Mn, S32 for Cd and Zn), and five  
424 locations for one element (S7, S9, S11, S19, and S28). Overall, the results of the present study  
425 were comparable to previously conducted studies around the world at locations with similar  
426 properties with the specific remark that the Pb content of soils was less elevated than some other

427 study locations with a history of anthropogenic impact. Studies aiming the site characterization and  
428 human health risk assessment on identified hotspots (particularly at S36 as identified via  
429 comparison to legislations and by statistical investigations) and PTEs (particularly for As that is  
430 elevated at most locations) are recommended.

431

#### 432 **Acknowledgements**

433 The present research was supported in part by the grant “Kazakhstan soil microbiome:  
434 agricultural characteristics and perspectives” in the framework of the agreement #104 dated  
435 12.02.2015 of the program #055 “Scientific and/or Scientific-Technical Activities” within  
436 subprogram #101 “Research Grant Funding” from the Ministry of Education and Science of the  
437 Republic of Kazakhstan and by Nazarbayev University. The authors also acknowledge financial  
438 support from Nazarbayev University Small Grant Program with code SOE2018020.

439

#### 440 **Conflict of Interest:**

441 The authors declare that they have no conflict of interest.

442

443       **References**

- 444       Alekseenko, V., Alekseenko, A. (2014). The abundance of chemical elements in urban soils.  
445               *Journal of Geochemical Exploration*, 147, 245-249.
- 446       Bayandinova, S., Mamutov, Z., Issanova, G. (2018). *Man-Made Ecology of East*  
447               *Kazakhstan*, Springer, Singapore.
- 448       Cheng, S. (2003). Heavy metal pollution in China: origin, pattern and control. *Environmental*  
449               *Science and Pollution Research*, 10, 192 – 198.
- 450       Guney, M., Onay, T.T., Coptu, N.K. (2010). Impact of overland traffic on heavy metal levels in  
451               highway dust and soils of Istanbul, Turkey. *Environmental Monitoring and Assessment*,  
452               164, 101-110.
- 453       Huamain, C., Chunrong, Z., Cong, T., Yongguan, Z. (1999). Heavy Metal Pollution in Soils in  
454               China: Status and Countermeasures. *Ambio*, 2, 130-134.
- 455       Kuroda, H., Eckermann, R., Gauvreau, E., Haryanto, A., Lee Ch.-H. et al. (2005). *The Annual*  
456               *Report of Asian Development Bank*. Asian Development Bank, Singapore.
- 457       Liao, Q-L., Evans, L., Gu, X., Fan, D-F., Jin, Y., Wang, H. (2007). A regional geochemical  
458               survey of soils in Jiangsu Province, China: Preliminary assessment of soil fertility and soil  
459               contamination. *Geoderma*, 142, 18–28.
- 460       Makhmutova, E.V. (2018). Central Asia in Search for Its Own Way of Integration. *Mgimo*  
461               *Review of International Relations*, 4, 78–91.
- 462       Mazurek, R., Kowalska, J., Gasiorek, M., Zadrozny, P., Jozefowska, A. et al. (2017). Assessment  
463               of heavy metals contamination in surface layers of Roztocze National Park forest soils (SE  
464               Poland) by indices of pollution. *Chemosphere*, 168, 839-850.

465 MDDELCC (Ministère du Développement durable, Environnement et Lutte contre les  
466 Changements Climatiques du Québec) (2019). *Annex 2: Tableau 1: Politique de protection*  
467 *des sols et de réhabilitation des terrains contaminés*. Accessed on: 27 April 2019.  
468 <http://www.environnement.gouv.qc.ca/sol/terrains/guide-intervention/annexe2.pdf>.

469 Mueller, L., Saparov, A., Lischeid, G. (2014). *Novel Measurement and Assessment Tools for*  
470 *Monitoring and Management of Land and Water Resources in Agricultural Landscapes of*  
471 *Central Asia*. Springer, Switzerland.

472 Mukasheva, M.A., Mukasheva, G.Z., Nugumanova, S.M., Kazimova, A.E. (2013a). Pollution of  
473 the soil cover of the industrial city with heavy metals. *Journal of Chelyabinsk University*,  
474 298, 152-154.

475 Mukasheva, M.A., Surzhikov, D.V., Tykezhanova, G.M., Nugumanova, S.M., Kazimov, A.E.,  
476 Mukasheva, G.Z. (2013b). Assessment of the soil man-made pollution as the example of an  
477 industrial city. *KarGu Journal*, 69, 77-80.

478 Muzychenko, I., Jamalova, G., Mussina, U., Kazulis, V., Blumberga, D. (2017). Case Study of  
479 Lead Pollution in the Roads of Almaty. *Energy Procedia*, 113, 369-376.

480 Nezat, C. A., Hatch, S.A., Uecker, T. (2017). Heavy metal content in urban residential and park  
481 soils: A case study in Spokane. Washington, USA. *Applied Geochemistry*, 78, 186-193.

482 Nkongolo, K.K., Vaillancourt, A., Dobrzeniecka, S., Mehes, M., Beckett, P. (2008). Metal  
483 content in soil and black spruce (*Picea mariana*) trees in the Sudbury region (Ontario,  
484 Canada): low concentration of arsenic, cadmium, and nickel detected near smelter sources.  
485 *Bulletin of Environmental Contamination and Toxicology*, 80, 107-111.

486 MEPPRC (Ministry of Environmental Protection of the People's Republic of China) (1995).  
487 *Environmental Quality Standard for Soils*. GB 15618-1995. China Environmental Science  
488 Press, Beijing, China.

489 MEPRK (Ministry of Environmental Protection of the Republic of Kazakhstan) (2004). *The*  
490 *norms of maximum permissible concentrations of hazardous substances, organisms and*  
491 *other biological substances polluting the soil*. Consignment Order No. 99 of the Ministry of  
492 Health of the Republic of Kazakhstan and No. 21 of the MEPRK, Astana, Kazakhstan.

493 Ramazanova, R. H. (2011). *The impact of anthropogenic emissions sources on the heavy metals*  
494 *content in the soil and plants at the south and southeast of Kazakhstan*. XI International  
495 Scientific and Practical Conference "Problems and Trends of Modern Society"  
496 International Academy of Science and Higher Education, London, UK.

497 RPR (Russian Federation Rospotrebnadzorb) (2006). *GN 2.1.7.2041-06: the maximum*  
498 *permissible concentrations (MPC) of chemicals in the soil*.

499 Santos-Frances, F., Martinez-Grana, A., Alonso Rojo, P., Garcia Sanchez, A. (2017).  
500 Geochemical Background and Baseline Values Determination and Spatial Distribution of  
501 Heavy Metal Pollution in Soils of the Andes Mountain Range (Cajamarca-Huancavelica,  
502 Peru). *International Journal of Environmental Research and Public Health*, 14, 859-881.

503 Slessarev, E., Lin, Y., Bingham, N., Johnson, J., Dai, Y., Schimel, J. et al. (2016). Water balance  
504 creates a threshold in soil pH at the global scale. *Nature*, 540, 567-569.

505 Tepanosyan, G., Sahakyan, L., Belyaeva, O., Maghakyan, N., Saghatelyan, A. (2017). Human  
506 health risk assessment and riskiest heavy metal origin identification in urban soils of  
507 Yerevan, Armenia. *Chemosphere*, 184, 1230 – 1240.

508 Toth, G., Hermann, T., Szatmari, G., Pasztor, I. (2016). Maps of heavy metals in the soils of the  
509 European Union and proposed priority areas for detailed assessment. *Science of the Total*  
510 *Environment*, 565, 1054-1062.

511 Turner, A., Lewis, M. (2018). Lead and other heavy metals in soils impacted by exterior legacy  
512 paint in residential areas of south west England. *Science of the Total Environment*, 619–  
513 620, 1206–1213.

514 Ungureanu, T., Iancu, G. O., Pintilei, M., Chicos, M. M. (2017). Spatial distribution and  
515 geochemistry of heavy metals in soils: A case study from the NE area of Vaslui county,  
516 Romania. *Journal of Geochemical Exploration*, 176, 20–32.

517 USEPA (United States Environmental Protection Agency) 1992. *Behavior of Metals in Soils*.  
518 USEPA Office of Solid Waste and Emergency Response, EPA/540/S-92/018, Washington,  
519 DC, USA.

520 Wei, B., Yang, L. (2010). A review of heavy metal contaminations in urban soils, urban road  
521 dusts and agricultural soils from China. *Microchemical Journal*, 94, 99–107.

522 Yan, X., Gao, D., Zhang, F., Zeng, C., Xiang, W., Zhang, M. (2013). Relationships between  
523 Heavy Metal Concentrations in Roadside Topsoil and Distance to Road Edge Based on  
524 Field Observations in the Qinghai-Tibet Plateau, China. *International Journal of*  
525 *Environmental Research and Public Health*, 10, 762-775.

526 Yapiyev, V., Gilman, C.P. Kabdullayeva, T., Suleimanova, A., Shagadatova, A. et al. (2018).  
527 Top soil physical and chemical properties in Kazakhstan across a north-south gradient.  
528 *Scientific Data*, 5, 180242.



529 Zhao, Y., Wang, Z., Sun, W., Huang, B., Shi, X. et al. (2010). Spatial interrelations and multi-  
530 scale sources of soil heavy metal variability in a typical urban–rural transition area in  
531 Yangtze River Delta region of China. *Geoderma*, 156, 216–227.  
532  
533

534 **List of Figures**

535

536 Fig. 1. Study area with locations of sampling sites

537 Fig. 2. Soil pH and SOC values (0-15 cm depth) through the north-south transect (based on data  
538 from Yapiyev *et al.* (2018))

539 Fig. 3. Box plot for concentrations of PTEs in soil samples (0-15 cm depth) (boxes: first, second,  
540 and third quartiles; whiskers:  $\pm 1.5 \times$  interquartile range)

541

542 **List of Tables**

543

544 Table 1. Range of concentrations of potentially toxic elements (PTEs) in the soils of present study  
545 (0-15 cm depth), background levels of PTEs in Canadian and Chinese soils, and regulatory  
546 standards for PTEs for Kazakhstan, Russia, Canada, and China ( $\text{mg.kg}^{-1}$ )

547 Table 2. Selected descriptive statistics for PTE concentrations

548 Table 3. Pearson correlation matrix between individual PTEs

549 Table 4. Pearson correlations between PTEs and soil physiochemical characteristics


550 Table 5. Results for two sample *t*-test between two sets of points with distance  $< 10$  km and  $> 10$   
551 km to the nearest settlement

552 Table 6. Pearson correlations between PTEs and distance of sampling sites to nearest settlement


553 Table 7. Results for Shapiro-Wilk test and outlier points with PTE concentrations

554 Supplementary Table 1. Sampling points with data for nearby settlements, concentrations of  
555 selected PTEs (0-15 cm depth), and soil properties (from Yapiyev *et al.* (2018))


556



558 **Fig. 1.** Study area with locations of sampling sites




559



560

561 **Fig. 2.** Soil pH and SOC values (0-15 cm depth) through the north-south transect (based on data  
 562 from Yapiyev *et al.* (2018))

563



564

565 **Fig. 3.** Box plot for concentrations of PTEs in soil samples (0-15 cm depth) (boxes: first, second,  
 566 and third quartiles; whiskers:  $\pm 1.5 \times$  interquartile range)

567

568 **Table 1.** Range of concentrations of potentially toxic elements (PTEs) in the soils of present study  
 569 (0-15 cm depth), background levels of PTEs in Canadian and Chinese soils, and regulatory  
 570 standards for PTEs for Kazakhstan, Russia, Canada, and China (mg.kg<sup>-1</sup>)  
 571

<b>PTE</b>	<b>Range for the present study</b>	<b>Natural background values (Quebec, Canada) (MDDELCC 2019)</b>	<b>Natural background values (China) (Huamain <i>et al.</i> 1999)</b>	<b>National standards for Kazakhstan (MEPRK 2004) and Russia (RPR 2006)</b>	<b>Residential limits (Quebec, Canada) (MDDELCC 2019)</b>	<b>Chinese Environmental Quality Standards (MEPPRC 1995)</b>
As	8.70 - 42.0	6	≤15	2	30	30
Cd	0.12 - 0.85	1.5	≤0.20	0.5	5	0.3
Co	10.8 - 80.6	25	N/A	5	50	N/A
Cr	33.7 - 217	100	≤90	6	250	200
Cu	18.6 - 105	50	≤35	33	100	100
Mn	353 - 3.84E3	1,000	N/A	1,500	1,000	N/A
Ni	23.0 - 159	50	≤40	4	100	50
Pb	8.60 - 161	50	≤35	32	500	300
Zn	23.6 - 264	140	≤100	23	500	250

572 N/A – not available  
 573

574 **Table 2.** Selected descriptive statistics for PTE concentrations  
 575

<b>PTE</b>	<b>Average</b>	<b>Min</b>	<b>Max</b>	<b>Standard deviation</b>	<b>Coefficient of variation</b>	<b>Skewness</b>	<b>Kurtosis</b>
As	20.8	8.70	42.0	7.67	36.9%	0.59	0.007
Cd	0.38	0.12	0.85	0.18	47.9%	0.88	0.529
Co	29.8	10.8	80.6	15.8	53.1%	1.58	2.753
Cr	120	33.7	217	46.9	39.1%	0.28	-0.762
Cu	53.6	18.6	105	20.8	38.7%	0.65	-0.291
Mn	1.68E3	353	3.84E3	762	45.4%	0.83	0.543
Ni	69.6	23.0	159	28.7	41.2%	0.87	0.875
Pb	38.7	8.60	161	23.8	61.6%	3.62	17.883
Zn	140	23.6	264	46.4	33.1%	0.37	0.910

576  
 577

578 **Table 3.** Pearson correlation matrix between individual PTEs  
 579

	<b>As</b>	<b>Cd</b>	<b>Co</b>	<b>Cr</b>	<b>Cu</b>	<b>Mn</b>	<b>Ni</b>	<b>Pb</b>	<b>Zn</b>
<b>As</b>	1	0.481**	0.580**	0.809**	0.699**	0.602**	0.763**	0.307	0.642**
<b>Cd</b>		1	0.449**	0.622**	0.610**	0.623**	0.482**	0.295	0.583**
<b>Co</b>			1	0.699**	0.577**	0.767**	0.859**	0.501**	0.513**
<b>Cr</b>				1	0.594**	0.713**	0.873**	0.321*	0.671**
<b>Cu</b>					1	0.628**	0.623**	0.417**	0.763**
<b>Mn</b>						1	0.745**	0.670**	0.725**
<b>Ni</b>							1	0.293	0.658**
<b>Pb</b>								1	0.424**
<b>Zn</b>									1

580 \* Significant at the 0.05 level

581 \*\* Significant at the 0.01 level

582



583 **Table 4.** Pearson correlations between PTEs and soil physiochemical characteristics  
 584

	<b>As</b>	<b>Cd</b>	<b>Co</b>	<b>Cr</b>	<b>Cu</b>	<b>Mn</b>	<b>Ni</b>	<b>Pb</b>	<b>Zn</b>	<b>pH</b>	<b>SOC</b>
<b>pH</b>	-0.213	0.012	-0.070	-0.302	0.086	-0.139	-0.127	-0.046	-0.107	1	-0.350*
<b>SOC</b>	0.516**	0.170	0.549**	0.564**	0.273	0.510**	0.669**	0.128	0.394*	-0.350*	1

585 \* Significant at the 0.05 level

586 \*\* Significant at the 0.01 level

587

588 **Table 5.** Results for two sample *t*-test between two sets of points with distance < 10 km and > 10  
 589 km to the nearest settlement  
 590

<b>Element</b>	<b>Average concentration for d &lt; 10 km (mg × kg<sup>-1</sup>)</b>	<b>Average concentration for d &gt; 10 km (mg × kg<sup>-1</sup>)</b>	<b><i>p</i>-value</b>	<b><i>t</i>-statistic</b>	<b>Null hypothesis</b>
As	22.5	19.1	0.161	1.432	Accepted
Cd	0.41	0.35	0.314	1.021	Accepted
Co	33.3	26.2	0.158	1.439	Accepted
Cr	135	105	0.044	2.087	Rejected
Cu	53.2	54.1	0.893	-0.134	Accepted
Mn	1.80E+03	1.56E+03	0.334	0.978	Accepted
Ni	76.7	62.5	0.118	1.601	Accepted
Pb	36.6	40.8	0.583	-0.554	Accepted
Zn	141	139	0.889	0.14	Accepted

591  
 592  
 593

594 **Table 6.** Pearson correlations between PTEs and distance of sampling sites to nearest settlement  
595

	<b>As</b>	<b>Cd</b>	<b>Co</b>	<b>Cr</b>	<b>Cu</b>	<b>Mn</b>	<b>Ni</b>	<b>Pb</b>	<b>Zn</b>
<b>Distance</b>	-0.228	-0.257	-0.327	-0.415	0.060	-0.267	-0.364	-0.051	-0.196
<b>r</b>	0.052	0.066	0.107	0.172	0.004**	0.071	0.133	0.003**	0.038*

596 \* Significant at the 0.05 level

597 \*\* Significant at the 0.01 level

598

599 **Table 7.** Results for Shapiro-Wilk test and outlier points with PTE concentrations  
 600

<b>PTE</b>	<b><i>p</i>-value</b>	<b>W</b>	<b>Distribution <sup>a</sup></b>	<b>Outlier points <sup>b</sup> (concentration in mg × kg<sup>-1</sup>)</b>
As	0.150	0.9585	Normal	S9 (42.0)
Cd	0.011	0.9251	Non-normal	S32 (0.85), S10 (0.78), S36 (0.76)
Co	0.000	0.8527	Non-normal	S36 (80.6), S35 (75.5)
Cr	0.246	0.9649	Normal	S7 (217)
Cu	0.050	0.9447	Non-normal	S28 (105)
Mn	0.070	0.9490	Normal	S36 (3.84E+03), S10 (3.30E+03)
Ni	0.046	0.9437	Non-normal	S36 (159)
Pb	0.000	0.6824	Non-normal	S11 (161)
Zn	0.431	0.9725	Normal	S19 (264), S32 (243)

<sup>a</sup> Normality is rejected if  $p < 0.050$

<sup>b</sup> determined via Q-Q plots and histograms

601  
 602

603 **Supplementary Table 1.** Sampling points with data for nearby settlements, concentrations of selected PTEs (0-15 cm depth), and  
 604 soil properties from Yapiyev *et al.* (2018)  
 605

Sample ID	Soil type	Location	Nearest settlement	Distance (km)	Population	Elemental concentration (mg × kg <sup>-1</sup> )										pH (-)	SOC (%)
						As	Cd	Co	Cr	Cu	Mn	Ni	Pb	Zn			
S1	Chernozem	Road street Potanina	Petropavlovsk city	0	201,446	9.66	0.14	23.3	54.7	18.6	642	30.9	14.0	73.2	7.07	0.90	
S2	Chernozem	Highway A 1	Beiterek village	2.7	9,679	13.1	0.14	22.7	67.3	28.8	1.38E3	46.4	26.7	87.9	6.38	1.87	
S3	Chernozem	Highway A 1	Astrakhanka village	3.2	6,313	21.2	0.41	37.6	157	64.5	2.32E3	94.4	38.2	186	6.93	4.09	
S4	Chernozem	Highway A 1	Amangel'dy village	20.3	7,569	17.8	0.31	26.0	128	39.9	1.21E3	72.0	27.7	99.5	7.64	1.82	
S5	Chernozem	Highway A 1	Rostovka village	17.6	2,290	26.8	0.36	43.1	196	67.9	2.34E3	107	41.8	189	7.45	3.06	
S6	Chernozem	Highway A 343	Kokshetau town	35.6	135,106	21.0	0.26	26.1	133	44.9	1.18E3	74.4	26.6	130	7.37	2.81	
S7	Chernozem	Highway A 1	Kokshetau town	1.3	135,106	29.4	0.46	44.6	217	83.4	2.18E3	112	40.0	169	5.76	2.16	
S8	Chernozem	Highway A 1	Kenesary village	1.4	1,598	32.7	0.42	55.4	185	86.9	2.11E3	119	40.0	181	7.24	2.41	
S9	Chernozem	Highway A 1	Shchuchinsk town	6.3	44,106	42.0	0.51	44.9	192	80.1	2.42E3	96.3	39.6	182	6.33	3.07	
S10	Chernozem	Highway A 1, Gas station "Sinooil"	Makinka village	2.3	2,013	23.5	0.78	31.8	200	51.1	3.30E3	73.1	70.9	210	5.32	1.63	
S11	Chernozem	Highway A 1	Akkol village	17.5	14,217	17.4	0.31	54.1	109	55.1	3.00E3	58.8	161	128	7.46	1.40	
S12	Kastanozem	Highway A 1	Bazoygir village	3.3	2,526	26.6	0.29	34.6	131	53.4	2.07E3	87.5	42.9	140	7.53	2.10	
S13	Kastanozem	Highway M 36	Astana city	0	613,006	16.5	0.25	18.4	79.5	38.5	1.09E3	53.4	23.8	109	7.34	1.96	
S14	Kastanozem	Highway M 36	Anar village	29.4	1,117	27.7	0.34	26.5	128	54.2	1.43E3	69.1	30.3	135	7.69	2.19	
S15	Kastanozem	Highway M 36	Anar village	1.4	1,117	24.1	0.40	22.0	112	43.4	1.31E3	60.7	31.8	115	6.51	1.62	
S16	Kastanozem	Highway M 36	Temirtau town	22	169,590	8.9	0.21	15.3	72.6	38.8	982	45.5	31.5	134	7.44	2.36	
S17	Kastanozem	Highway M 36	Karaganda town	3.5	459,778	13.5	0.31	16.4	87.6	37.4	1.15E3	49.2	27.0	132	7.26	2.00	
S18	Calcisol + Solonetz	Highway M 36	Akbastau village	17.6	3,300	18.8	0.29	19.2	92.6	40.6	1.38E3	52.8	30.4	130	6.30	1.34	
S19	Calcisol + Solonetz	Highway M 36	Aksu-Ayuly village	10.9	4,586	32.5	0.49	32.6	157	90.1	1.44E3	86.0	61.8	264	6.91	0.92	
S20	Calcisol + Solonetz	Highway M 36	Aksu-Ayuly village	35.1	4,586	14.0	0.17	14.9	71.5	33.2	966	38.4	25.6	98.7	6.76	1.30	
S21	Arenosol	Highway M 36	Akshatau village	10.4	1,149	13.8	0.26	16.7	80.4	40.4	1.02E3	47.6	20.3	109	6.62	0.98	

S22	Arenosol	Highway M 36	Akzhal village	22.8	3,397	24.5	0.24	16.5	129	32.7	1.16E3	57.2	26.4	110	7.71	0.69
S23	Arenosol + Solonetz	Highway M 36	Balkhash town	55.6	68,883	14.9	0.12	16.3	68.6	36.1	1.07E3	48.4	23.6	92.3	7.99	0.59
S24	Arenosol + Solonetz	Highway M 36	Konirat village	3.1	3,103	13.1	0.73	14.4	116	25.6	353	33.3	8.6	23.6	7.91	0.62
S25	Arenosol + Solonetz	Highway M 36	Balkhash lake, recreation area Gulf Stream, Balkhash town	28.9	68,883	8.70	0.14	10.8	44.8	26.4	854	35.0	13.6	80.9	8.35	0.61
S26	Arenosol	Highway M 36	Balkhash lake, Balkhash town	64.4	68,883	14.4	0.15	17.6	69.4	37.3	1.09E3	44.7	32.1	107	8.40	0.52
S27	Solonchak + Solonetz	Highway M 36	Saryshagan village	15.7	4,429	17.8	0.45	23.4	100	52.6	1.71E3	59.7	39.3	135	8.14	0.67
S28	Arenosol	Highway M 36	Akbakay village	72.7	1,473	24.0	0.52	22.0	90.0	105	1.76E3	51.8	48.9	144	8.53	0.43
S29	Arenosol	Highway M 36	Balkhash lake chemical plant "Jambul Cement"	9.2	721	20.6	0.32	22.8	110	68.6	1.45E3	50.7	48.1	124	9.18	0.45
S30	Arenosol	Highway M 36	Shyganak village	13.9	2,402	16.8	0.40	18.3	83.3	42.7	1.18E3	53.3	36.7	136	8.65	0.38
S31	Arenosol	Highway M 36	Aksuek village	42.1	1,231	11.7	0.36	13.8	33.7	66.3	726	23.0	27.9	127	8.25	0.87
S32	Arenosol	Highway A 358	Birlik village	16.7	3,157	16.8	0.85	30.6	127	85.6	2.88E3	65.7	44.7	243	8.76	0.77
S33	Chemozem + Solonetz	Highway A 358	Kenes village	0.5	2,332	16.2	0.36	33.9	157	44.2	1.91E3	99.1	31.4	161	7.96	1.00
S34	Regosol	Highway A 358	Chu river 200m, Tole Bi village	1.3	19,000	20.4	0.29	24.5	97.7	43.7	1.69E3	63.4	32.9	117	7.35	2.07
S35	Regosol	Highway P 29	Aspara village	5.1	1,086	17.5	0.43	75.5	143	58.2	1.90E3	104	32.7	134	8.18	1.69
S36	Regosol	Highway P 29	Oytal village	13.4	4,181	33.0	0.76	80.6	190	91.4	3.84E3	159	65.3	195	7.78	3.61
S37	Umbrisol	Highway A 2	Zhambul village	4.8	6,633	29.0	0.53	36.4	163	63.5	2.13E3	97.7	46.9	191	8.33	2.48
S38	Umbrisol	Highway A 2	Kokdonen village	0.5	2,063	21.4	0.29	29.6	104	44.9	1.76E3	72.5	36.3	146	8.27	2.46
S39	Umbrisol	Highway A 2	Terenezek village	4.7	2,060	31.4	0.59	37.8	180	68.3	2.06E3	103	48.5	166	7.83	2.15
S40	Arenosol	Highway A 2, Gas station "Sinooil"	Talas village	2.3	2,921	28.0	0.51	39.8	144	60.2	2.71E3	87.8	51.0	181.3	7.70	2.80

