

DRAFT VERSION AUGUST 2, 2021

Typeset using L^AT_EX twocolumn style in AASTeX62

Elemental Abundances of the Hot Atmosphere of Luminous Infrared Galaxy Arp 299

JUNJIE MAO,^{1,2,3} PING ZHOU,⁴ AURORA SIMIONESCU,^{3,5,6} YUANYUAN SU,⁷ YASUSHI FUKAZAWA,^{8,9,10} LIYI GU,^{3,11}
HIROKI AKAMATSU,³ ZHENLIN ZHU,^{3,5} JELLE DE PLAA,³ FRANÇOIS MERNIER,^{12,3} AND JELLE S. KAASTRA^{3,5}¹*Department of Physics, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan*²*Department of Physics, University of Strathclyde, Glasgow G4 0NG, UK*³*SRON Netherlands Institute for Space Research, Sorbonnelaan 2, 3584 CA Utrecht, the Netherlands*⁴*School of Astronomy and Space Science, Nanjing University, Nanjing 210023, PR China*⁵*Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands*⁶*Kavli Institute for the Physics and Mathematics of the Universe (WPI), The University of Tokyo, Kashiwa, Chiba 277-8583, Japan*⁷*Department of Physics and Astronomy, University of Kentucky, 505 Rose Street, Lexington, KY, 40506, USA*⁸*Department of Physical, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan*⁹*Hiroshima Astrophysical Science Center, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan*¹⁰*Core Research for Energetic Universe, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan*¹¹*RIKEN High Energy Astrophysics Laboratory, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan*¹²*European Space Agency (ESA), European Space Research and Technology Centre (ESTEC)**Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands*

ABSTRACT

Hot atmospheres of massive galaxies are enriched with metals. Elemental abundances measured in the X-ray band have been used to study the chemical enrichment of supernova remnants, elliptical galaxies, groups and clusters of galaxies. Here we measure the elemental abundances of the hot atmosphere of luminous infrared galaxy Arp 299 observed with *XMM-Newton*. To measure the abundances in the hot atmosphere, we use a multi-temperature thermal plasma model, which provides a better fit to the Reflection Grating Spectrometer data. The observed Fe/O abundance ratio is subsolar, while those of Ne/O and Mg/O are slightly above solar. Core-collapse supernovae (SNcc) are the dominant metal factory of elements like O, Ne, and Mg. We find some deviations between the observed abundance patterns and theoretical ones from a simple chemical enrichment model. One possible explanation is that massive stars with $M_{\star} \gtrsim 23 - 27 M_{\odot}$ might not explode as SNcc and enrich the hot atmosphere. This is in accordance with the missing massive SNcc progenitors problem, where very massive progenitors $M_{\star} \gtrsim 18 M_{\odot}$ of SNcc have not been clearly detected. It is also possible that theoretical SNcc nucleosynthesis yields of Mg/O yields are underestimated.

Keywords: X-rays: galaxies – galaxies: individual: Arp 299 – galaxies: abundance – techniques: spectroscopic

1. INTRODUCTION

Hot atmospheres of massive galaxies are enriched with metals (Werner et al. 2019). Metals are mainly produced by stars before and during their splendid death as supernovae (SN). Generally speaking, core-collapse supernovae (SNcc) of massive ($M_{\star} \gtrsim 8 M_{\odot}$) stars are the main metal factory of elements like O, Ne, and Mg (Nomoto et al. 2013). Type-Ia supernovae (SNIa) dom-

inate the enrichment of the Fe-peak elements like Fe and Ni (Hitomi Collaboration et al. 2017). Low- and intermediate-massive stars ($M_{\star} \lesssim 7 M_{\odot}$) in the asymptotic giant branch (AGB) phase contribute most to light elements like N (Kobayashi et al. 2006; Werner et al. 2006; Mao et al. 2019).

In chemical enrichment models, it is common to assume that massive stars from $\sim 10 M_{\odot}$ to $\gtrsim 40 - 50 M_{\odot}$ can explode as SNcc and enrich the surrounding environment (e.g., Kobayashi et al. 2006; Sanders & Fabian 2006; de Plaa et al. 2007; Mernier et al. 2016; Mao et al. 2019). If the SNcc progenitor does not contain metals in its atmosphere (i.e. $Z_{\text{init}} = 0$), very massive stars up

Corresponding author: Junjie Mao
jmao2018@hiroshima-u.ac.jp

to $M_* = 140 M_\odot$ can also contribute to the chemical enrichment via SNcc (Nomoto et al. 2013). It is still debated whether very massive stars with higher initial metallicity could make successful SN explosions (Heger et al. 2003).

From the observational perspective, there is a missing massive SNcc progenitor problem as reviewed by Smartt (2015). Massive stars, especially those with $M_* \gtrsim 18 M_\odot$ might explode as a hypernova (HN), explode as a faint supernova ejecting a small amount of heavy elements, or directly form a black hole without a visible supernova (Nomoto et al. 2013). If the bulk of massive stars above a certain mass limit do not explode as SNcc or HN, we expect to see mismatches between the observed and theoretical abundance ratios.

Elemental abundances measured in the X-ray band have been used to decode the chemical enrichment of supernova remnants (e.g., Zhou & Vink 2018), elliptical galaxies, groups and clusters of galaxies (e.g., Hitomi Collaboration et al. 2017; Mernier et al. 2018; Simionescu et al. 2019; Mao et al. 2019). Here, we present the elemental abundances of the hot atmosphere of a starburst galaxy, which form stars intensively.

Arp 299, also known as NGC 3690¹, is a pair of interacting galaxies with a star-formation rate of $119 M_\odot \text{ yr}^{-1}$ (Smith et al. 2018) at the distance of 48.5 ± 3.4 Mpc (James et al. 2014). It is one of the most powerful nearby starburst galaxies (Alonso-Herrero et al. 2009). With $L_{\text{IR}} = 6.3 \times 10^{14} L_\odot$ (Pereira-Santaella et al. 2011), it is qualified as a luminous infrared galaxy (Sanders & Mirabel 1996). From 1992 to 2010, seven SNe have been recorded in Arp 299 (Anderson et al. 2011). Six of them are identified as SNcc and one is unclassified (Anderson et al. 2011). According to the Open Supernova Catalog (Guillochon et al. 2017)², three more SNe have been found since 2010: SN2018lrd, SN2019lqo, and SN2020fkb. All of these three are SNcc. Nevertheless, due to the obscuration by large amounts of dust in Arp 299, Mattila et al. (2012) argue that $\sim 80\%$ of the SNe are estimated to be missed by observations in the optical band.

Arp 299 hosts a hot atmosphere that dominates the soft X-ray spectrum below 2 keV (e.g., Anastasopoulou et al. 2016). About 20 ultraluminous X-ray sources (Zezas et al. 2003; Anastasopoulou et al. 2016) and at least one Compton-thick nucleus (Ptak et al. 2015) dominate the hard X-ray spectrum at 2 – 10 keV and > 10 keV, respectively. Previous X-ray studies do not focus

Table 1. *XMM-Newton* observation log.

Date	ObsID	Effective exposures
2001-05-06	0112810101	9.6 ks (RGS), 13.0 ks (pn)
2011-12-15	0679381101	9.8 ks (RGS), 6.4 ks (pn)
2020-05-08	0861250101	42.9 ks (RGS), 39.3 ks (pn)
2020-05-22	0861250201	34.9 ks (RGS), 26.3 ks (pn)
2020-11-22	0861250301	35.7 ks (RGS), 11.6 ks (pn)

on accurate abundance measurement of O, Ne, and Mg, which are mainly enriched by SNcc. Here, we measure the elemental abundances of the hot atmosphere of Arp 299 with the high-quality X-ray spectra obtained with *XMM-Newton*. We compare the observed abundance ratios with various chemical enrichment models. These models include different SN yields, the initial mass functions (IMFs), the initial metallicities and the mass ranges of the SNcc progenitors, and mixing fractions of HN or SNIa.

A supplementary package is available at Zenodo³. This package includes data and scripts used to create the figures presented in this paper.

2. XMM-NEWTON OBSERVATIONS

We use all the X-ray spectra observed with *XMM-Newton* (Table 1), including three new observations obtained in Cycle 19 (PI: J. Mao). The data are reduced following the standard procedure using *XMM-Newton* Science Analysis Software (SAS) v19.0. We construct a time-averaged X-ray spectrum with the high-resolution soft X-ray spectra (7.5 – 27 Å) obtained with Reflection Grating Spectrometer (RGS, den Herder et al. 2001) and the relatively low-resolution hard X-ray spectra (1.6 – 10.5 Å) obtained with the positive-negative (pn) junction CCD (charge-coupled device) camera of the European Photon Imaging Camera (EPIC).

Lightcurves from the CCD9 of RGS and the 10 – 12 keV energy band of EPIC/pn are created with a time bin of 100 s. Time intervals contaminated by background soft-proton flares are identified as those above the 3σ level and excluded from the following data reduction. As shown in the left panel of Figure 1, we extract RGS spectra in rectangular regions centered on the source with the width along the cross-dispersion direction equals ~ 48 arcsec (i.e., 90 % of the point spread function). We extract EPIC/pn spectra in a circular region with a radius of 24 arcsec. The RGS modeled

¹ IC 694 is a small elliptical or lenticular galaxy about an arcmin northwest, which is not part of Arp 299.

² <https://sne.space/>

³ <https://doi.org/10.5281/zenodo.5148020>




Figure 1. X-ray to IR image of Arp 299. *Left:* *XMM-Newton* soft X-ray (0.3 – 2 keV) image with the RGS (rectangles) and EPIC/pn (circle) source regions. The three new observations in 2020 (Table 1) have similar RGS roll angles so that one rectangular box appears thick. *Right:* The true color image of Arp 299 with soft X-ray in purple, optical (*Hubble*, 814 nm) in green, and infrared (*Spitzer*, 3.6 μm) in red.

background and pn local background spectra are subtracted. All the RGS spectra are combined, as well as the EPIC/pn spectra. By matching the flux in the common 7.5 – 10.5 \AA wavelength range, the EPIC/pn spectra are scaled by 0.70 with respect to RGS to account for the different instrument normalization and the different aperture (the left panel of Figure 1). The right panel of Figure 1 shows a true color image of Arp 299. The soft X-ray (0.3 – 2 keV) image (in purple) is created with the *XMM-Newton image*⁴ script. The optical image (in green) is taken with the Advanced Camera for Surveys Wide Field Channel (F814W filter, 814 nm) aboard *Hubble Space Telescope*. The infrared image (in red) is taken with the Infrared Array Camera (Channel 1, 3.6 μm) aboard *Spitzer* Space Telescope.

3. SPECTRAL ANALYSIS

For the spectral analysis, we use the SPEX code (Kaastra et al. 2020, v3.06.00), which includes the most recent atomic data for the Fe-L complex (Gu et al. 2020). We use *C*-statistics (Kaastra 2017) and statistical uncertainties are quoted at the 68% confidence level. With $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$, and $\Omega_\Lambda = 0.7$, the redshift of the target is 0.0112 and a distance scale of 12.7 kpc/arcmin.

The hot atmosphere is modeled with multi-temperature thermal plasma models. A power-law component (de-

noted as PL) is included to account for the point sources. Three Fe K lines with rest-frame energies of 6.4 keV, 6.7 keV, and 6.97 keV are also included. These Fe K lines are associated with the nucleus of NGC 3690, X-ray binaries and supernova remnants (Ptak et al. 2015; Anastasopoulou et al. 2016). Since the power-law component dominates the flux above $\gtrsim 2$ keV, the absolute abundance of the thermal plasma with respect to hydrogen cannot be well determined. We set the reference element to oxygen. The proto-solar abundance table of Lodders et al. (2009) is used⁵. Both the thermal and power-law components are corrected for absorptions of both the host galaxy and Milky Way. For the Galactic absorption, we use $N_{\text{H}}^{\text{MW}} = 9.35 \times 10^{19} \text{ cm}^{-2}$ (Willington et al. 2013). Similar to Smith et al. (2018), we use different hydrogen column densities for the thermal and power-law components for the host galaxy absorption.

We ran a total of eight spectral fittings considering (1) various differential emission measure distributions, (2) mimicking the non-equilibrium scenario, (3) varying power-law photon indices, and (4) coupling the Fe and Ni abundances. Detailed results are presented in Appendix A. In Figure 2, we show the best-fit model to the observed X-ray spectrum of Arp 299. Most of the

⁵ The abundances denoted as X and X/O designate $(X/H)_{\text{gas}}/(X/H)_{\odot}$ and $(X/O)_{\text{gas}}/(X/O)_{\odot}$, respectively. They correspond to the ratios of X/H and X/O in the hot atmosphere to those in the proto-solar abundance table of Lodders et al. (2009).

⁴ <https://www.cosmos.esa.int/web/xmm-newton/images>




Figure 2. The best-fit model (Model M4 in Table 2) to the X-ray spectrum of Arp 299 observed with *XMM-Newton*. The high-resolution RGS spectrum is shown in red, while the EPIC/pn spectrum is shown in purple. The 1σ uncertainties are shown in grey. The RGS spectrum is rebinned here for clarity. Vertical dashed lines in blue mark key diagnostics emission lines in the spectrum.

previous studies (Huo et al. 2004; Anastasopoulou et al. 2016; Smith et al. 2018) described the hot atmosphere as a single-temperature thermal (i.e., isothermal) component. Our best-fit results favor a multi-temperature description of the hot atmosphere. This should better reflect the multi-temperature nature of the hot atmospheres of starburst galaxies (Strickland & Stevens 2000).

4. METAL ABUNDANCES

Accurate abundance measurement of individual elements is essential to understand the chemical enrichment process of the hot atmosphere of Arp 299. In the present work, we measure the abundance ratios of $N/O = 0.70_{-0.29}^{+0.38}$, $Ne/O = 1.16 \pm 0.21$, $Mg/O = 1.27 \pm 0.26$, $Si/O = 0.97 \pm 0.23$, $S/O = 1.78 \pm 0.57$, $Fe/O = 0.40_{-0.05}^{+0.06}$, and $Ni/O = 0.66_{-0.27}^{+0.31}$. Our measurement takes advantage of the resolving power of the RGS spectra for emission lines of N, O, and Fe-L. Previous studies either do not measure individual elements separately or only cover a few elements. Huo et al. (2004) measured merely the Fe abundance ($0.12_{-0.05}^{+0.21}$) of the thermal plasma model MEKAL (Mewe et al. 1995), which does not contain the state-of-the-art atomic data. Anastasopoulou et al. (2016) used VAPEC and measure the abundances of Ne ($1.24_{-0.29}^{+0.34}$), Mg ($1.14_{-0.20}^{+0.25}$), and Fe ($0.26_{-0.03}^{+0.04}$). Smith

et al. (2018) used VMEKAL and measured merely the $\alpha/Fe (= 3.25 \pm 2.48)$ abundance ratio.

We also note that the abundance ratios measured in the X-ray band are more robust than the absolute abundance with respect to hydrogen, especially for relatively cool thermal plasmas with its weak bremsstrahlung (free-free) emission dominated by the power-law component of the host galaxy. The absolute oxygen abundance (O/H) of the H II regions that are closest to the seven SNe exploded in 1992 – 2010 ranges from ~ 0.53 to ~ 0.70 solar (Anderson et al. 2011). The absolute oxygen abundance (O/H) of the interstellar medium of NGC 3690 is ~ 0.66 solar (Heckman et al. 2015).

In other well studied X-ray bright starburst galaxies, super-solar Ne/O and Mg/O have been reported. For NGC 253, the Mg/O and Ne/O abundance ratios are $\sim 2 - 3$ times solar for the central and surrounding regions (Bauer et al. 2007). For M82, while the stellar abundances are almost solar, the hot atmosphere Mg/O and Ne/O abundance ratios are $\sim 2 - 5$ times solar (Origlia et al. 2004). After taking into account the charge-exchange effect (Zhang et al. 2014), these two abundance ratios are still about twice solar with $Ne/O = 2.2 \pm 0.3$ and $Mg/O = 2.1 \pm 0.3$, respectively.

5. METAL ENRICHMENT VIA SUPERNOVAE

To interpret the observed abundance ratios, we build a simplified chemical enrichment model based on SNcc and SNIa nucleosynthesis yields. SNIa is included here to take into account the potential contribution of a “prompt” population of SNIa (Maoz et al. 2014) that explode within a few hundreds of million years after the peak star-formation activity. The theoretical abundance ratios are calculated via

$$\frac{z_i}{z_j} = \frac{\bar{y}_i^{\text{SNcc}} + r^{\text{SNIa}} y_i^{\text{SNIa}} A_j n_j}{\bar{y}_j^{\text{SNcc}} + r^{\text{SNIa}} y_j^{\text{SNIa}} A_i n_i}, \quad (1)$$

where $z_{i,j}$ are the abundance of the i th or j th element, \bar{y}_i^{SNcc} the IMF-weighted SNcc yields, r^{SNIa} the number ratio of SNIa with respect to SNcc, y_i^{SNIa} the SNIa yields, $A_{i,j}$ the atomic weight of the i th or j th element and $n_{i,j}$ are the elemental abundance by number in the (proto-)solar abundance table.

We do not have accurate abundances of Fe-peak abundances (e.g., Cr, Mn, and Ni) other than Fe. Therefore, we cannot break degeneracy of dozens of SNIa yields available in the literature. For simplicity, we adopt the widely used W7 yields of Iwamoto et al. (1999) when investigating the contribution from a “prompt” population of SNIa.

The IMF weighted SNcc yields⁶ are calculated via:

$$\bar{y}_i^{\text{SNcc}} = \frac{\int_{m_{\text{lo}}}^{m_{\text{up}}} \phi(m) y_i(m) dm}{\int_{m_{\text{lo}}}^{m_{\text{up}}} \phi(m) dm}, \quad (2)$$

where $\phi(m) \propto m^\Gamma$ is the IMF, m_{lo} and m_{up} are the lower and upper mass limits of the progenitors, which depends on its initial metallicity. The Salpeter IMF (Salpeter 1955, $\Gamma = -2.35$) is adopted here as the default. SNcc yields are sourced from Nomoto et al. (2013) and Sukhbold et al. (2016). Nomoto et al. (2013, N13) provide SNcc yields of massive stars with different initial metallicities ($Z = 0, 0.001, 0.004, 0.008, 0.02, \text{ and } 0.05$). The mass range of the progenitor are $11 - 140 M_\odot$ for $Z = 0$ and $13 - 40 M_\odot$ for $Z > 0$. Taking advantage of a one dimensional neutrino transport model for the explosion, Sukhbold et al. (2016, S16) provide SNcc yields of massive ($12.25 - 120 M_\odot$) stars with solar metallicity ($Z = 0.02$). Both the W18 and N20 models of Sukhbold et al. (2016) are considered here. The former was calibrated so that a Nomoto & Hashimoto (1988) progenitor explodes like SN 1987A, while the latter was calibrated so that a Utrobin et al. (2015) progenitor explodes like SN 1987A.

As shown in Figure 3, the calculated IMF weighted abundance ratios are not within the $\gtrsim 1\sigma$ uncertainties

of the observed ones. The closest match is the one with N13 yields for massive ($13 - 40 M_\odot$) stars with solar metallicity ($Z = 0.02$). In the following, we tried to vary several underlying parameters based on the N13 $Z = 0.02$ model.

First, we tried to vary the IMF power-law index. For Arp 299, the relatively high frequency of “stripped-envelope” SNe (type Ib and type I Ib) with respect to normal type II SNe suggests that the IMF of the host galaxy is biased toward the production of high-mass stars (Anderson et al. 2011)⁷. In Figure 3, the purple filled circles illustrate the effect of varying IMF power-law index. A top-heavy IMF with $\Gamma = -2.0$ can decrease the abundance ratios (X/O) by $\lesssim 0.06$, while a bottom-heavy IMF with $\Gamma = -2.7$ can increase the abundance ratios (X/O) by $\lesssim 0.06$. Both cases do not improve the mismatch.

Varying the upper mass limit of the massive progenitors can bring the calculated abundance ratios closer to the observed ones. The blue curve and filled circles in Figure 3 illustrate the effect of varying m_{up} in Eq. 2. The observed abundance ratios N/O, Ne/O, Mg/O, S/O, and Ni/O can be better explained if massive progenitors up to $\sim 23 - 27 M_\odot$ explode as SNcc and enrich the hot atmosphere with metals. The Si/O–Mg/O and Fe/O–Mg/O patterns favor $m_{\text{up}} \gtrsim 30 M_\odot$ with the observed Mg/O ratio larger than the theoretical ones. If the theoretical SNcc nucleosynthesis yields of Mg/O yields are underestimated (Griffith et al. 2021), say by 20%, a more consistent match might be found for all six abundance patterns. In this case, the upper limit of the SNcc progenitors can be larger than $\sim 23 - 27 M_\odot$.

We further consider a more complicated scenario where massive progenitors with $M_\star \sim 13 - 20 M_\odot$ explode as SNcc while a fraction of massive progenitors with $M_\star \sim 20 - 40 M_\odot$ explode as HN⁸. The pink curve and filled circles in Figure 3 illustrate this mixing effect. The abundance pattern of Ne/O–Mg/O can break the degeneracy between the HN fraction and m_{up} cut-off. Taking into account that SNcc are the dominant metal factory of O, Ne, and Mg (i.e., contributions from AGBs and SNIa are negligible), the Ne/O–Mg/O pattern is the best one to explore the difference of theoretical chemical enrichment models of SNcc.

We also consider an alternative complicated scenario where massive progenitors with a mixture of SNcc (from massive stars with $M_\star \sim 13 - 40 M_\odot$) and SNIa. Here

⁷ Note that “stripped-envelope” SNe can also be produced by binary stars with a less massive progenitor (Smartt 2009).

⁸ Nomoto et al. (2013) HN yields are available for $M_\star = 20, 25, 30, \text{ and } 40 M_\odot$ with solar metallicity.

⁶ Only single-star SN nucleosynthesis yields are considered here.




Figure 3. Elemental abundance patterns of the hot atmosphere of Arp 299. The observed abundance ratios (with 1σ uncertainties) are shown as the black crosses. Chemical enrichment models with core-collapse supernovae (SNcc) yields from Sukhbold et al. (2016, S16) and Nomoto et al. (2013, N13) are shown as squares and hexagons, respectively. For the N13 models, we further consider different IMF power-law indices -2.0 (left) and -2.7 (right); different cut-off masses for the SNcc progenitors with the upper mass limit M_{up} : $36 - 39 M_{\odot}$ (leftmost), $30 - 35 M_{\odot}$, $28 - 29 M_{\odot}$, $25 - 27 M_{\odot}$, $23 - 24 M_{\odot}$, and $20 - 22 M_{\odot}$ (rightmost); different hypernova fractions from 0.5 (leftmost) to 0.0 (rightmost) for massive stars with $M_{\star} \sim 20 - 40 M_{\odot}$ (and SNcc for stars with $M_{\star} \sim 13 - 20 M_{\odot}$), and different SNIa fractions from 0.0 (bottom) to 0.4 (top).

we consider the number of SNIa to that of SNcc (r_{SNIa} in Eq. 2) ranges from 0.0 to 0.4 . The green curve and filled squares in Figure 3 illustrate this mixing effect. The abundance patterns of N/O–Mg/O and Ne/O–Mg/O are barely affected by the SNIa fraction. As shown in Mernier et al. (2016); Simionescu et al. (2019); Mao et al. (2019), even for rather old systems like elliptical galaxies, groups and clusters of galaxies, where a significant fraction of SNIa has contributed to the chemical enrichment process, SNcc is still the dominant metal factory of O, Ne, and Mg. For heavier elements including Si, the increasing SNIa fraction will increase the abundance ratios to O.

In short, we find a hint of deviation between the observed abundance patterns and theoretical ones from a simple chemical enrichment model. The deviation can be explained if massive stars with $M_{\star} \gtrsim 23 - 27 M_{\odot}$ do not explode as SNcc. This is in accordance with the

missing massive SNcc progenitors problem, where SNcc observed so far are not produced by massive stars with $M_{\star} \gtrsim 18 M_{\odot}$ (Smartt 2015). It is also possible that theoretical SNcc nucleosynthesis yields of Mg/O yields are underestimated (Griffith et al. 2021). Statistical uncertainties of the current data are still large, which can be significantly improved with the next generation of X-ray spectrometer like XRISM/Resolve (XRISM Science Team 2020) and Athena/X-IFU (Barret et al. 2018). Last but not least, we caution that the observed abundance ratios might not be directly compared to our simple chemical enrichment models. The metals produced by SNcc are distributed not only in the X-ray emitting hot atmosphere but also in the cooler phase of the interstellar medium (gas and dust). Inflows and outflows can further complicate the distribution and dilution of the metals.

This work is based on observations obtained with XMM-Newton, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA. We thank the referee for the careful reading of the manuscript and useful suggestions to improve the quality of this work. JM acknowledges useful discussions with Jiangtao Li and H. Yamaguchi. PZ acknowledges the support from NSFC grant 11590781. AS is supported by the Women In Science Excel (WISE) programme of the Netherlands Organisation for Scientific Research (NWO), and acknowledges the World Pre-

mier Research Center Initiative (WPI) and the Kavli IPMU for the continued hospitality. YS acknowledges support from Chandra Grants AR8-19020A and GO1-22126X. SRON is supported financially by NWO, the Netherlands Organization for Scientific Research.

Facilities: XMM, HST, Spitzer

Software: SPEX v3.06.00 (Kaastra et al. 2020)

APPENDIX

A. SPECTRAL MODELS

A total of eight spectral fittings were performed in the present work. The first three sets (S, T, M0 in Table 2) in Table 2 adopt different differential emission measure distributions for the hot atmosphere: single-temperature (Model S), two-temperature (Model T), and a multi-temperature (Model M1) model with a Gaussian (log-normal) differential emission measure (DEM) distribution (GDEM, de Plaa et al. 2006),

$$Y(x) = \frac{Y_0}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-x_0)^2}{2\sigma^2}\right), \quad (\text{A1})$$

where Y and Y_0 are the emission measures, $x = \log_{10} T$, $x_0 = \log_{10}(T_0)$, and T_0 is peak temperature of the DEM distribution in units of keV. The single-temperature (i.e., isothermal) model is nested in Eq. A1 with $\sigma = 0$. Both the multi-temperature and two-temperature DEM distributions yield significantly better statistics.

Adding another single-temperature cool component to Model M1 can further improve C -statistics by ~ -16 at the cost of 2 less degrees of freedom (Model M2). The main improvement comes from O VII. Setting $\sigma(\log T)$ of this cool component free does not improve the fit at all with $\sigma(\log T)$ remains at zero. With still some residuals for the O VII lines, we set the temperature ratio T_b/T_e for the cool component (Model M3). For collisionally ionized equilibrium plasmas, the ionization balance temperature T_b equals the electron temperature T_e . A non-unity value can mimic non-equilibrium plasmas. For Model M3, we obtain $T_b/T_e \sim 1.6$, where the forbidden to resonant line ratio of He-like O VII triplet was boosted to better match the observed data. But C -statistics is merely improved by ~ -0.8 with one less degree of freedom. This indicates a minor contribution from non-equilibrium plasmas. The charge exchange process can also increase the the forbidden to resonant line ratio of He-like O VII triplet, as in M82 (Zhang et al. 2014). Considering the availability of H I and molecular gas in Arp 299 (e.g., Heckman et al. 1999; Sliwa et al. 2012), it is possible that charge exchange between the hot and cold gas play a role here. Unfortunately, the quality of the current data set is not sufficient to constrain the free parameters of the possible charge exchange process, such as the interacting atomic or molecular species, collision velocities, and different capture mechanisms to the l -subshells. In addition, based on Model M2, we decouple the turbulence velocity (v_{RMS}) of the two thermal components in Model M4. This yields the best C -statics among all eight models.

For all previous models (S to M4), the photon index of the power-law component was fixed to 1.67 as measured by Anastasopoulou et al. (2016) with joint *Chandra* and *NuSTAR* observations. In Model M5, we set the photon index free, which yields $\Gamma \sim 1.59$ but C -statistics is merely improved by ~ -0.8 with one less degree of freedom (compared to Model M2).

Last but not least, based on Model M2, we couple the Ni and Fe abundances in Model M6. This is driven by the imperfect Ni-L complex atomic data in SPEX v3.06.00. The best-fit results of Models M2 to M6 also provide an estimate of the systematic uncertainties, which are smaller than the statistical uncertainties of the current data. That is to say, the key abundance ratios (Ne/O, Mg/O, and Fe/O) do not vary significantly in these models.

REFERENCES

- Alonso-Herrero, A., García-Marín, M., Monreal-Ibero, A., et al. 2009, *A&A*, 506, 1541.
doi:10.1051/0004-6361/200911813
- Anastasopoulou, K., Zezas, A., Ballo, L., et al. 2016, *MNRAS*, 460, 3570. doi:10.1093/mnras/stw1200

Table 2. Best-fit parameters and statistics. The hot atmosphere is modeled with either a single-temperature (Model S), two-temperature (Model T), or multi-temperature (Models M1–M6) differential emission measure plasma model. The power-law (PL) component and Fe K lines are required for the point-sources. The expected C -statistics (Kaastra 2017) are 1135 ± 49 (total), 1010 (RGS), and 125 (pn). Statistical uncertainties at the 68% confidence level are provided for the best-fit (Model M4). Frozen and coupled parameters are shown as (f) and (c), respectively.

Model	S	T	M1	M2	M3	M4	M5	M6
Hot atmosphere								
N_{H}^{HA} (10^{21} cm $^{-2}$)	0.9	1.2	1.3	1.5	1.7	$1.6^{+0.5}_{-0.3}$	1.6	1.5
N_{hot} (10^{64} cm $^{-3}$)	1.7	1.5	2.1	2.2	2.4	$2.4^{+0.8}_{-0.5}$	2.4	2.3
T_{hot}	0.59	0.62	0.61	0.67	0.66	0.67 ± 0.04	0.68	0.68
$\sigma_{\text{hot}}(\log T)$	0.50	0.33	0.35	0.31 ± 0.05	0.36	0.32
$v_{\text{RMS, hot}}$	830	930	810	840	840	840 ± 60	830	840
N_{cool} (10^{64} cm $^{-3}$)	...	0.3	...	0.8	3.3	$0.6^{+2.1}_{-0.3}$	0.9	0.9
T_{cool}	...	0.18	...	0.11	0.086	0.12 ± 0.04	0.11	0.11
$T_{\text{b}}/T_{\text{e}}(\text{cool})$...	1.0 (f)	...	1.0 (f)	1.6	1.0 (f)	1.0 (f)	1.0 (f)
$v_{\text{RMS, cool}}$...	930 (c)	...	840 (c)	840 (c)	380^{+250}_{-190}	830 (c)	840 (c)
N/O	0.85	0.58	0.87	0.69	0.53	$0.70^{+0.38}_{-0.29}$	0.70	0.68
Ne/O	1.21	1.47	1.44	1.22	1.20	1.16 ± 0.21	1.20	1.21
Mg/O	1.00	1.21	1.81	1.36	1.35	1.27 ± 0.26	1.34	1.29
Si/O	1.07	1.17	1.42	1.04	1.02	0.97 ± 0.23	1.05	0.96
S/O	3.8	3.9	2.2	1.89	1.77	1.78 ± 0.57	1.84	1.73
Fe/O	0.24	0.31	0.53	0.42	0.42	$0.40^{+0.06}_{-0.05}$	0.43	0.41
Ni/O	0.97	1.06	1.28	0.73	0.80	$0.66^{+0.31}_{-0.27}$	0.64	0.41 (c)
Point sources								
N_{H}^{PS} (10^{21} cm $^{-2}$)	4.0	4.0	5.1	6.3	6.8	$6.7^{+3.2}_{-1.5}$	5.7	6.6
N_{PL} (10^{49} ph s $^{-1}$ keV $^{-1}$)	7.87	7.85	7.17	8.09	7.43	7.48 ± 0.02	6.56	7.43
Γ_{PL}	1.67 (f)	1.67 (f)	1.67 (f)	1.67 (f)	1.67 (f)	1.67 (f)	1.59	1.67 (f)
$N_{6.4 \text{ keV}}$ (10^{47} ph s $^{-1}$)	11.8	11.8	12.6	12.4	12.5	12.43 ± 0.14	12.1	12.5
$N_{6.7 \text{ keV}}$ (10^{47} ph s $^{-1}$)	7.1	7.1	7.2	7.3	7.3	7.3 ± 1.3	7.1	7.4
$N_{7.0 \text{ keV}}$ (10^{47} ph s $^{-1}$)	2.1	2.2	2.9	2.6	2.7	2.6 ± 1.1	2.4	2.7
Statistics								
C_{stat} (total)	1420.0	1353.2	1311.5	1295.8	1294.6	1292.8	1294.6	1297.1
d.o.f. (total)	1080	1078	1079	1077	1076	1076	1076	1078
C_{stat} (RGS)	1268.4	1203.7	1171.7	1157.7	1156.7	1155.1	1158.0	1157.6
C_{stat} (pn)	151.6	149.4	139.8	138.1	137.9	137.7	136.6	139.5

- Anderson, J. P., Haberman, S. M., & James, P. A. 2011, *MNRAS*, 416, 567. doi:10.1111/j.1365-2966.2011.19080.x
- Barret, D., Lam Trong, T., den Herder, J.-W., et al. 2018, *Proc. SPIE*, 10699, 106991G. doi:10.1117/12.2312409
- Bauer, M., Pietsch, W., Trinchieri, G., et al. 2007, *A&A*, 467, 979. doi:10.1051/0004-6361:20066340
- de Plaa, J., Werner, N., Bleeker, J. A. M., et al. 2007, *A&A*, 465, 345. doi:10.1051/0004-6361:20066382
- de Plaa, J., Werner, N., Bykov, A. M., et al. 2006, *A&A*, 452, 397. doi:10.1051/0004-6361:20053864
- den Herder, J. W., Brinkman, A. C., Kahn, S. M., et al. 2001, *A&A*, 365, L7. doi:10.1051/0004-6361:20000058
- Griffith, E. J., Sukhbold, T., Weinberg, D. H., et al. 2021, arXiv:2103.09837
- Gu, L., Shah, C., Mao, J., et al. 2020, *A&A*, 641, A93. doi:10.1051/0004-6361/202037948
- Guillochon, J., Parrent, J., Kelley, L. Z., et al. 2017, *ApJ*, 835, 64. doi:10.3847/1538-4357/835/1/64
- Heckman, T. M., Armus, L., Weaver, K. A., et al. 1999, *ApJ*, 517, 130. doi:10.1086/307193
- Heckman, T. M., Alexandroff, R. M., Borthakur, S., et al. 2015, *ApJ*, 809, 147. doi:10.1088/0004-637X/809/2/147
- Heger, A., Fryer, C. L., Woosley, S. E., et al. 2003, *ApJ*, 591, 288. doi:10.1086/375341
- Hitomi Collaboration, Aharonian, F., Akamatsu, H., et al. 2017, *Nature*, 551, 478. doi:10.1038/nature24301
- Huo, Z. Y., Xia, X. Y., Xue, S. J., et al. 2004, *ApJ*, 611, 208. doi:10.1086/422164
- Iwamoto, K., Brachwitz, F., Nomoto, K., et al. 1999, *ApJS*, 125, 439. doi:10.1086/313278
- James, B. L., Aloisi, A., Heckman, T., et al. 2014, *ApJ*, 795, 109. doi:10.1088/0004-637X/795/2/109
- Kaastra, J. S. 2017, *A&A*, 605, A51. doi:10.1051/0004-6361/201629319
- Kaastra, J. S., Raassen, A. J. J., de Plaa, J., et al. 2020, *Zenodo*, doi:10.5281/zenodo.3939056
- Kobayashi, C., Umeda, H., Nomoto, K., et al. 2006, *ApJ*, 653, 1145. doi:10.1086/508914
- Lodders, K., Palme, H., & Gail, H.-P. 2009, *Landolt Börnstein*, 4B, 712. doi:10.1007/978-3-540-88055-4_34
- Mao, J., de Plaa, J., Kaastra, J. S., et al. 2019, *A&A*, 621, A9. doi:10.1051/0004-6361/201730931
- Maoz, D., Mannucci, F., & Nelemans, G. 2014, *ARA&A*, 52, 107. doi:10.1146/annurev-astro-082812-141031
- Mattila, S., Dahlen, T., Efstathiou, A., et al. 2012, *ApJ*, 756, 111. doi:10.1088/0004-637X/756/2/111
- Mernier, F., de Plaa, J., Pinto, C., et al. 2016, *A&A*, 595, A126. doi:10.1051/0004-6361/201628765
- Mernier, F., Biffi, V., Yamaguchi, H., et al. 2018, *SSRv*, 214, 129. doi:10.1007/s11214-018-0565-7
- Mewe, R., Kaastra, J. S., & Liedahl, D. A., 1995, *Legacy*, 6, 16.
- Nomoto, K. & Hashimoto, M. 1988, *PhR*, 163, 13. doi:10.1016/0370-1573(88)90032-4
- Nomoto, K., Kobayashi, C., & Tominaga, N. 2013, *ARA&A*, 51, 457. doi:10.1146/annurev-astro-082812-140956
- Origlia, L., Ranalli, P., Comastri, A., et al. 2004, *ApJ*, 606, 862. doi:10.1086/383018
- Pereira-Santaella, M., Alonso-Herrero, A., Santos-Lleo, M., et al. 2011, *A&A*, 535, A93. doi:10.1051/0004-6361/201117420
- Ptak, A., Hornschemeier, A., Zezas, A., et al. 2015, *ApJ*, 800, 104. doi:10.1088/0004-637X/800/2/104
- Salpeter, E. E. 1955, *ApJ*, 121, 161. doi:10.1086/145971
- Sanders, D. B. & Mirabel, I. F. 1996, *ARA&A*, 34, 749. doi:10.1146/annurev.astro.34.1.749
- Sanders, J. S. & Fabian, A. C. 2006, *MNRAS*, 371, 1483. doi:10.1111/j.1365-2966.2006.10779.x
- Simionescu, A., Nakashima, S., Yamaguchi, H., et al. 2019, *MNRAS*, 483, 1701. doi:10.1093/mnras/sty3220
- Sliwa, K., Wilson, C. D., Petitpas, G. R., et al. 2012, *ApJ*, 753, 46. doi:10.1088/0004-637X/753/1/46
- Smartt, S. J. 2009, *ARA&A*, 47, 63. doi:10.1146/annurev-astro-082708-101737
- Smartt, S. J. 2015, *PASA*, 32, e016. doi:10.1017/pasa.2015.17
- Smith, B. J., Campbell, K., Struck, C., et al. 2018, *AJ*, 155, 81. doi:10.3847/1538-3881/aaa1a6
- Strickland, D. K. & Stevens, I. R. 2000, *MNRAS*, 314, 511. doi:10.1046/j.1365-8711.2000.03391.x
- Sukhbold, T., Ertl, T., Woosley, S. E., et al. 2016, *ApJ*, 821, 38. doi:10.3847/0004-637X/821/1/38
- Utrobin, V. P., Wongwathanarat, A., Janka, H.-T., et al. 2015, *A&A*, 581, A40. doi:10.1051/0004-6361/201425513
- Werner, N., Böhringer, H., Kaastra, J. S., et al. 2006, *A&A*, 459, 353. doi:10.1051/0004-6361:20065678
- Werner, N., McNamara, B. R., Churazov, E., et al. 2019, *SSRv*, 215, 5. doi:10.1007/s11214-018-0571-9
- Willingale, R., Starling, R. L. C., Beardmore, A. P., et al. 2013, *MNRAS*, 431, 394. doi:10.1093/mnras/stt175
- XRISM Science Team 2020, arXiv:2003.04962
- Zezas, A., Ward, M. J., & Murray, S. S. 2003, *ApJL*, 594, L31. doi:10.1086/378144
- Zhang, S., Wang, Q. D., Ji, L., et al. 2014, *ApJ*, 794, 61. doi:10.1088/0004-637X/794/1/61
- Zhou, P. & Vink, J. 2018, *A&A*, 615, A150. doi:10.1051/0004-6361/201731583