

Synthesis and Reactivity of Bis-tris(pyrazolyl)borate Lanthanide/Aluminum Heterobimetallic Trihydride Complexes

Tajrian Chowdhury, Fáinché Murphy, Alan R. Kennedy, Claire Wilson, Joy H. Farnaby,* and Catherine E. Weetman*

Cite This: *Inorg. Chem.* 2024, 63, 9390–9394

Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Molecular heterobimetallic hydride complexes of lanthanide (Ln) and main-group (MG) metals exhibit chemical properties unique from their monometallic counterparts and are highly reactive species, making their synthesis and isolation challenging. Herein, molecular Ln/Al heterobimetallic trihydrides $[\text{Ln}(\text{Tp})_2(\mu\text{-H})_2\text{Al}(\text{H})(\text{N}'')]]$ [**2-Ln**; Ln = Y, Sm, Dy, Yb; Tp = hydrotris(1-pyrazolyl)borate; $\text{N}'' = \text{N}(\text{SiMe}_3)_2$] have been synthesized by facile insertion of aminoalane $[\text{Me}_3\text{N}\cdot\text{AlH}_3]$ into the Ln–N amide bonds of $[\text{Ln}(\text{Tp})_2(\text{N}'')]]$ (**1-Ln**). Thus, this is a simple synthetic strategy to access a range of Ln/Al hydrides. Reactivity studies demonstrate that **2-Ln** is a heterobimetallic hydride, with evidence for the cooperative nature of **2-Ln** shown by the catalytic amine–borane dehydrocoupling under ambient conditions in contrast to its monomeric counterparts.

Heterobimetallic hydride complexes are desirable synthetic targets because they are highly reactive species and exhibit unique chemical reactivity properties distinct from their monometallic counterparts.¹ However, their high reactivity makes the synthesis and isolation of heterobimetallic hydrides challenging. This is particularly true in the case of lanthanide (Ln)/aluminum (Al) hydrides, and as such careful ligand design is required to stabilize the highly electrostatic Ln and Al cations, which are bridged by small hydrides. Bis-substituted cyclopentadienyl (Cp^R) ligands have enabled the synthesis of Ln/Al heterobimetallic hydrides predominantly by salt metathesis routes, e.g., $[\{\text{Ln}(\text{Cp}^R)_2(\text{X})\}_n]$ (X = halide) with LiAlH_4 .^{1a} However, the hydride and Ln/Al heterobimetallic hydride chemistry in Trofimenko's isolable scorpionate nitrogen-donor tris(pyrazolyl)borate (Tp^R) ligand environment is much less well established.²

Ln homometallic hydrides $[\{\text{Ln}(\text{Tp}^R)(\text{H})_2\}_n]$ supported by a single (Tp^R) ligand have been isolated, in which the size of the substituents in the 3 and 5 positions of the Tp ligand dictates the nuclearities of the complex (R = ⁱPr, $n = 3$; R = Me, $n = 4$; R = H, $n = 6$).^{2c} Notably only for $[\{\text{Ln}(\text{Tp}^{\text{Me}_2})\text{-}(\text{H})_2\}_4]$ was Ln larger than Y, and for $\text{Tp}^R = \text{Tp}^{3\text{-}t\text{Bu},5\text{-}Me}$, no Ln(III) complexes were isolable. More importantly, $\text{Tp}^{3\text{-}t\text{Bu},5\text{-}Me}$ enabled the isolation of the first molecular ytterbium(II) hydride (**[I]**, Figure 1). These Tp^R -supported Ln hydrides are highly reactive; complex **[I]** undergoes facile activation chemistry at room temperature,^{2a} while **[II-Y]** mediated the hydrogenation and coupling of carbon monoxide (Figure 1).^{2b} The $\text{Tp}^{3\text{-}t\text{Bu},5\text{-}Me}$ ligand environment can also support monomeric Ln/Al heterobimetallic dihydrides $[\text{Ln}(\text{Tp}^{3\text{-}t\text{Bu},5\text{-}Me})(\mu\text{-HAlMe}_3)_2]$, where Ln = Y and Lu (**[III]**, Figure 1).²ⁱ Complex **[III]** was synthesized by the reaction of $[\text{Ln}(\text{Tp}^{3\text{-}t\text{Bu},5\text{-}Me})(\text{Me})_2]$ with 2 equiv of HAlMe_2 and shown to deprotonate amines to access a rare imide-bridged Ln–Al hydride (**[IV]**, Figure 1).^{2i,3} The insertion of substituted alanes into Ln–Me σ -bonds has been shown to be a viable route to




Figure 1. Reactive Ln hydride and Ln/Al heterobimetallic hydride complexes supported by Tp^R and Cp^R ligands.

Ln/Al heterobimetallic dihydrides in the bis- Cp^* ($\text{Cp}^* = \text{C}_5\text{Me}_5$) ligand environment. The reaction of $\text{HAl}(\text{N}'')_2$ ($\text{N}'' = \text{N}(\text{SiMe}_3)_2$) with $[(\text{Cp}^*)_2\text{Y}(\text{Me})(\text{THF})]$ yielded $[(\text{Cp}^*)_2\text{Y}(\mu\text{-H})_2\text{Al}(\text{N}'')(\text{Me})]$ (**[V]**), which was the first example of a Ln catalyst in the hydroalumination of 1-octene.⁴

Using Tp^R , the Ln scope is limited to Y and Lu, and there are no bis-Tp-stabilized heterobimetallic Ln/Al complexes directly analogous to the isolable bis- Cp^R complexes. Recently, monomeric bis-Tp complexes $[\text{Ln}(\text{Tp})_2(\text{X})]$ (Ln = Y, Dy, Yb; X = triflate)⁵ have shown themselves to be a versatile tool for accessing new Ln–element bonds (E = O,^{5a} N^{Sb}) because the anion can be readily exchanged via metathesis or protonolysis. Herein, we report the facile synthesis of molecular Ln/Al


Special Issue: Ligand-Metal Complementarity in Rare Earth and Actinide Chemistry

Received: March 4, 2024

Revised: April 23, 2024

Accepted: April 23, 2024


Published: April 29, 2024



heterobimetallic trihydrides in the bis-Tp ligand environment, which is applicable to a wide range of Ln.

We began by extending the scope of the $[\text{Ln}(\text{Tp})_2(\text{X})]$ complexes to include Sm(III), allowing for a representative range of Ln(III) ions to be studied. Following the same method as previously reported,⁵ $[\text{Sm}(\text{Tp})_2(\text{N}'')]$ (**1-Sm**) was obtained by metathesis between the heteroleptic samarium (III) triflate $[\text{Sm}(\text{Tp})_2(\text{OTf})]$ ($\text{OTf} = \text{CF}_3\text{SO}_3$) with $\text{K}(\text{N}'')$ in toluene. To target Ln/Al hydride species, we opted to use Lewis base-stabilized alanes, $[\text{L}\cdot\text{AlH}_3]$ ($\text{L} = \text{NMe}_3$, N-heterocyclic carbene) because these should undergo insertion chemistry into the Ln–N'' bond akin to the formation of **[III]** and **[V]**.^{2i,4} The reactions of IDippAlH_3 [$\text{IDipp} = 1,3\text{-bis}(2,6\text{-diisopropylphenyl})\text{imidazol-2-ylidene}$] with **1-Ln** ($\text{Ln} = \text{Y, Dy, Yb}$) were slow and resulted in complicated mixtures. Major products were identified as $[\text{Ln}(\text{Tp})_3]$ ^{5b,6} and $[(\text{IDipp})\text{Al}(\text{N}'')(\text{H})_2]$ (**Al-IDipp**; sections A4.1 and B4.4).⁷ In contrast, the reactions of **1-Ln** ($\text{Ln} = \text{Y, Sm, Dy, Yb}$) with aminoalane $[\text{Me}_3\text{N}\cdot\text{AlH}_3]$ in toluene resulted in the evolution of NMe_3 gas and the formation of Ln/Al heterobimetallic trihydride complexes $[\text{Ln}(\text{Tp})_2(\mu\text{-H})_2\text{Al}(\text{H})(\text{N}'')]$ (**2-Ln**; $\text{Ln} = \text{Y, 95\%}$; Sm, 50\% ; Dy, 92\% ; Yb, 94\% ; Scheme 1a).

Scheme 1. (a) Synthesis of Ln/Al Heterobimetallic Trihydride Complexes 2-Ln and (b) Substrate Insertion Reactivity of 2-Ln with Carbodiimides and Benzophenone



Multinuclear NMR data for the diamagnetic **2-Y** and paramagnetic **2-Ln** ($\text{Ln} = \text{Sm, Dy, Yb}$) complexes can be readily assigned by relative integration (Table S1) and are comparable to the literature.⁵ Amide (N'') transfer from Ln to Al is confirmed by ^1H and $^{29}\text{Si}\{^1\text{H}\}$ INEPT NMR spectroscopy because the N'' shifts downfield in line with transfer to the more electronegative element [e.g., 0.13 and -11.96 ppm (**1-Y**) to 0.25 and -2.38 ppm (**2-Y**), respectively]. The metal hydrides of **2-Ln** are observed at 5.07 ppm (**2-Y**), 0.24 ppm (**2-Sm**), and -92.98 ppm (**2-Yb**) in the ^1H NMR spectrum, with no hydride resonances observed for **2-Dy**. Variable-temperature (VT) NMR studies of **2-Y** (Figure S23) maintained the magnetically equivalent hydride environment across the temperature range of 220–353 K, suggesting fast exchange of the hydrides in solution. The Fourier transform infrared (FT-IR) spectra of **2-Ln** showed weak absorptions at 2450–2470 cm^{-1} , which are characteristic of the Tp ligand borohydride stretching frequencies, and weak absorptions between 1800 and 1810 cm^{-1} and between 1730 and 1740 cm^{-1} , consistent with the terminal and bridging hydrides, respectively, reported for Ln/Al heterobimetallic hydrides.^{1a}

Colorless single crystals of **2-Ln** suitable for single-crystal X-ray diffraction (SCXRD) were grown from either saturated hexane solutions ($\text{Ln} = \text{Y, Dy}$) or saturated toluene solutions in the presence of hexane antisolvent ($\text{Ln} = \text{Yb}$) at -35 °C. The structures of **2-Ln** are all monomeric with an 8-coordinate Ln(III) ion bound to two κ^3 -coordinated Tp ligands and two bridging hydrides, which are arranged around the Ln(III) ion in a distorted square-antiprismatic geometry [**2-Ln** = Dy (Figure 2), Y, Yb, and Sm (Figures S110–S112); see section




Figure 2. Molecular structure of **2-Dy**. Hydrogen atoms, except for H, HA, and HB, are omitted for clarity, and the pyrazolyl carbon atoms of Tp are displayed in wireframe. Displacement ellipsoids are drawn at 50% probability.

B4 for structural comparisons]. Two hydrides bridge the Ln(III) ion to a tetrahedral Al(III) ion, which is also bound to a terminal hydride and the N'' anion. Bridging and terminal hydrides were found and refined freely, with the Ln–H–Al bond distances comparable to previously reported Ln/Al heterobimetallic hydrides.^{1a} The Ln...Al separation is similar across the **2-Ln** series [**2-Dy** 3.1826(6), **2-Y** 3.1822(5), and **2-Yb** 3.1439(9) Å]. The close Ln–Al separation (approximately the sum of the Ln and Al covalent radii) is indicative of strong Ln–($\mu\text{-H}$)–Al bonding, consistent with previous bridging hydrides and three-center, two-electron bonds.⁸ The closest structural comparator of **2-Ln** is **[V]**, which exhibits the same μ^2 binding mode of the bridging hydrides.⁴ Notably, **2-Dy** is the first crystallographically characterized example of the Dy–($\mu\text{-H}$)–Al unit. The Ln–N($\kappa^3\text{-Tp}$) bond distances in **2-Ln** are consistent with those of **1-Ln** and $[\text{Ln}(\text{Tp})_2]^+$ in the literature.⁵ The Al–N(N'') bond distances of 1.8430(19)–1.844(3) Å in **2-Ln** are similar to those in **[V]** [1.837(2) Å],⁴ TM/Al heterobimetallic hydrides containing TM–($\mu\text{-H}$)₂–Al(N'')_n⁹ and the homoleptic complex $[\text{Al}(\text{N}'')_3]$ [1.813(3) Å].¹⁰

The reactivity scope of **2-Ln** was investigated to understand the nature of the heterobimetallic hydride. The reaction of **2-Ln** with 1 equiv of **IDipp** again led to the formation of **Al-IDipp**⁷ and multiple products. Importantly, no discrete Ln homometallic hydrides, putative “[$\{\text{Ln}(\text{Tp})_2(\mu\text{-H})_2\}_2$]”, were observed spectroscopically (section B2.2). Previously, the reaction chemistry of $\text{Ln}(\text{L})_2(\mu\text{-X})_2\text{AlX}_2$ ($\text{L} = \text{ligand}$; $\text{X} = \text{monoanionic substituent, e.g., H, CH}_3$) showed that these complexes are best described as an ion pair, $[\text{Ln}(\text{L})_2]^+[\text{AlX}_4]^-$, because they undergo facile metathesis reactions.¹¹ As such, we examined the potential ion-pair character of **2-Ln**. The reaction of **2-Ln** with $\text{K}(\text{N}'')$ should yield the starting material **1-Ln**; however, this resulted in complicated reaction mixtures containing **1-Ln**, $[\text{Ln}(\text{Tp})_3]$,^{5b,6} $[\{\text{K}(\mu\text{-N}'')(\mu\text{-H})\text{Al}(\text{H})_2\}_n]$,¹²

HN^{''}, and H₂ gas and, in addition for **2-Yb**, reduction to [Yb(Tp)₂]¹³ (sections B2.3–B2.7). Complexes **2-Ln** do not undergo facile ion exchange, indicating that the strong Ln–(μ-H)–Al interaction persists in solution (consistent with the VT NMR data). Thus, **2-Ln** should be considered a bimetallic hydride.

To prove this hypothesis, we studied the insertion chemistry of **2-Ln** toward unsaturated substrates. Amides **1-Ln** do not react with carbodiimides to yield the putative guanidates “[Ln(Tp)₂{κ²-(R)NC(N'')N(R)}]”. The reaction of RN=C=NR (R = ⁱPr or Cy) with **2-Ln** in benzene-*d*₆ (Scheme 1b) resulted in formation of the formamidinate complexes [Ln(Tp)₂{κ²-(R)NCHN(R)}] (**3-Ln**) and [Al{κ²-(R)NCHN(R)}₂(N'')] (**3-Al**) (sections B2.10–B2.13). The ¹H NMR resonances of **3-Al** are comparable to those of [Al{κ²-(R)NCHN(R)}₂(X)] (X = hydride, halide) in the literature.¹⁴ Preparative-scale reactions allowed for the separation of **3-Al**, and complexes **3-Ln** (R = Cy) were isolated in moderate yields (Y, 45%; Sm, 36%; Dy, 51%; Yb, 49%; Scheme 1b). Multinuclear NMR data for diamagnetic **3-Y** and paramagnetic **3-Ln** (Ln = Sm, Dy, or Yb) complexes can be readily assigned by relative integration. The FT-IR spectra of **3-Ln** (Ln = Dy, Yb) confirm the transfer of the hydride to the carbodiimide because strong absorptions between 1556 and 1559 cm⁻¹ are assigned to the N–C(H)–N linkage stretching frequencies (ν_{NCN}) in **3-Ln**. ν_{NCN} in **3-Ln** is lower in wavenumber compared to ν_{NCN}, 1650–1670 cm⁻¹, as observed for homoleptic [Ln(N,N'-R₂-formamidinate)₃]¹⁵ and heteroleptic [Ln(N,N'-R₂-formamidinate)₂(X)] complexes, with X monoanions and aromatic R groups.¹⁶

Yellow crystals of **3-Ln** (Ln = Dy, Yb) suitable for SCXRD were grown from saturated toluene solutions in the presence of a hexane antisolvent at –35 °C. Both **3-Yb** (Figure 3) and **3-**




Figure 3. Molecular structure of **3-Yb**. Hydrogen atoms except H10 are omitted for clarity, and the pyrazolyl carbon atoms of Tp are displayed in wireframe. Displacement ellipsoids are drawn at 50% probability.

Dy (Figure S113) structures are monomeric, containing an 8-coordinate Ln(III) ion bound to two κ³-coordinated Tp ligands and a κ²-coordinated bidentate (Cy)NCHN(Cy) anion, arranged around the Ln(III) ion in a distorted square-antiprismatic geometry. The reduction of the carbodiimide was confirmed by the presence of H10 on the sp² carbon (C10) in the now notably bent formamidinate fragment [N7–C10–N7' 118.6(15)° (**3-Dy**) and 119.8(6)° (**3-Yb**)]. A comparison of the C–N bond lengths also confirms the reduced nature of the formamidinate fragment because C10–N7 falls between the typical values for C–N single and double bonds [C10–N7 1.326(14) Å (**3-Dy**) and 1.315(5) Å (**3-Yb**) vs single (~1.47

Å) and double (~1.15 Å) C–N bonds], consistent with delocalization of the N–C–N fragment in **3-Ln**.¹⁷ The Ln–N(κ³-Tp) bond distances in **3-Ln** are consistent with those of **1-Ln**, **2-Ln**, and [Ln(Tp)₂(X)] in the literature.^{5b,6} The Ln–N7 bond distances in **3-Ln** [Ln = Dy, 2.376(8) Å; Yb, 2.373(3) Å] are consistent with the Ln–N bond distances in related literature complexes.^{15,16,18}

Notably, **2-Ln** behaves differently from the TM/Al heterobimetallic trihydride complex [Ti(Cp)₂(μ-H)₂Al(H){C–(SiMe₃)₃}] because reaction with excess ⁱPrN=C=NⁱPr resulted in a formamidinate-bridged Ti/Al complex, [Ti–(Cp)₂(μ-H){μ-(ⁱPr)N(CH)N(ⁱPr)}Al(H){C(SiMe₃)₃}].¹⁹ Analogous to the insertion chemistry of carbodiimides, the insertion of sterically hindered carbonyls like ketones is common in both Ln–H and MG–H chemistry.^{1b,d,h,12b} NMR-scale reactions of **2-Ln** with 3 equiv or more of benzophenone in benzene-*d*₆ yielded the major products [Ln(Tp)₂(OCHPh₂)] and dimeric [{Al(N'')(H)(μ-O–(CHPh₂))₂}] (**4-Al**) (see sections B2.14–B2.17).

Inspired by the Ln/Al heterobimetallic hydride hydroalumination catalyst [V] (Figure 1),⁴ the applicability of **2-Ln** in catalysis was examined to establish the cooperative behavior between the two metal centers. The catalytic dehydrocoupling of [Me₂HN·BH₃] has attracted significant interest due to hydrogen storage applications and as precursors to boron nitride materials.²⁰ Ln and Al hydrides have both been reported to catalyze the dehydrocoupling of [Me₂HN·BH₃] at elevated temperatures, thus providing a good basis for this preliminary study.^{1e,21} The dehydrocoupling of [Me₂HN·BH₃] by 10 mol % **2-Ln** (Ln = Y, Sm, Dy, Yb) proceeds at room temperature to yield [Me₂N·BH₂]₂ as the major product (see section C). Notable differences were found across the **2-Ln** series, with consumption of amine–borane occurring most efficiently in the order Yb > Y ~ Dy > Sm, in line with the relative Lewis acidity of the Ln(III) ion. It is of note that **1-Y** and [Me₃N·AlH₃] do not catalyze this reaction under these conditions; therefore, cooperativity between the two metal centers enables this transformation to occur at room temperature.

In conclusion, the facile synthesis and characterization of Ln/Al heterobimetallic trihydride complexes **2-Ln** (Ln = Y, Sm, Dy, Yb) are reported. The reactivity of **2-Ln** was investigated to understand the cooperative nature of the two different metal centers. Attempts to break apart **2-Ln** by the addition of Lewis bases or ion-exchange chemistry resulted in multiple products. However, upon reaction with unsaturated organic substrates, clean conversion to discrete Ln(III) and Al(III) hydride insertion products was obtained. Highlighting this is the fact that **2-Ln** is best described as a bimetallic hydride with a strong Ln–(μ-H)–Al interaction. Preliminary catalytic investigations of **2-Ln** also showed heterobimetallic cooperativity, enabling the dehydrocoupling of [Me₂HN·BH₃] at room temperature. Exploration of the reactivity and catalytic activity of these Ln/Al heterobimetallic trihydrides is ongoing to fully understand and utilize the synergistic metal–metal interaction.

■ ASSOCIATED CONTENT

Data Availability Statement

Additional data that support the findings of this study are available from the University of Strathclyde KnowledgeBase at 10.15129/f6a428f0-48e2-400e-81c2-465639e81845.

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.inorgchem.4c00824>.

Data that support the findings of this study and additional references²² (PDF)

Accession Codes

CCDC 2313687–2313695 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION**Corresponding Authors**

Catherine E. Weetman – Department of Pure and Applied Chemistry, University of Strathclyde, Glasgow G1 1XL, U.K.; orcid.org/0000-0001-5643-9256; Email: catherine.weetman@strath.ac.uk

Joy H. Farnaby – Joseph Black Building, School of Chemistry, University of Glasgow, Glasgow G12 8QQ, U.K.; orcid.org/0000-0002-1224-7823; Email: joy.farnaby@glasgow.ac.uk

Authors

Tajrian Chowdhury – Joseph Black Building, School of Chemistry, University of Glasgow, Glasgow G12 8QQ, U.K.; orcid.org/0000-0002-8728-5277

Fáinché Murphy – Department of Pure and Applied Chemistry, University of Strathclyde, Glasgow G1 1XL, U.K.; orcid.org/0009-0009-3176-260X

Alan R. Kennedy – Department of Pure and Applied Chemistry, University of Strathclyde, Glasgow G1 1XL, U.K.; orcid.org/0000-0003-3652-6015

Claire Wilson – Joseph Black Building, School of Chemistry, University of Glasgow, Glasgow G12 8QQ, U.K.; orcid.org/0000-0002-0090-5374

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.inorgchem.4c00824>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank Craig Irving for assistance with NMR spectroscopy. J.H.F. acknowledges the University of Glasgow for funding and the award of a College of Science & Engineering Ph.D. scholarship to T.C. C.E.W. thanks the University of Strathclyde for the award of a Chancellor's Fellowship and the award of a Ph.D. studentship (to F.M.).

REFERENCES

- (1) (a) Ephritikhine, M. Synthesis, Structure, and Reactions of Hydride, Borohydride, and Aluminohydride Compounds of the f-Elements. *Chem. Rev.* **1997**, *97* (6), 2193–2242. (b) Barbier-Baudry, D.; Dormond, A.; Visseaux, M. Early lanthanide organometallic hydrides: electronic and steric control of the stability. *J. Organomet. Chem.* **2000**, *609* (1), 21–28. (c) Barbier-Baudry, D.; Dormond, A. New Trends in Organolanthanide Hydride Chemistry. In *Recent Advances in Hydride Chemistry*; Peruzzini, M., Poli, R., Eds.; Elsevier, 2001; Chapter 8, pp 249–270. (d) Konkol, M.; Okuda, J. Non-metallocene hydride complexes of the rare-earth metals. *Coord. Chem. Rev.* **2008**, *252* (15), 1577–1591. (e) Fegler, W.; Venugopal, A.; Kramer, M.; Okuda, J. Molecular Rare-Earth-Metal Hydrides in Non-Cyclopentadienyl Environments. *Angew. Chem., Int. Ed.* **2015**, *54* (6), 1724–1736. (f) Harder, S. Molecular early main group metal hydrides: synthetic challenge, structures and applications. *Chem. Commun.* **2012**, *48* (91), 11165–11177. (g) Butler, M. J.; Crimmin, M. R. Magnesium, zinc, aluminium and gallium hydride complexes of the transition metals. *Chem. Commun.* **2017**, *53* (8), 1348–1365. (h) Roy, M. M. D.; Omaña, A. A.; Wilson, A. S. S.; Hill, M. S.; Aldridge, S.; Rivard, E. Molecular Main Group Metal Hydrides. *Chem. Rev.* **2021**, *121* (20), 12784–12965.
- (2) (a) Ferrence, G. M.; McDonald, R.; Takats, J. Stabilization of a Discrete Lanthanide(II) Hydrido Complex by a Bulky Tp^R Ligand. *Angew. Chem., Int. Ed.* **1999**, *38* (15), 2233–2237. (b) Cheng, J.; Ferguson, M. J.; Takats, J. Synthesis and Reaction of [(Tp^{RPr2})LnH₂]₃ (Ln = Y, Lu) with CO: Trinuclear Cluster-Bound Propenolate en Route to Selective Formation of Propene. *J. Am. Chem. Soc.* **2010**, *132* (1), 2–3. (c) Cheng, J.; Saliu, K.; Ferguson, M. J.; McDonald, R.; Takats, J. Variable nuclearity scorpionate-supported lanthanide polyhydrides: [(Tp^{R,R'})LnH₂]_n (n = 3, 4 and 6). *J. Organomet. Chem.* **2010**, *695* (25–26), 2696–2702. (d) Onishi, M.; Itoh, K.; Hiraki, K.; Oda, R.; Aoki, K. Syntheses, properties and structural characteristics of several new (polypyrazolylborato) samarium(III) complexes: agostic interaction of the BH₂ proton with samarium. *Inorg. Chim. Acta* **1998**, *277* (1), 8–13. (e) Saliu, K. O.; Takats, J.; Ferguson, M. J. Yb(II) [Yb(Tp^{3-tBu,5-Me})₂] toluene solvate. *Acta Crystallogr. E* **2009**, *65* (6), m643–m644. (f) Saliu, K. O.; Maunder, G. H.; Ferguson, M. J.; Sella, A.; Takats, J. Synthesis and structure of heteroleptic ytterbium (II) tetrahydroborate complexes. *Inorg. Chim. Acta* **2009**, *362* (12), 4616–4622. (g) Momin, A.; Bonnet, F.; Visseaux, M.; Maron, L.; Takats, J.; Ferguson, M. J.; Le Goff, X.-F.; Nief, F. Synthesis and structure of divalent thulium borohydrides, and their application in ε-caprolactone polymerisation. *Chem. Commun.* **2011**, *47* (44), 12203–12205. (h) Galler, J. L.; Goodchild, S.; Gould, J.; McDonald, R.; Sella, A. Tris-pyrazolylborate complexes of redox inactive lanthanides - the structures of [(Tp^{Me2})₂NdX] (X = Cl, NPh₂, dpm, H₂BET₂). *Polyhedron* **2004**, *23* (2–3), 253–262. (i) Schädle, C.; Schädle, D.; Eichele, K.; Anwander, R. Methylaluminum-Supported Rare-Earth-Metal Dihydrides. *Angew. Chem., Int. Ed.* **2013**, *52* (50), 13238–13242.
- (3) Schädle, D.; Anwander, R. Rare-earth metal and actinide organoimide chemistry. *Chem. Soc. Rev.* **2019**, *48* (24), 5752–5805.
- (4) Schädle, C.; Anwander, R. Rare-Earth-Metal-Promoted Hydroalumination. *Eur. J. Inorg. Chem.* **2013**, *2013* (19), 3302–3306.
- (5) (a) Chowdhury, T.; Horsewill, S. J.; Wilson, C.; Farnaby, J. H. Heteroleptic lanthanide(III) complexes: synthetic utility and versatility of the unsubstituted bis-scorpionate ligand framework. *Aust. J. Chem.* **2022**, *75* (9), 660–675. (b) Chowdhury, T.; Wilson, C.; Maichle-Mössner, C.; Anwander, R.; Farnaby, J. H. Lanthanide Amide Complexes Supported by the Bis-tris(pyrazolyl)borate Ligand Environment. *Eur. J. Inorg. Chem.* **2024**, *27* (11), No. e202300731.
- (6) (a) Stainer, M. V. R.; Takats, J. Solution structure of [Yb(Tp)₃]; rare example of solution rigid lanthanide complex. *J. Am. Chem. Soc.* **1983**, *105* (3), 410–415. (b) Apostolidis, C.; Rebizant, J.; Kanellakopulos, B.; Von Ammon, R.; Dornberger, E.; Müller, J.; Powietzka, B.; Nuber, B. Homoscorpionates (Tp complexes) of the trivalent 4f ions. The crystal and molecular structure of [Ln(Tp)₃] (Ln = Pr, Nd). *Polyhedron* **1997**, *16* (7), 1057–1068.
- (7) Hock, A.; Schneider, H.; Krauß, M. J.; Radius, U. Hydride Amide and Hydride Phenolate Complexes of NHC Coordinated Aluminum. *Z. Anorg. Allgem. Chem.* **2018**, *644* (21), 1243–1251.
- (8) Green, J. C.; Green, M. L. H.; Parkin, G. The occurrence and representation of three-centre two-electron bonds in covalent inorganic compounds. *Chem. Commun.* **2012**, *48* (94), 11481–11503.
- (9) Stennett, C. R.; Wagner, C. L.; Fetting, J. C.; Vasko, P.; Power, P. P. Reductions of M(Nⁿ)₃ (M = V, Cr, Fe): Terminal and Bridging Low-Valent First-Row Transition Metal Hydrido Complexes and “Metallo-Transamination. *Inorg. Chem.* **2021**, *60* (15), 11401–11411.

- (10) Tang, J. A.; Masuda, J. D.; Boyle, T. J.; Schurko, R. W. Ultra-wide-line ^{27}Al NMR Investigation of Three- and Five-Coordinate Aluminum Environments. *ChemPhysChem* **2006**, *7* (1), 117–130.
- (11) Schädle, C.; Fischbach, A.; Herdtweck, E.; Törnroos, K. W.; Anwänder, R. Reactivity of Yttrium Carboxylates Toward Alkylaluminum Hydrides. *Chem. Eur. J.* **2013**, *19* (48), 16334–16341.
- (12) (a) Veith, M.; König, P.; Rammo, A.; Huch, V. Cubane-Like Li_4H_4 and $\text{Li}_3\text{H}_3\text{Li}(\text{OH})$: Stabilized in Molecular Adducts with Alanes. *Angew. Chem., Int. Ed.* **2005**, *44* (37), 5968–5971. (b) Pollard, V. A.; Orr, S. A.; McLellan, R.; Kennedy, A. R.; Hevia, E.; Mulvey, R. E. Lithium diamidodihydridoaluminates: bimetallic cooperativity in catalytic hydroboration and metallation applications. *Chem. Commun.* **2018**, *54* (10), 1233–1236.
- (13) Chowdhury, T.; Evans, M. J.; Coles, M. P.; Bailey, A. G.; Peveler, W. J.; Wilson, C.; Farnaby, J. H. Reduction chemistry yields stable and soluble divalent lanthanide tris(pyrazolyl)borate complexes. *Chem. Commun.* **2023**, *59* (15), 2134–2137.
- (14) (a) Cole, M. L.; Jones, C.; Junk, P. C.; Kloth, M.; Stasch, A. Synthesis and Characterization of Thermally Robust Amidinato Group 13 Hydride Complexes. *Chem. Eur. J.* **2005**, *11* (15), 4482–4491. (b) Aldabbagh, A. K.; Takjoo, R.; Najafi, A.; Guo, Z.; Hossain, M. E.; Rad, N. E.; Wang, J.; Junk, P. C. Syntheses and structural characterisation of some heteroleptic aluminium(III) formamidinates. *Aust. J. Chem.* **2022**, *75* (4), 265–275.
- (15) Cole, M. L.; Deacon, G. B.; Forsyth, C. M.; Junk, P. C.; Konstas, K.; Wang, J. Steric Modulation of Coordination Number and Reactivity in the Synthesis of Lanthanoid(III) Formamidinates. *Chem. Eur. J.* **2007**, *13* (29), 8092–8110.
- (16) (a) Ali, S. H.; Deacon, G. B.; Junk, P. C.; Hamidi, S.; Wiecko, M.; Wang, J. Lanthanoid Pseudo-Grignard Reagents: A Major Untapped Resource. *Chem. Eur. J.* **2018**, *24* (1), 230–242. (b) Werner, D.; Zhao, X.; Best, S. P.; Maron, L.; Junk, P. C.; Deacon, G. B. Bulky Ytterbium Formamidinates Stabilise Complexes with Radical Ligands, and Related Samarium “Tetracyclone” Chemistry. *Chem. Eur. J.* **2017**, *23* (9), 2084–2102.
- (17) Rumble, J. *CRC Handbook of Chemistry and Physics*, 104th ed.; CRC Press, 2023; p 1604.
- (18) (a) Bashirov, D. A.; Kolybalov, D. S.; Sukhikh, T. S.; Konchenko, S. N. Synthesis, Structure, and Photoluminescent Properties of Lanthanide (Ln = Dy, Tb) Chlorides and Thiophenolates Supported by Formamidinate Ligands. *J. Struct. Chem.* **2020**, *61* (8), 1219–1226. (b) Salehisaki, M.; Rad, N. E.; Deacon, G. B.; Wang, J.; Guo, Z.; Junk, P. C. Synthesis and reactivity of rare-earth- N,N' -(diphenyl)formamidinate and rare-earth- N,N' -bis(2,4-dimethylphenyl)formamidinate complexes. *Inorg. Chim. Acta* **2022**, *539*, 120997.
- (19) Brown, A. C.; Altman, A. B.; Lohrey, T. D.; Hohloch, S.; Arnold, J. Hydride oxidation from a titanium–aluminum bimetallic complex: insertion, thermal and electrochemical reactivity. *Chem. Sci.* **2017**, *8* (7), 5153–5160.
- (20) (a) Melen, R. L. Dehydrocoupling routes to element–element bonds catalysed by main group compounds. *Chem. Soc. Rev.* **2016**, *45* (4), 775–788. (b) Rossin, A.; Peruzzini, M. Ammonia–Borane and Amine–Borane Dehydrogenation Mediated by Complex Metal Hydrides. *Chem. Rev.* **2016**, *116* (15), 8848–8872. (c) Reuter, M. B.; Seth, D. M.; Javier-Jiménez, D. R.; Finfer, E. J.; Beretta, E. A.; Waterman, R. Recent advances in catalytic pnictogen bond forming reactions via dehydrocoupling and hydrofunctionalization. *Chem. Commun.* **2023**, *59* (10), 1258–1273.
- (21) (a) Cui, P.; Spaniol, T. P.; Maron, L.; Okuda, J. Dehydrogenation of Amine–Borane $\text{Me}_2\text{NH}\cdot\text{BH}_3$ Catalyzed by a Lanthanum–Hydride Complex. *Chem. Eur. J.* **2013**, *19* (40), 13437–13444. (b) Rad'kov, V.; Dorcet, V.; Carpentier, J.-F.; Trifonov, A.; Kirillov, E. Alkylaluminum Complexes of Amidine–Amidopyridinate Ligands. Intramolecular $\text{C}(\text{sp}^3)\text{--H}$ Activation and Reactivity Studies. *Organometallics* **2013**, *32* (5), 1517–1527. (c) Less, R. J.; Simmonds, H. R.; Dane, S. B. J.; Wright, D. S. Stoichiometric and catalytic reactions of LiAlH_4 with Me_2NHBH_3 . *Dalton Trans.* **2013**, *42* (18), 6337–6343. (d) Less, R. J.; Simmonds, H. R.; Wright, D. S. Reactivity and catalytic activity of *tert*-butoxy-aluminium hydride reagents. *Dalton Trans.* **2014**, *43* (15), 5785–5792. (e) Weetman, C.; Ito, N.; Unno, M.; Hanusch, F.; Inoue, S. NHI- and NHC-Supported Al(III) Hydrides for Amine–Borane Dehydrocoupling Catalysis. *Inorganics* **2019**, *7* (8), 92–102.
- (22) (a) Murphy, F.; Kennedy, A. R.; Weetman, C. E. Synthesis and Structural Comparisons of NHC–Alanes. *Inorganics* **2023**, *11* (1), 13. (b) Ruff, J. K.; Hawthorne, M. F. The Amine Complexes of Aluminum Hydride. I. *J. Am. Chem. Soc.* **1960**, *82* (9), 2141–2144. (c) Jafarpour, L.; Stevens, E. D.; Nolan, S. P. A sterically demanding nucleophilic carbene: 1,3-bis(2,6-diisopropylphenyl)imidazol-2-ylidene. Thermochemistry and catalytic application in olefin metathesis. *J. Organomet. Chem.* **2000**, *606* (1), 49–54. (d) Baker, R. J.; Davies, A. J.; Jones, C.; Kloth, M. Structural and spectroscopic studies of carbene and *N*-donor ligand complexes of Group 13 hydrides and halides. *J. Organomet. Chem.* **2002**, *656* (1), 203–210. (e) SHELXL; Bruker AXS Inc.: Madison, WI, 2016. (f) CrysAlisPro; Rigaku Ltd.: Yarnton, Oxfordshire, England, 2019. (g) Sheldrick, G. M. Crystal structure refinement with SHELXL. *Acta Crystallogr. C* **2015**, *71* (1), 3–8. (h) Dolomanov, O. V.; Bourhis, L. J.; Gildea, R. J.; Howard, J. A. K.; Puschmann, H. OLEX2: a complete structure solution, refinement and analysis program. *J. Appl. Crystallogr.* **2009**, *42* (2), 339–341.