

Communication

Reactivity and Stability of a Ring-Expanded N-Heterocyclic Carbene Copper(I) Boryl Imidinate

Rex S. C. Charman , Thomas M. Horsley Downie, Thomas H. Jerome, Mary F. Mahon and David J. Liptrot * 

Department of Chemistry, University of Bath, Claverton Down, Bath BA2 7AY, UK

* Correspondence: d.j.liptrot@bath.ac.uk

Abstract: Frustrated Lewis pairs (FLPs) have evolved from a revolutionary concept to widely applied catalysts. We recently reported the ring-expanded N-heterocyclic carbene supported copper(I) boryliminomethanide, (6-Dipp)CuC(=N^tBu)Bpin and noted it reacted with heterocumulenes in a fashion reminiscent of FLPs. We thus set out to explore its reactivity with a range of other substrates known to react with FLPs. This was undertaken by a series of synthetic studies using NMR spectroscopy, mass spectrometry, IR spectroscopy, and single crystal X-ray crystallography. (6-Dipp)CuC(=N^tBu)Bpin was investigated for its reactivity towards water, hydrogen, and phenylacetylene. Its solution stability was also explored. Upon heating, (6-Dipp)CuC(=N^tBu)Bpin decomposed to (6-Dipp)CuCN, which was characterised by SC-XRD and NMR spectroscopy, and pinB^tBu. Although no reaction was observed with hydrogen, (6-Dipp)CuC(=N^tBu)Bpin reacted with water to form (6-Dipp)CuC(=N(H)^tBu)B(OH)pin, which was structurally characterised. In contrast to its FLP-reminiscent heterolytic cleavage reactivity towards water, (6-Dipp)CuC(=N^tBu)Bpin acted as a Brønsted base towards phenyl acetylene generating (6-Dipp)CuCCPh, which was characterised by SC-XRD, IR, and NMR spectroscopy, and HC(=N^tBu)Bpin

Keywords: frustrated Lewis pair; copper; N-heterocyclic carbene



Citation: Charman, R.S.C.; Horsley Downie, T.M.; Jerome, T.H.; Mahon, M.F.; Liptrot, D.J. Reactivity and Stability of a Ring-Expanded N-Heterocyclic Carbene Copper(I) Boryl Imidinate. *Inorganics* **2022**, *10*, 135. <https://doi.org/10.3390/inorganics10090135>

Academic Editors: Stephen Mansell and Simon Aldridge

Received: 16 August 2022

Accepted: 2 September 2022

Published: 7 September 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).


1. Introduction

The “Frustrated Lewis Pair” (FLP) concept has contributed to a revolution in small molecule activation with transition metal-free systems, as well as in diverse other fields, including organic synthesis [1,2]. Though the original concept relied upon the combination of strong Lewis acid (LA) and base fragments (LB) that could not form a classical Lewis adduct, there is an increasing recognition that almost any set of appropriately oriented lone pairs and vacant orbitals can be considered within the FLP conceptual framework [3].

The major attraction of FLPs is their ability to activate molecules in a transition metal-mimetic fashion. For example, Stephan’s first report of an FLP was noted for the reversible binding of dihydrogen [4] that could be exploited in hydrogenation [5]. Since this initial report, a plethora of other E-H bonds have been activated across FLPs. Of particular relevance to this work is the heterolytic activation of water to generate a Lewis acid-hydroxide fragment and a protonated Lewis base [6]. A similar reaction has also been observed for some C-H acidic fragments, most notably acetylenes which generate a Lewis acid-acetylide and protonated Lewis base pair [7]. Carbon dioxide and analogous heterocumulenes are also activated by FLPs, generally generating a LB-C(O)O-LA moiety [8,9].

One structural motif which has seen some interest for its FLP-type activity is the carbiminoborane scaffold which comprises a R₂B-C(R)=NR fragment where the 3-coordinate boron centre is Lewis acidic, and the 2-coordinate imine nitrogen, which retains a lone pair, constitutes the Lewis base. Figueroa and co-workers reported that 1,1-hydroboration of an isonitrile generated a carbiminoborane, Cy₂BC(H)=NAr (Ar = 2,6-(ⁱPr)₂C₆H₃)₂C₆H₃), that displayed FLP-like reactivity towards carbon dioxide, nitriles, and acetylenes

(Scheme 1) [10,11]. For example, the reaction with CO₂ gave rise to a 5-membered N-C-B-O-C ring that was proposed to form via a 1,2-alkyl transfer of the cyclohexyl group on boron from Cy₂BC(H)=NAr·CO₂. In contrast, E-H bonds in water and acetylene were cleaved in a heterolytic fashion to give CyB(E)C(Cy)(H)=N(H)Ar (E = CC^tBu, OH) once again via a FLP-activation/cyclohexyl migration mechanism. Erker and co-workers reported that a mixture of an isonitrile and (Fmes)BH₂ (Fmes = 2,4,6-(F₃C)₃C₆H₂) would react with a wide range of organic unsaturates, chemistry they attributed to an iminoborane, Fmes(H)BC(H)NR (R = H₂CPh, 2,6-Me₂C₆H₃), formed in situ [12]. For example, Fmes(H)BC(H)NCH₂Ph reacted with anisyl acetylene via a C-H cleavage FLP-type intermediate, Fmes(H)B(CCC₆H₄-4-OMe)C(H)N(H)CH₂Ph (Scheme 1). We recently reported that the reaction of heterocumulenes with a copper(I) boryl generated isonitriles [13]. These species reacted further with the copper(I) boryl to furnish a copper(I) boryliminomethanide, (6-Dipp)CuC(=N^tBu)Bpin, which reacted with heterocumulenes in an analogous fashion to that reported for the systems of Erker and Figueroa.



Scheme 1. A number of borylimines described in the literature and their FLP-type reactivity towards Brønsted acids and heterocumulenes. Fmes = 2,4,6-(F₃C)₃C₆H₂, Dipp = 2,6-(^tPr)₂C₆H₃.

Herein we report the further exploration of this copper(I) boryliminomethanide, including our attempts at its structural characterisation, its long-term stability, and its reactivity towards a number of other substrates.

2. Results and Discussion

We began our work by resynthesising (6-Dipp)CuC(=N^tBu)Bpin (compound **1**) via the reaction of tert-butyl isocyanide with isolated (6-Dipp)CuBpin. Single crystals of this compound were grown by vapour diffusion of hexane into a toluene solution of **1**. Unfortunately, despite repeated data collections, a satisfactory structural refinement beyond confirmation of connectivity (Figure 1a) could not be obtained for this compound. Thus, the metric data for **1**, provided below, will not be discussed or interpreted in any detail. On one occasion, adventitious moisture resulted in hydrolysis of **1**, and the crystal structure of

the resulting compound was obtained to show (6-Dipp)CuC(B(OH)(pin)=N(H)^tBu (1·H₂O) (Figure 1b).





Figure 1. (a) Ball and stick drawing arising from a preliminary structural determination of **1** (for additional information regarding the limitations of this diffraction experiment, see ESI). (b) Crystal structure of **1**·H₂O (30% probability ellipsoids) where lattice solvent and hydrogen atoms (except for those bonded to O1 and N3) have been omitted for visual clarity. Wireframe view has been employed for Dipp groups, also for visual simplicity. Selected bond lengths (Å), angles (°), and geometric parameters: (a) C1–Cu1 1.934(7); Cu1–C29 1.937(9); C29–B1 1.59(2); C29–N3 1.265(14); N1–C1–N2 116.2(7); C1–Cu1–C29 174.0(5); N3–C29–B1 105.0(9). (b) N1–C1 1.3431(15); N2–C1 1.3424(16); C1–Cu1 1.9331(12); Cu1–C29 1.9141(12); C29–B1 1.6481(18); B1–O1 1.4632(17); B1–O2 1.4887(16); B1–O3 1.4651(17); C29–N3 1.2937(17); N3–C30 1.4873(17); N1–C1–N2 116.77(11); C1–Cu1–C29 173.62(5); Cu1–C29–B1 118.59(9); Cu1–C29–N3 130.24(10); C29–N3–C30 132.50(11); B1 τ₄ 0.93.

These results provide strong supporting evidence for the structure we previously proposed for the boryliminomethanide. In **1**·H₂O, the O–H bond of water has been cleaved with the transfer of a proton to the imine N atom and a hydroxy group to the B atom (Scheme 2). Similar reactivity was observed by Figueroa and co-workers for Cy₂BC(H)=NAr and (9-BBN)C(H)=NAr [10,11] and is typical for FLPs [6]. This reaction demonstrates how **1**, with a Lewis acidic boron centre in proximity to a Lewis basic nitrogen centre, can act as a reactive B/N FLP-type system in E–H cleavage reactions consistent with its previously reported reactivity towards heterocumulenes.



Scheme 2. The reaction between compound **1** and water to yield **1**·H₂O.

Compound **1**·H₂O comprises a 2-coordinate copper atom in a near-linear coordination environment (C1-Cu1-C29 173.62(5)°) with C_{carbene}-Cu bond lengths (C1-Cu1 1.9331(12) Å) and N-C_{carbene}-N parameters commensurate (N1-C1 1.3431(15); N2-C1 1.3424(16) Å, N1-C1-N2 116.77(11)°) with other (6-Dipp)CuX systems. The boratoimidiniummethanide fragment is relatively unusual, with limited literature examples to compare to. Nevertheless, in the context of the FLP-esque activity of **1**, its Cu-C(Bpin)=N^tBu core can be compared to the previously reported isocyanate activation product **1**·PhNCO. The copper-carbon bond in this fragment is only marginally shorter in **1**·H₂O (Cu1-C29 1.9141(12) vs. 1.925(2) Å), and the C-B bond length is very similar (C29-B1 1.6481(18) vs. 1.656(3) Å). The C-N bond lengths are similar, although the bond is slightly shorter in **1**·H₂O (1.2937(17) vs. 1.314(3) Å). These data allow attribution of both systems as largely retaining a C=N double bond, but the slightly shorter bond in **1**·H₂O is unsurprising in light of the more conjugated system in **1**·PhNCO [13]. We previously reported that the ¹¹B NMR spectrum of **1** contained a broad resonance at 28.3 ppm (FWHM = 610 Hz) [13], whereas a similar analysis of **1**·H₂O provided a sharp peak at 5.3 ppm (FWHM = 36 Hz).

During an attempt to recrystallise a toluene solution of **1** in a boiling water bath, we noted the onset of what we suspected to be decomposition due to changes in the colour of the solution and the formation of a precipitate. To characterise the products of this decomposition, we heated a C₆D₆ solution of **1**, generated in situ, to 80 °C overnight to produce a colourless solution and a crystalline material. The ¹¹B NMR spectrum of this reaction mixture contained a resonance at 35.0 ppm that was attributed to pinB^tBu based on comparison to literature data [14]. The crystalline material was found to be suitable for single crystal X-ray crystallography, which showed it to be (6-Dipp)CuCN (compound **2**, Figure 2).





Figure 2. Molecular structure (30% probability ellipsoids) of **2**. Hydrogens have been omitted for clarity. Wireframe view has been employed for Dipp groups, also for visual simplicity. Selected bond lengths (Å), angles (°): N1-C1 1.3402(13); C1-Cu1 1.9187(17); Cu1-C16 1.905(2); C16-N2 1.110(3); N1-C1-N1¹ 117.55(15); N1-C1-Cu1 121.20(8); C1-Cu1-C16 177.49(9); Cu1-C16-N2 179.9(2). Symmetry operations (1); $x, \frac{1}{2} - y, z$.

This reactivity can be summarised as a 1,3-*tert*-butyl transfer from the imine nitrogen to the pinB- fragment, as shown in Scheme 3. Moreover, we found that samples of **1** left over the course of months at ambient temperature would undergo this reaction and were, thus, not stable long term.



Scheme 3. The decomposition of compound **1** on heating to generate compound **2**.

Compound **2** comprises a 2-coordinate copper atom bonded to both the 6-Dipp and cyanide ligands. It is essentially linear at copper (C1-Cu1-C16 177.49(9)°), as was previously observed by Nolan and co-workers for (IPr)CuCN (177.5(3)°) [15]. Compared to (IPr)CuCN, the copper-cyanide fragment of **2** also has a similar C-N distance (C16-N2 1.110(3) vs. 1.139(6) Å) and Cu-C-N bond angle (Cu1-C16-N2 179.9(2) vs. 177.9(7)°). The steric demand of the 6-Dipp ligand, however, provides for slightly longer C_{carbene}-Cu (C1-Cu1 1.9187(17) vs. 1.900(4) Å) and Cu-C_{ciano} (Cu1-C16 1.905(2) vs. 1.862(5) Å) distances. Attempts to observe a ¹³C NMR signal for the cyanide carbon were unsuccessful. Nolan and co-workers observed a resonance at 142.4 ppm to this nucleus [15], but similar resonances in the ¹³C NMR spectrum of **3** (146.7, 142.4 ppm) were assigned to substituted aromatic carbons based on **3**'s ¹H-¹³C HMBC spectrum, which showed that these carbon nuclei coupled to the iso-propyl methine and other aromatic hydrogens on the ligand.

Seeking to further extend the FLP-esque reactivity of **1**, we investigated its reaction with a number of other molecules. The addition of an atmosphere of H₂ to a C₆D₆ solution of **1** provided no evidence of reaction, and a ¹H NMR spectrum containing both free H₂ and compound **1**, and heating of the reaction only led to the onset of decomposition. The addition of excess *tert*-butyl isocyanide was similar, providing no evidence of reactivity, unlike the systems described by Erker. A reaction was observed, however, with an acetylene.

The reaction of C-H acidic substrates with FLPs often gives access to C-H cleavage products analogous to the O-H cleavage product, 1·H₂O. The reaction of compound **1** with a monosubstituted alkyne was thus also investigated. The addition of one equivalent of phenyl acetylene to **1** provided a ¹H NMR spectrum consistent with a reaction having occurred, with the complete attenuation of the resonance associated with the acetylene C-H and a 6-Dipp region consistent with the formation of one major new ligand-containing compound. Material deposited from this reaction was subjected to single crystal X-ray crystallography, which showed it to be (6-Dipp)CuC≡CPh, compound **3** (Figure 3).

Compound **3** is thus the result not of C-H activation across the FLP fragment but of the Cu-C bond in **1** acting as a Brønsted base towards the acidic C-H bond of the acetylene. This reaction should occur with concomitant formation of HC(=N^{*t*}Bu)Bpin, evidence for which was provided by subjecting the reaction mixture to mass spectrometry (Scheme 4). An ion with an *m/z* of 86.0962 was observed in the positive mode spectrum and assigned to the [M+H]⁺ ion of H₂C=N^{*t*}Bu. This ion was proposed to form through the hydrolysis of the C-B bond of HC(=N^{*t*}Bu)Bpin by residual moisture in the solvent used to make up the sample.





Figure 3. Molecular structure (30% probability ellipsoids) of **3**. Hydrogen atoms and the second disordered component have been omitted for visual clarity. Wireframe view has been employed for Dipp groups, also for visual simplicity. Selected bond lengths (Å), angles (°): N1-C1 1.339(3); N2-C1 1.474(3); C1-Cu1 1.919(2); Cu1-C29 1.867(2); C29-C30 1.208(3); C30-C31 1.450(4); N1-C1-N2 117.49(18); N1-C1-Cu1 116.50(15); N2-C1-Cu1 126.01(16); C1-Cu1-C29 176.50(10); Cu1-C29-C30 176.6(2); C29-C30-C31 174.0(5).




Scheme 4. The Brønsted basic nature of compound **1** in its reaction with phenyl acetylene to yield **3**.

Compound **3** is once again an example of a near linear (C1-Cu1-C29 176.50(10)°) copper(I) complex. The related NHC compound (IPr)CuCCPh has been reported by both Kleeberg and co-workers [16] and the Jones group [17]. The metric parameters of these systems are largely identical, and those reported by Jones, which were also collected at 150 K, will be discussed here. Once again, the ring-expansion to 6-Dipp results in an elongation of the C_{carbene}-Cu (C1-Cu1 1.919(2) vs. 1.890(4) Å) but not the Cu-C_{acetylene}

(Cu1-C29 1.867(2) vs. 1.861(4) Å) bond lengths. The C≡C triple bond is also essentially unchanged (C29-C30 1.208(3) vs. 1.209(5) Å), and both systems display a similar distortion of the Cu-C≡C-Ph fragment away from linearity (Cu1-C29-C30 176.6(2) vs. 169.3(4)°; C29-C30-C31 174.0(5) vs. 177.4(4)°), although this is less pronounced in **3**. This latter observation may reflect the effect of the marginally longer C_{carbene}-Cu bond, or the electronic variation between the 6-Dipp and IPr ligands. Although the ¹H NMR spectrum of **3** is unremarkable, comprising the expected resonances for the 6-Dipp ligand and a phenyl group, the ¹³C NMR spectrum was more useful. Resonances at 122.6 and 105.3 were attributed to the alkyne carbon atoms, specifically as the carbon bonded to copper and to the phenyl substituent, respectively. Further evidence for the presence of an intact alkyne fragment in **3** could be found in the IR spectrum, which showed a sharp peak at 2086.92 cm⁻¹ that was assigned to a C≡C bond stretch.

The difference in the reactivity of compound **1** towards water and phenyl acetylene is notable, especially in the light of the much lower pK_a of O-H bonds compared to C^{sp}-H bonds. We tentatively attribute this deviation to the effect of the lone pairs on the oxygen atom of water, which are capable of forming an adduct with the Bpin fragment of compound **1**, thus predisposing it to the FLP-type reactivity observed to yield **1**·H₂O. In contrast, the absence of such a hard donor group on phenyl acetylene precludes such a coordination, and thus, the reaction occurs at the Cu-C bond to yield the acetylide, **3**. Further evidence in favour of this hypothesis comes from repeated attempts to rationally synthesise **1**·H₂O, which yielded material contaminated with (6-Dipp)CuOBpin. The formation of (6-Dipp)CuOBpin can be considered as the result of the Cu-C bond in **1**·H₂O deprotonating its {pinBOH} fragment and extruding a fragment of the formula H₂CN^tBu, although the precise mechanism by which this occurs is unclear (Scheme 5). We interpreted this to indicate that **1**·H₂O is the kinetic product of the activity of **1** towards water, a protic substrate with active lone pairs, but such a pathway is inhibited in the case of acetylenes, owing to the weak interaction between the C-C π-bond and vacant p-orbital on boron, thus resulting only in direct deprotonation to give **3**.



Scheme 5. The generation of (6-Dipp)CuOBpin from **1**·H₂O.

In conclusion, compound **1**, (6-Dipp)CuC(=N^tBu)Bpin, shows less reliable FLP activity than we initially hoped. Despite its previously reported activation of isocyanates when generated and used quickly, compound **1** is unstable in our hands. Thermolysis and long-term storage both led to decomposition to generate compound **2**, (6-Dipp)CuCN, the first structurally characterised ring-expanded NHC-copper(I) cyanide complex. Furthermore, no reaction towards H₂ was observed. Finally, compound **1** showed divergent reactivity towards different protic reagents. The serendipitous addition of water provided O-H activation across the boryliminomethanide fragment to generate an iminium borate, **1**·H₂O, in a fashion reminiscent of an FLP. In contrast, the addition of a monosubstituted acetylene provided the protonation of the boryliminomethanide to generate the corresponding boryliminomethane and (6-Dipp)CuCCPh, **3**, the first ring-expanded NHC-copper(I) acetylide to be structurally characterised. We tentatively attribute this deviation in reactivity to the presence of a lone pair on the oxygen atom of water that facilitates O-H cleavage across the B/N fragment via the initial formation of an adduct with the boron atom. Thus, we were able to map the range of decomposition and bond activation pathways available to

compound **1** which comprise E-H activation across the boryliminomethanide, 1,3-*tert*-butyl transfer, and Brønsted basicity of the boryliminomethanide.

3. Materials and Methods

General Considerations and Starting Materials

All reactions dealing with air- and moisture-sensitive compounds were carried out under argon atmosphere using standard Schlenk line and glovebox techniques. NMR experiments using air sensitive compounds were conducted in J. Young's tap NMR tubes prepared and sealed in a glovebox under argon. Toluene and hexane were purified using an MBraun Solvent Purification System and stored over 4 Å molecular sieves. C₆D₆ and *d*₈-THF were dried over a potassium mirror prior to vacuum transfer into a sealed ampoule and stored in a glove box under argon. All NMR data, unless otherwise stated, were acquired at 298 K on an Agilent ProPulse instrument for ¹H (500 MHz), ¹³C{¹H} (126 MHz) and ¹¹B (160 MHz). ¹H and ¹³C NMR spectra were referenced using residual solvent resonances. X-ray data were collected on a New Xcalibur EosS2 diffractometer with Mo source ($\lambda = 0.71073$), or on a RIGAKU SuperNova with Cu source ($\lambda = 1.54184$). Using Olex2 [18] the structures were solved with the SHELXT [19] and refined with the ShelXL [19] refinement package using Least Squares minimisation. Mass spectrometry was performed using a Bruker MicrOTOF Electrospray Time-Of-Flight Mass Spectrometer coupled to an Agilent High Performance Liquid Chromatography unit. Elemental analyses were performed by Elemental Microanalysis Ltd., Okehampton, Devon, U.K. IR data was collected with a Bruker ALPHA II spectrometer with ATR accessory inside an inert atmosphere glovebox. *Tert*-Butyl isocyanide was purchased from VWR and was used as received. Bis(pinacolato)diboron and phenyl acetylene were purchased from Fluorochem and used as received. (6-Dipp)CuO^{*t*}Bu and (6-Dipp)CuBpin were prepared according to literature procedures [13,20].

Synthesis of compound 1·H₂O, (6-Dipp)CuC(=N(H)^{*t*}Bu)B(OH)pin

In a vial in a glovebox, ^{*t*}BuNC (11.4 μ L, 0.101 mmol) was added to a solution of (6-Dipp)CuBpin (30 mg, 0.050 mmol) in C₆H₆ (0.5 mL), forming a yellow solution. Volatiles were removed in vacuo, precipitating a yellow powder. The solid was re-dissolved in C₆D₆ (0.5 mL) and added to a solution of H₂O (0.9 μ L, 0.050 mmol) in C₆D₆ (0.5 mL). A pale brown solution formed, which was transferred to J. Young's tap NMR tube. ¹H, ¹¹B, and ¹³C NMR spectra were taken, confirming the formation of compound 1·H₂O, (6-Dipp)CuC(B(OH)(pin)=N(H)^{*t*}Bu. Colourless crystalline material precipitated out of solution overnight, which was confirmed to be (6-Dipp)CuOBpin by ¹H and ¹¹B NMR spectroscopy, yield 11.8 mg (38%).

¹H NMR (500 MHz, C₆D₆) δ 10.59 (s, 1H, N-H), 7.24–7.20 (m, 2H, *para*-H), 7.12 (d, $J = 7.8$ Hz, 4H, *meta*-H), 3.08–3.01 (m, 4H, CH(CH₃)₂), 2.80 (t, $J = 5.3$ Hz, 4H, NCH₂), 1.57 (s, 6H, OC(CH₃)₂), 1.56–1.52 (m, 14H, CH(CH₃)₂ and NCH₂CH₂), 1.20 (d, $J = 7.0$ Hz, 12H, CH(CH₃)₂), 1.11 (s, 6H, OC(CH₃)₂), 0.63 (s, 9H, NC(CH₃)₃), –0.89 (s, 1H, OH). ¹³C NMR (126 MHz, C₆D₆) δ 203.8 (CuC), 145.7 (Ar-C), 145.6 (Ar-C), 141.9 (Ar-C), 129.7 (Ar-C), 125.3 (Ar-C), 78.0 (OC(CH₃)₂), 54.1 (NC(CH₃)₃), 46.8 (NCH₂), 29.5 (NC(CH₃)₃), 29.0 (CH(CH₃)₂), 27.1 (OC(CH₃)₂), 26.7 (OC(CH₃)₂), 25.2 (CH(CH₃)₂), 25.1 (CH(CH₃)₂), 20.3 (NCH₂CH₂). ¹¹B NMR (160 MHz, C₆D₆) δ 5.3 (FWHM = 36 Hz). MS (ESI) [M+H] C₃₉H₆₄BCuN₃O₃: 696.4337, found: 696.4360 (err [ppm] = 0.44).

Synthesis of compound 2, (6-Dipp)CuCN

In a vial in a glovebox, ^{*t*}BuNC (9.5 μ L, 0.084 mmol) was added to a solution of (6-Dipp)CuBpin (25 mg, 0.042 mmol) in C₆H₆ (0.5 mL), forming a yellow solution. Volatiles were removed in vacuo precipitating a yellow powder. The solid was re-dissolved in C₆D₆ (1 mL), and the solution was placed in a sealed ampoule. The solution was heated to 80 °C overnight, forming a brown solution. Upon cooling to room temperature, a small amount of colourless material crystallised out of the solution, which was confirmed by ¹H and ¹³C NMR spectroscopy to be compound 2, (6-Dipp)CuCN. The crystalline material was investigated by SC-XRD.

^1H NMR (500 MHz, d_8 -THF) δ 7.37–7.35 (m, 2H, *para*-H), 7.26 (d, $J = 7.6$ Hz, 4H, *meta*-H), 3.58 (residual THF), 3.46 (t, $J = 5.9$ Hz, 4H, NCH₂), 3.12 (hept, $J = 6.9$ Hz, 4H, CH(CH₃)₂), 2.45 (water impurity), 2.35 (p, $J = 5.8$ Hz, 2H, NCH₂CH₂), 1.73 (residual THF), 1.34 (d, $J = 7.0$ Hz, 12H, CH(CH₃)₂), 1.30 (d, $J = 7.0$ Hz, 12H, CH(CH₃)₂). ^{13}C NMR (126 MHz, d_8 -THF) δ 201.9 (CuC_{carbene}), 146.7 (Ar-C), 142.4 (Ar-C), 130.3 (Ar-C), 125.6 (Ar-C), 47.3 (NCH₂), 29.6 (CH(CH₃)₂), 24.9 (CH(CH₃)₂), 21.3 (NCH₂CH₂). MS (ESI) [M+Na] C₂₉H₄₀CuN₃Na: 516.2416, found: 516.2423 (err [ppm] = −1.86).

Synthesis of compound 3, (6-Dipp)CuC≡CPh,

In a vial in a glovebox, a solution of (6-Dipp)CuO^{*t*}Bu (30 mg, 0.055 mmol) in C₆H₆ (0.5 mL) was added to B₂pin₂ (14 mg, 0.055 mmol). Volatiles were removed in vacuo, precipitating a white powder. The powder was washed with hexane (2 mL) and dried in vacuo. The remaining solid was dissolved in C₆H₆ (0.5 mL) and ^{*t*}BuNC (12.5 μL , 0.111 mmol) was added, forming a yellow solution. Volatiles were removed in vacuo, precipitating a yellow powder. The solid was re-dissolved into C₆D₆ (0.5 mL), and the reaction mixture was transferred to J. Young's tap NMR tube. An equivalent of phenyl acetylene (6.1 μL , 0.055 mmol) was added, forming an orange solution. Colourless material crystallised out of solution, which was confirmed by ^1H and ^{13}C NMR spectroscopy to be compound 4, (6-Dipp)CuCCPh. Yield 22 mg (70%). The crystalline material was investigated by SC-XRD.

^1H NMR (500 MHz, C₆D₆) δ 7.36–7.34 (m, 2H, *ortho*-H), 7.18–7.12 (m, 5H, Ar-H, residual solvent signal overlapped), 7.08–7.06 (m, 4H, *meta*-H), 6.84–6.80 (m, 2H, *meta*-H), 6.77–6.73 (m, 1H, *para*-H), 3.02 (h, $J = 6.9$ Hz, 4H, CH(CH₃)₂), 2.68 (t, $J = 5.9$ Hz, 4H, NCH₂), 1.52 (d, $J = 6.8$ Hz, 12H, CH(CH₃)₂), 1.45 (p, $J = 5.9$ Hz, 2H, NCH₂CH₂), 1.18 (d, $J = 7.0$ Hz, 12H, CH(CH₃)₂). ^{13}C NMR (126 MHz, C₆D₆) δ 203.4 (CuC), 145.6 (Ar-C), 141.6 (Ar-C), 132.3 (Ar-C), 129.7 (Ar-C), 129.6 (Ar-C), 127.6 (Ar-C), 124.9 (Ar-C), 124.5 (Ar-C), 122.6 (CuCC), 105.3 (CuCC), 46.1 (NCH₂), 28.9 (CH(CH₃)₂), 25.3 (CH(CH₃)₂), 24.6 (CH(CH₃)₂), 20.3 (NCH₂CH₂). Analysis calculated for C₃₆H₄₅CuN₂ (MW = 569.32 g/mol): Expected: C, 75.95; H, 7.97; N, 4.92%. Found: C, 74.08; H, 7.08; N, 4.75%. Despite repeated attempts, we were unable to obtain satisfactory elemental analysis for this compound. This was most likely due to the high sensitivity towards air and moisture. IR (ATR) cm^{−1} 2957.84 (C-H stretch), 2086.92 (C≡C stretch).

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/inorganics10090135/s1>, Table S1: Crystal data and structure refinement details; Figure S1: The ^1H NMR spectrum (500 MHz, C₆D₆) of the reaction of compound 1 with H₂O; Figure S2: The ^{11}B NMR spectrum (160 MHz, C₆D₆) of the reaction of compound 1 with H₂O; Figure S3: The $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum (126 MHz, C₆D₆) of the reaction of compound 1 with H₂O; Figure S4: The ^1H NMR spectrum (500 MHz, C₆D₆) of the thermolysis of compound 1; Figure S5: The ^{11}B NMR spectrum (160 MHz, C₆D₆) of the thermolysis of compound 1; Figure S6: The ^1H NMR spectrum (500 MHz, C₆D₆) of the reaction of compound 1 with phenyl acetylene; Figure S7: The ^{11}B NMR spectrum (160 MHz, C₆D₆) of the reaction of compound 1 with phenyl acetylene; Figure S8: The ^1H NMR spectrum (500 MHz, C₆D₆) of compound 1 with H₂ present in solution showing no reaction; Figure S9: The ^1H NMR spectrum (500 MHz, d_8 -THF) of compound 2; Figure S10: The $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum (126 MHz, d_8 -THF) of compound 2; Figure S11: The ^1H NMR spectrum (500 MHz, C₆D₆) of compound 3; Figure S12: The $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum (126 MHz, C₆D₆) of compound 3; and Figure S13: The IR spectrum of compound 3.

Author Contributions: Conceptualization, D.J.L.; data curation, R.S.C.C., T.M.H.D.; formal analysis, M.F.M.; funding acquisition, D.J.L.; investigation, R.S.C.C., T.M.H.D., T.H.J.; methodology, D.J.L.; project administration, D.J.L.; resources, R.S.C.C., T.M.H.D.; supervision, D.J.L., M.F.M.; visualization, D.J.L., M.F.M.; writing—original draft, T.M.H.D., R.S.C.C., D.J.L.; writing—review & editing, R.S.C.C., M.F.M., D.J.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Royal Society, grant number URF\R1\191066. The APC was funded by the kind contribution of MDPI.

Data Availability Statement: X-ray crystallography data can be found through the CCDC (2201279–2201281 for **1**·H₂O, **2** and **3**, respectively). The poor-quality data for **1** precluded submission to the CCDC and data are available from the authors on request. All other data can be found in the Supplementary Materials.

Acknowledgments: D.J.L. thanks the Royal Society for the support of a University Research Fellowship. We wish to thank the University of Bath and MC² for use of their analysis facilities.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Stephan, D.W.; Erker, G. Frustrated Lewis Pair Chemistry: Development and Perspectives. *Angew. Chem. Int. Ed.* **2015**, *54*, 6400–6441. [[CrossRef](#)] [[PubMed](#)]
- Stephan, D.W. Frustrated Lewis Pairs. *J. Am. Chem. Soc.* **2015**, *137*, 10018–10032. [[CrossRef](#)]
- Xu, M.; Jupp, A.R.; Qu, Z.-W.; Stephan, D.W. Alkali Metal Species in the Reversible Activation of H₂. *Angew. Chem. Int. Ed.* **2018**, *57*, 11050–11054. [[CrossRef](#)] [[PubMed](#)]
- Welch, G.C.; Juan, R.R.S.; Masuda, J.D.; Stephan, D.W. Reversible, Metal-Free Hydrogen Activation. *Science* **2006**, *314*, 1124–1126. [[CrossRef](#)]
- Stephan, D.W. Diverse Uses of the Reaction of Frustrated Lewis Pair (FLP) with Hydrogen. *J. Am. Chem. Soc.* **2021**, *143*, 20002–20014. [[CrossRef](#)]
- Scott, D.J.; Simmons, T.R.; Lawrence, E.J.; Wildgoose, G.G.; Fuchter, M.J.; Ashley, A.E. Facile Protocol for Water-Tolerant “Frustrated Lewis Pair”-Catalyzed Hydrogenation. *ACS Catal.* **2015**, *5*, 5540–5544. [[CrossRef](#)] [[PubMed](#)]
- Dureen, M.A.; Stephan, D.W. Terminal Alkyne Activation by Frustrated and Classical Lewis Acid/Phosphine Pairs. *J. Am. Chem. Soc.* **2009**, *131*, 8396–8397. [[CrossRef](#)] [[PubMed](#)]
- Stephan, D.W.; Erker, G. Frustrated Lewis pair chemistry of carbon, nitrogen and sulfur oxides. *Chem. Sci.* **2014**, *5*, 2625–2641. [[CrossRef](#)]
- Mömming, C.M.; Otten, E.; Kehr, G.; Fröhlich, R.; Grimme, S.; Stephan, D.W.; Erker, G. Reversible Metal-Free Carbon Dioxide Binding by Frustrated Lewis Pairs. *Angew. Chem. Int. Ed.* **2009**, *48*, 6643–6646. [[CrossRef](#)] [[PubMed](#)]
- Barnett, B.R.; Moore, C.E.; Rheingold, A.L.; Figueroa, J.S. Cooperative Transition Metal/Lewis Acid Bond-Activation Reactions by a Bidentate (Boryl)iminomethane Complex: A Significant Metal–Borane Interaction Promoted by a Small Bite-Angle LZ Chelate. *J. Am. Chem. Soc.* **2014**, *136*, 10262–10265. [[CrossRef](#)] [[PubMed](#)]
- Barnett, B.R.; Moore, C.E.; Rheingold, A.L.; Figueroa, J.S. Frustrated Lewis pair behavior of monomeric (boryl)iminomethanes accessed from isocyanide 1,1-hydroboration. *Chem. Commun.* **2015**, *51*, 541–544. [[CrossRef](#)] [[PubMed](#)]
- Li, J.; Daniliuc, C.G.; Kehr, G.; Erker, G. Three-Component Reaction to 1,4,2-Diazaborole-Type Heteroarene Systems. *Angew. Chem. Int. Ed.* **2021**, *60*, 27053–27061. [[CrossRef](#)] [[PubMed](#)]
- Horsley Downie, T.M.; Charman, R.S.C.; Hall, J.W.; Mahon, M.F.; Lowe, J.P.; Liptrot, D.J. A stable ring-expanded NHC-supported copper boryl and its reactivity towards heterocumulenes. *Dalton Trans.* **2021**, *50*, 16336–16342. [[CrossRef](#)] [[PubMed](#)]
- Procter, R.J.; Dunsford, J.J.; Rushworth, P.J.; Hulcoop, D.G.; Layfield, R.A.; Ingleson, M.J. A Zinc Catalyzed C(sp³)–C(sp²) Suzuki–Miyaura Cross-Coupling Reaction Mediated by Aryl-Zincates. *Chem. Eur. J.* **2017**, *23*, 15889–15893. [[CrossRef](#)] [[PubMed](#)]
- Fortman, G.C.; Slawin, A.M.Z.; Nolan, S.P. A Versatile Cuprous Synthone: [Cu(IPr)(OH)] (IPr = 1,3 bis(diisopropylphenyl)imidazol-2-ylidene). *Organometallics* **2010**, *29*, 3966–3972. [[CrossRef](#)]
- Plotzitzka, J.; Kleeberg, C. [(NHC)CuI–ER₃] Complexes (ER₃ = SiMe₂Ph, SiPh₃, SnMe₃): From Linear, Mononuclear Complexes to Polynuclear Complexes with Ultrashort CuI⋯CuI Distances. *Inorg. Chem.* **2016**, *55*, 4813–4823. [[CrossRef](#)] [[PubMed](#)]
- Jones, C.; Mills, D.P.; Rose, R.P.; Stasch, A.; Woodul, W.D. Synthesis and further reactivity studies of some transition metal gallyl complexes. *J. Organomet. Chem.* **2010**, *695*, 2410–2417. [[CrossRef](#)]
- Dolomanov, O.V.; Bourhis, L.J.; Gildea, R.J.; Howard, J.A.K.; Puschmann, H. OLEX2: A complete structure solution, refinement and analysis program. *J. Appl. Crystallogr.* **2009**, *42*, 339–341. [[CrossRef](#)]
- Sheldrick, G. SHELXT—Integrated space-group and crystal-structure determination. *Acta Crystallogr. Sect. A* **2015**, *71*, 3–8. [[CrossRef](#)] [[PubMed](#)]
- Jordan, A.J.; Wyss, C.M.; Bacsá, J.; Sadighi, J.P. Synthesis and Reactivity of New Copper(I) Hydride Dimers. *Organometallics* **2016**, *35*, 613–616. [[CrossRef](#)]