

RESEARCH ARTICLE

CRISPR/Cas9-mediated gene deletion of the *ompA* gene in symbiotic *Cedecea neteri* impairs biofilm formation and reduces gut colonization of *Aedes aegypti* mosquitoes

Shivanand Hegde¹, Pornjarim Nilyanimit², Elena Kozlova³, Enyia R. Anderson¹, Hema P. Narra³, Sanjeev K. Sahni³, Eva Heinz⁴, Grant L. Hughes^{1*}

1 Departments of Vector Biology and Tropical Disease Biology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom, **2** Center of Excellence in Clinical Virology, Chulalongkorn University, Bangkok, Thailand, **3** Department of Pathology, University of Texas Medical Branch, Galveston, Texas, United States of America, **4** Department of Vector Biology and Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, United Kingdom

* grant.hughes@lstm.ac.uk



OPEN ACCESS

Citation: Hegde S, Nilyanimit P, Kozlova E, Anderson ER, Narra HP, Sahni SK, et al. (2019) CRISPR/Cas9-mediated gene deletion of the *ompA* gene in symbiotic *Cedecea neteri* impairs biofilm formation and reduces gut colonization of *Aedes aegypti* mosquitoes. *PLoS Negl Trop Dis* 13(12): e0007883. <https://doi.org/10.1371/journal.pntd.0007883>

Editor: Mariangela Bonizzoni, Università degli Studi di Pavia, ITALY

Received: January 31, 2019

Accepted: October 26, 2019

Published: December 2, 2019

Copyright: © 2019 Hegde et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: GLH was supported by NIH grants (R21AI138074, R21AI124452 and R21AI129507), a Royal Society Wolfson Fellowship, the Western Gulf Center of Excellence for Vector-borne Diseases (CDC grant CK17-005), the Robert J. and Helen Kleberg Foundation and the Gulf Coast

Abstract

Background

Symbiotic bacteria are pervasive in mosquitoes and their presence can influence many host phenotypes that affect vectoral capacity. While it is evident that environmental and host genetic factors contribute in shaping the microbiome of mosquitoes, we have a poor understanding regarding how bacterial genetics affects colonization of the mosquito gut. The CRISPR/Cas9 gene editing system is a powerful tool to alter bacterial genomes facilitating investigations into host-microbe interactions but has yet to be applied to insect symbionts.

Methodology/Principal findings

To investigate the role of bacterial genetic factors in mosquito biology and in colonization of mosquitoes we used CRISPR/Cas9 gene editing system to mutate the outer membrane protein A (*ompA*) gene of a *Cedecea neteri* symbiont isolated from *Aedes* mosquitoes. The *ompA* mutant had an impaired ability to form biofilms and poorly infected *Ae. aegypti* when reared in a mono-association under gnotobiotic conditions. In adult mosquitoes, the mutant had a significantly reduced infection prevalence compared to the wild type or complement strains, while no differences in prevalence were seen in larvae, suggesting genetic factors are particularly important for adult gut colonization. We also used the CRISPR/Cas9 system to integrate genes (antibiotic resistance and fluorescent markers) into the symbionts genome and demonstrated that these genes were functional *in vitro* and *in vivo*.

Conclusions/Significance

Our results shed insights into the role of *ompA* gene in host-microbe interactions in *Ae. aegypti* and confirm that CRISPR/Cas9 gene editing can be employed for genetic manipulation of non-model gut microbes. The ability to use this technology for site-specific integration

Consortia. This work was also supported by a James W. McLaughlin postdoctoral fellowship at the University of Texas Medical Branch to SH and a Dutsadi Piphat Scholarship and Thailand Research Chair Grant from NSTDA (p-15-50004) to PN. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

of genes into the symbiont will facilitate the development of paratransgenic control strategies to interfere with arboviral pathogens such Chikungunya, dengue, Zika and Yellow fever viruses transmitted by *Aedes* mosquitoes.

Author summary

Microbiota profoundly affect their host but few studies have investigated the role of bacterial genetics in host-microbe interactions in mosquitoes. Here we applied the CRISPR/Cas9 gene editing system to knockout a membrane protein in *Cedecea neteri*, which is a dominant member of the mosquito microbiome. The mutant strain had an impaired capacity to form biofilms, infected larvae and adults at lower titers, and had a reduced prevalence in adults. The lower prevalence in adults, but not larvae, likely reflects the difference in the modes of bacterial acquisition from the larval water of these two life stages. Importantly from an applied perspective, we also demonstrated that this editing technology can be harnessed for site-specific integration of genes into the bacterial chromosome. In proof-of-principle studies we integrated either a fluorescent protein or gene conferring antibiotic resistance into the bacterial genome and showed these transgenes were functional in mosquitoes. The specificity, flexibility, and simplicity of this editing approach in non-model bacteria will be useful for developing novel symbiotic control strategies to mitigate the burden of arthropod-borne disease.

Introduction

Mosquitoes harbor a community of microbes within their guts. In general, the gut-associated microbiome of mosquitoes tends to have low species richness but can differ greatly between individuals and habitats [1–8]. Importantly, these microbes can modulate many host phenotypes, several of which can influence vectorial capacity [9–11]. As such, it is imperative that we understand how the microbiome is acquired and maintained within mosquito vectors. While environmental factors unquestionably influence the mosquito microbiome composition and abundance [2–4, 8], studies are elucidating the role of microbial interactions [5, 7, 12, 13] and host genetic factors [14–18] in shaping the microbiome. However, we have a poor understanding of bacterial factors that influence colonization of the mosquito gut and this is likely an underappreciated force influencing host-microbe interactions in mosquitoes.

In other invertebrates, several bacterial genes have been implicated in gut colonization. For example, a genome wide screen exploiting transposon-sequencing found a suite of genes from the bacterium *Snodgrassella alvi* involved in colonization of the honey bee gut [19]. These bacterial genes were classified into the broad categories of extracellular interactions, metabolism, and stress response [19]. Knockout of a purine biosynthesis gene in *Burkholderia* impaired biofilm formation and reduced bacterial colonization rates in a bean bug [20]. Biofilm formation was also shown to play a role in virulence of pathogenic *Pseudomonas* in artificial infections of *Drosophila*, with strains that lacked the capacity to form biofilms being more virulent to the host, although a hyperbiofilm strain was less virulent than the wild type (WT) strain [21]. In other blood feeding invertebrates, bacterial genetics also appears critical for host colonization. Knockout of the type II secretion system in *Aeromonas veronii* reduced infection in *Hirudo verbena* leeches [22]. In tsetse flies, the outer-membrane protein A (*ompA*) gene of *Sodalis glossinidius* is essential for symbiotic interactions [23]. *Sodalis* mutants lacking the

ompA gene poorly colonized the fly gut compared to the WT symbionts [23], likely due to the mutant strains reduced capacity to form biofilms [24]. Heterologous expression of the *ompA* gene from pathogenic *Escherichia coli* in *Sodalis* mutants induced mortality in the fly implicating this gene as a virulence factor in pathogenic bacteria [23]. Taken together, these studies suggest that bacterial genetic factors are critical for host colonization of invertebrates and that biofilm formation facilitates symbiotic associations in insects.

In mosquitoes, few studies have investigated how bacterial genetics affect gut colonization. However, evidence from experimental evolution studies suggests bacterial genetics plays a critical role. In two separate studies, *Enterobacter* was selected for increased persistence in the gut of *Anopheles gambiae* mosquitoes, the major malaria vector in sub-Saharan Africa, by repeatedly infecting mosquitoes with strains that persisted in the gut for longer periods of time [25, 26]. Transcriptomics comparisons of effective and ineffective colonizers in liquid media identified 41 genes that were differentially expressed between these two strains [26], further implicating the importance of bacterial genetics in mosquito infection, however the role of these genes in colonization of the mosquito gut has not been resolved. In a separate study, *in vitro* screening of a transposon mutant library of *Enterobacter* identified a *waaL* gene mutant that was insensitive to oxidative stress [27]. The *waaL* gene encodes an O antigen ligase which is needed for attachment of the O antigen to lipopolysaccharide. The mutant was found to have lower rates of colonization of the midguts of *Anopheles* mosquitoes [27].

Gene knockouts approaches in bacteria provide compelling evidence of the role of bacterial genes in host-microbe interactions [22–24, 27–29]. In general, most studies use transposon mutagenesis for gene knockout, which requires screening of the mutant library. A targeted gene knockout approach is highly desirable to investigate the functionality of bacterial genes in host-microbe interactions. In the past few years, the CRISPR/Cas9 gene editing system has been employed to modify bacterial genomes [30–32]. While much of the work has been done in model bacterial species [31–37], editing approaches have expanded into non-model bacterial systems [38–43]. Despite this expansion, the approach has been used less frequently for host-associated microbes [39, 44], and rarely for arthropod symbionts. In the vector biology field, gene knockout approaches can be used to interrogate the role of bacterial genes responsible for host-microbe interactions, whilst the ability to integrate genes into the bacterial symbiont genome has great potential for applied paratransgenic control strategies [10, 45–47]. To date, manipulation of non-model symbionts that associate with insect vectors has been accomplished by plasmid transformation [48–55] or stable transformation of the genome using transposons or integrative plasmids [56–63], but the use of CRISPR/Cas9 gene editing in insect gut symbionts has yet to be accomplished. For paratransgenic strategies, stable site-specific integration of transgenes into the symbiont genome is critical. Therefore, the application of CRISPR/Cas9 gene editing technology to non-model bacteria that associate with insect vectors will stimulate research in this field.

We therefore undertook studies to develop CRISPR/Cas9 genome editing approaches in *Cedecea neteri* isolated from *Aedes* mosquitoes. We used the Scarless Cas9 Assisted Recombinering (no-SCAR) method to disrupt the *ompA* gene of the non-model *C. neteri* [35]. After characterization of the mutant *in vitro*, we examined the role of the *ompA* gene in host-microbe interactions by re-infecting bacteria into mosquitoes in a mono-association. To demonstrate that the CRISPR/Cas9 gene-editing system could be useful for applied symbiotic control approaches we inserted genes conferring antibiotic resistance or a fluorescent protein into the bacterial genome and re-infected the altered strains back into mosquitoes. Our result sheds insights into the role of the *ompA* gene in host-microbe interactions in *Ae. aegypti* and confirm that CRISPR/Cas9 gene editing can be a powerful tool for genetic manipulation of native gut-associated microbes of mosquitoes.

Results

C. neteri biofilm formation in *Ae. aegypti* guts

Over the course of conducting mono-axenic infections in *Ae. aegypti* mosquitoes with a *Cedecea* symbiont, we repeatedly observed a conglomeration of bacterial cells in the anterior and posterior midgut (Fig 1, S1 Fig) that had a similar appearance to biofilms observed in the guts of other insects [21, 24]. We also infected mosquitoes with the *E. coli* BL21(DE3) lab strain as a control, but we did not see any evidence of infection (Fig 1, S1D–S1F Fig) although infection with this bacterium enabled mosquito development [64]. The *E. coli* BL21(DE3) lab strain does not have the capacity to form biofilms [65], possibly explaining its inability to infect mosquitoes. We therefore set out to examine the role of bacterial genetics in biofilm formation and host colonization of gut-associated bacteria of *Aedes* mosquitoes. We used multilocus sequence typing (MLST) to confirm the species of our isolate, which indicated the bacterium was *C. neteri* (S2 Fig). Several genes have been implicated in biofilm formation [21, 24], but we chose to knockout the *ompA* gene of *C. neteri* given that this gene has been demonstrated to influence biofilm formation and gut colonization of *Sodalis* [23, 24], an *Enterobacteriaceae* symbiont of tsetse flies. We used the CRISPR/Cas9 genome editing system to mutate the symbiont genome to demonstrate this approach could be employed for non-model symbiotic bacteria that associate with mosquitoes.

Genome editing in *C. neteri* bacteria isolated from mosquitoes

To edit the *Cedecea* isolate that resides within the gut of *Aedes* mosquitoes, we employed the no-SCAR gene editing approach that had been developed in *E. coli* [35]. To optimize the approach in our hands, we performed initial experiments in *E. coli* to delete a ~1 kb region of the *ompA* gene (Fig 2A). As the no-SCAR approach exploits the λ -Red recombineering system to repair double stranded breaks, we transformed bacteria with a double stranded DNA template that had regions of homology flanking the gRNA site (250 bp for each arm). Using this approach, we successfully deleted a 1001 bp fragment of the *ompA* gene. From the colonies we screened, we saw an editing at a frequency of 6.25% (N = 48) (Fig 2A). For *C. neteri*, we altered




Fig 1. Midgut infection of *C. neteri* and *E. coli* in mono-associations of *Aedes* mosquitoes. *C. neteri* forms a biofilm in the gut of 3–4 day old *Ae. aegypti* adult mosquitoes (left) while no bacteria were observed in the gut of mosquitoes reared with *E. coli* under gnotobiotic conditions (right). Bacteria possessed the pRAM-mCherry plasmid, which expresses the mCherry fluorescent protein and conferred resistance to kanamycin. Blue—host nuclei stained by DAPI. Green—host actin cytoskeleton stained with phalloidin. The scale bar is 70 μ m.

<https://doi.org/10.1371/journal.pntd.0007883.g001>




Fig 2. CRISPR/Cas9 genome editing in bacteria. A schematic of the editing approach and screening of putative mutants in (A) *E. coli* and (B) *C. neteri*. A ~1kb fragment of *E. coli* BL21 (DE3) was deleted using no-SCAR protocol. The 250 bp of the left arm (LA) and right arm (RA) was assembled to generate the 500 bp donor DNA. The transformants were screened via colony PCR with primers binding in regions flanking the deletion. Similar to the strategy employed in *E. coli*, the knockout of the *ompA* gene from *C. neteri* isolated from the mosquito gut was created by deleting the 598 bp fragment. The grey area indicates the PAM site in the *ompA* gene and arrow shows the cleavage site in the genome. (C) The sequence of the *ompA* mutation in *E. coli* and *C. neteri* was confirmed by Sanger sequencing. The sequence above the gene within the dotted line has been deleted. The chromatogram shows the 10 bp flanking the deletion.

<https://doi.org/10.1371/journal.pntd.0007883.g002>

our editing procedure to delete a 598 bp fragment from the *ompA* gene. This was done to enhance the efficiency of obtaining mutants [35] and accommodate the PAM site which was at a different location in the *ompA* gene in *C. neteri*. Using a donor template designed for the *C. neteri ompA* gene that had flanking homology arms of similar length as the previous experiment done in *E. coli*, we obtained mutant knockouts at a rate of 32% (N = 50) (Fig 2B). For both bacterial species, Sanger sequencing across the integration site indicated the deletion occurred at the expected loci in the bacterial genome (Fig 2C; S1 Appendix).

Characterization of the *C. neteri ompA* mutant

We quantified the growth rates of the $\Delta ompA$ mutant in comparison to the WT *C. neteri* and the $\Delta ompA/ompA$ complement in liquid LB media. We saw minimal differences in the growth between the WT, the $\Delta ompA$ mutant or the $\Delta ompA/ompA$ complement (Fig 3A). To examine the stability of the deletion, we subcultured the $\Delta ompA$ mutant on LB media for 10 generations and performed PCR to amplify across the deletion. At alternative generations, PCR analysis indicated the deletion was present indicating genomic stability at this site (Fig 3B).




Fig 3. *In vitro* characterization of the *ompA* mutation. (A) The *C. neteri* $\Delta ompA$ mutant had a similar growth rate compared to both the WT and the $\Delta ompA/ompA$ complement in liquid LB media. Five technical replicates were used to create growth curves. (B) The stability of mutant was evaluated *in vitro* by continuous subculturing in LB media. Genomic DNA from alternative subcultures was used as template for PCR using primers that amplified across the deletion. The stability assay was repeated twice. Two separate gel images were merged to create figure 3B (passage 8 was run on a separate gel to passages 0–6). (C) Biofilm formation was assessed using the CV biofilm assay for the WT, $\Delta ompA$ mutant and the $\Delta ompA/ompA$ complement. Two biological replicates were completed. (D) Quantification of the relative biofilm formation normalized by the number of bacteria per well (N = 3). Error bars represent standard error. The assay was repeated twice.

<https://doi.org/10.1371/journal.pntd.0007883.g003>

Previously, *ompA* has been shown to be important in biofilm formation as *Sodalis* deletion mutants were unable to form biofilms [24]. Therefore, we characterized *in vitro* biofilm formation using the crystal violet (CV) biofilm assay. From visual inspection, it was clear the $\Delta ompA$ mutant had distinctly less biofilm deposition compared to either the WT or the $\Delta ompA/ompA$ complement (Fig 3C). After quantification and normalization to account for any difference in growth between the strains, biofilm formation was confirmed to be significantly different between the $\Delta ompA$ mutant and the WT or complement (Fig 3D; Tukey’s multiple comparisons test, $P < 0.0001$). There was no significant difference between the WT and the $\Delta ompA/ompA$ complement (Tukey’s multiple comparisons test $P = 0.2$).

The role of *ompA* gene in mosquito infection

To examine the importance of the *ompA* gene on bacterial colonization of mosquitoes, we infected *Ae. aegypti* mosquitoes in a mono-association under gnotobiotic conditions [64]. This infection method was used to avoid other gut-associated microbes influencing host colonization rates [7] and it also enabled straightforward quantification of introduced bacteria by measuring colony forming units (CFUs). No significant changes were seen in the prevalence of infection (number of mosquitoes infected) in the larval stage (Fig 4A, Fisher’s exact test; WT




Fig 4. The $\Delta ompA$ mutant poorly infected mosquitoes. Infection of *C. neteri* strains (WT, $\Delta ompA$ mutant and $\Delta ompA/ompA$ complement; the former two possessed the pRAM-Cherry plasmid while the latter possessed the pRAM-Cherry-Ent-*OmpA* plasmid) reared in a mono-association using a gnotobiotic rearing approach for larvae (A and C) and adults (B and D). L4 larvae and 3–4 days post emergence adults were screened for bacterial load by plating on selective LB media with kanamycin to quantify the bacteria. The prevalence of infection (number of mosquitoes infected) between the treatments was calculated comparing the number of infected to uninfected larvae (A) or adults (B). Density of bacteria (CFU/mosquito) in larvae (C) and adults (D). The assay was repeated twice. Results display pooled data from each independent replicate. Box and whiskers show the median, the 25th and 75th percentiles and the minimum and maximum values.

<https://doi.org/10.1371/journal.pntd.0007883.g004>

compared to $\Delta ompA$ $P = 0.24$ and $\Delta ompA$ compared to $\Delta ompA/ompA$ $P = 0.24$) with rates of infection consistently high (WT 100%, $\Delta ompA$ 96%, and $\Delta ompA/ompA$ 100%). In adults, the prevalence of infection was significantly different (Fig 4B, Fisher’s exact test; WT compared to $\Delta ompA$ $P < 0.0001$ and $\Delta ompA$ compared to $\Delta ompA/ompA$ $P < 0.0001$), with only 45% of




Fig 5. The $\Delta ompA$ mutant does not affect growth rates or development of mosquitoes. The growth rate (time to pupation) (A) and development (percentage of L1 larvae to reach adulthood) (B) was observed in mosquitoes infected with *C. neteri* strains (WT, $\Delta ompA$ mutant and $\Delta ompA/ompA$ complement) reared in a mono-association. The experiment was done twice with a minimum of 15 individuals. Sample size for panel A indicates number of individuals, while for B indicates the number of replicate flasks. Each flask has 20 mosquitoes.

<https://doi.org/10.1371/journal.pntd.0007883.g005>

adults infected by the $\Delta ompA$ mutant compared to 95% and 88% by the WT and $\Delta ompA/ompA$ complement, respectively. In larvae, we saw a significant reduction in bacterial titer in the mutant compared to both the WT (Kruskal-Wallis test with Dunn's test; $P < 0.05$) and the $\Delta ompA/ompA$ complement (Kruskal-Wallis test with Dunn's test; $P < 0.05$) (Fig 4C) with median values of 1.5×10^5 , 2.3×10^4 , and 1.5×10^5 for the WT, $\Delta ompA$, and $\Delta ompA/ompA$ complement respectively. Similarly, in adults, there was a significant reduction in bacterial infection in the $\Delta ompA$ mutant compared to either the WT or $\Delta ompA/ompA$ complement (Kruskal-Wallis test with Dunn's test; $P < 0.001$) (Fig 4D), with median value of 8.1×10^2 , 0, and 7.5×10^2 for the WT, $\Delta ompA$, and $\Delta ompA/ompA$ complement respectively. However, when considering only the infected mosquitoes for analysis, we saw no significant difference between the treatments (S3 Fig, Kruskal-Wallis test with Dunn's test; $P > 0.99$). For both the larvae and adult density quantifications, the non-parametric test (Kruskal-Wallis test) was used due to non-normal distribution of data (Shapiro-Wilks test; $P < 0.001$). We also monitored the growth rates of mosquitoes administered with the WT, $\Delta ompA$ mutant and $\Delta ompA/ompA$ complement. No significant differences were seen in the time to pupation (Fig 5A) or percentage of first instar larvae that reached adulthood (Fig 5B) between any of the bacterial strains.

Integration of genes into the *C. neteri* chromosome

We undertook experiments to demonstrate the CRISPR/Cas9 gene-editing approaches can be used to integrate genes into the chromosome of non-model bacteria that associate with mosquitoes. We created two independent transgenic strains that had either a gene encoding mCherry fluorescence or a gene encoding resistance to the antibiotic gentamicin inserted into the bacterial chromosome. Before undertaking these integration experiments we confirmed that *C. neteri* was susceptible to gentamicin. These genes were integrated into the genome using the same gRNA that was used for deletional mutagenesis (S1 Table), and as such, these

insertions also disrupted the *ompA* gene. Sequencing across the integration site indicated the insertion of these genes occurred within the *ompA* gene and thereby disrupted its function (Fig 6A and 6B). Continual subculturing was undertaken for both strains and molecular analysis indicated the stability of these lines for ten generations (Fig 6C and 6D). Expression of mCherry fluorescence and growth of the $\Delta ompA::gentamicin$ strain on media containing gentamicin demonstrated the integrated genes were functional *in vitro* (Fig 6E and 6F).

To examine the functionality of the integrated genes in the mosquito we administered either WT, $\Delta ompA::mCherry$, or $\Delta ompA::gentamicin$ to conventionally reared 3–4 day old adult female *Ae. aegypti* in a sugar meal for 3 days or larvae cleared of their microbiota. For gnotobiotic infection we used bacteria expressing mCherry from a plasmid. The dissected gut from 3–4 day old adults showed a higher percentage of WT bacteria compared to either of the integrated mutants. After screening midgut samples from each treatment, we found that mosquitoes infected with WT bacteria had the highest infection prevalence (69%) and that the mCherry and gentamicin knockin mutants were found only in 4% and 33% of the samples, respectively (S4 Fig, S4 Table). In addition, biofilms were seen mainly in mosquitoes infected with WT bacteria (31%) whilst midguts infected with mutants had few or no biofilms (0–2%) (S4 Fig, S4 Table). In sugar fed adult mosquitoes, $\Delta ompA::mCherry$ bacteria were observed in the gut of mosquitoes with a distinct punctate distribution, whereas no signal was seen in auto-fluorescence controls (WT *C. neteri* infected mosquitoes) (Fig 6G). The *C. neteri ompA::gentamicin* was successfully rescued from mosquitoes reared on gentamicin and stably infected mosquitoes over time at a density of approximately 1×10^4 CFUs/mosquito. Consistent with our previous result (Fig 4B), WT bacteria initially infected mosquitoes at higher titers compared to the mutant (T test; day 0 $P < 0.001$). However, after 4 days rearing on antibiotic the total bacterial load in mosquitoes administered WT *C. neteri* was significantly reduced compared to the $\Delta ompA::gentamicin$ (T test; day 4 $P < 0.05$) while the prevalence of mosquitoes with culturable microbiota was reduced to 80%. After 6 days rearing on antibiotic, the $\Delta ompA::gentamicin$ density was significantly elevated compared to the WT (T test; day 6 $P < 0.001$) only one mosquito was infected, which had a low density infection (10 CFUs/mosquito) (Fig 6H).

Discussion

We harnessed the CRISPR/Cas9 gene editing system to create knockout mutants in a *C. neteri* gut symbiont of *Aedes* mosquitoes to examine the role of bacterial genetics in biofilm formation and gut colonization. A deletion of the *ompA* gene of *C. neteri* decreased bacterial colonization of mosquitoes after infection in a mono-association. Strikingly, we found this effect was most pronounced in adult mosquitoes with more than half of the mosquitoes not possessing any culturable mutants, whereas there was no difference in prevalence of infection between the mutant and WT bacteria in larvae. The reduced prevalence of mutant bacteria in adults likely reflects differences in microbial colonization of each mosquito life stage. Larvae are continually subjected to bacteria in the larval water habitat while adults only have a short time frame to acquire bacteria from the aquatic environment immediately after eclosion. Alternatively, the reduced prevalence in adults could be due an impaired ability of mutant bacteria to be transstadially transmitted. Several bacterial species have been shown to exploit this process to transfer between life stages [66–69]. When only analysing adult mosquitoes where bacteria did colonize the host, we saw no differences in the density of the mutant strain compared to the WT or complement, suggesting that *ompA* is acting at the colonization stage but has minimal effect on post-colonization processes. However, when examining midguts using fluorescent microscopy, in general, we observed reduced loads of the mutant strains. When




Fig 6. Integration of mCherry and gentamicin into the *C. neteri* genome. Sanger sequence across the integration site, stability of the inserted gene and *in vitro* expression of the inserted gene for the $\Delta ompA::mCherry$ (A-C) and the $\Delta ompA::gentamicin$ (B-D) strains. The chromatogram shows the sequence spanning the inserted sites. Strains were continually subcultured for 10 passages and PCR was done to examine the stability of the insert (C; $\Delta ompA::mCherry$ plus WT, D; $\Delta ompA::gentamicin$ passaged with (ab+) or without (ab-) gentamicin plus WT). mCherry fluorescence (E) or ability to grow on selective media containing gentamicin (F) confirmed the expression of the transgene *in vitro*. Mosquitoes were inoculated with the *C. neteri* strains to confirm expression of the transgene *in vivo*. Dissected midgut infected with $\Delta ompA::mCherry$ (left) or negative control (right; WT bacteria without expression plasmid) (G). Midguts were stained with phalloidin (green) and DAPI (blue). The WT and $\Delta ompA::gentamicin$ *C. neteri* strains were fed to adult mosquitoes for 3 days in a sugar meal before gentamicin was administered to mosquitoes in sugar without bacteria (H). Mosquitoes were collected every second day and CFUs assessed. Pairwise comparisons were conducted at each time point using a T test.

<https://doi.org/10.1371/journal.pntd.0007883.g006>

quantifying bacterial load by CFU we used whole mosquitoes. It may be possible that mutant bacteria were residing in other tissues in the adult but poorly re-infected the midgut. If this occurred, it would indicate involvement of *ompA* in transstadial transmission. The greater variability seen in the prevalence of adults compared to the larval is consistent with other sequence-based studies that indicate adult stages have greater variability in species composition of their microbiota, whereas the microbiome of immature stages is similar to the microbiota in larval water habitat [2–5, 8, 70].

Mutant bacteria colonized mosquitoes at higher densities when administered to adults as opposed to larvae. There are several possible explanations for this finding. The first relates to the method of inoculation with adults being administered bacteria in a sugar meal while larvae were exposed to bacteria in their aquatic environment. The different inoculation process itself may influence titer but also when sugar feeding, adults had the opportunity for repeated infections whereas emerging adults only had a narrow window for inoculation as they did not have further access to the larval water habitat after eclosion. The second explanation relates to differences in the microbiome of these mosquitoes. The mosquitoes inoculated as adult were reared conventionally, and as such, had an intact microbiome, while larvae reared in the gnotobiotic system only possessed the individual *Cedecea* strains that were administered. For the latter group there was no opportunity for the native WT bacteria (either of the same or different species) to rescue the mutant phenotype. In the *Sodalis-tsetse* system, mutant bacteria were capable of infecting flies that had an intact microbiome but were unable to infect *Sodalis*-free *tsetse* flies [23], suggesting WT *Sodalis* facilitated colonization of the mutant strain. In mono-axenic infections, the *C. neteri* mutant strain was able to infect *Ae. aegypti*, indicating that *ompA* is not essential for infection in the mosquito-*Cedecea* system.

Our results, in conjunction with studies in the *Sodalis-tsetse* system [23, 24], suggests that biofilm formation may be a strategy employed by bacteria to colonize the gut of insects. In pathogenic infections in mammals, biofilms enable bacteria to colonize new niches, promote infection, and are associated with virulence [71]. Although less is known regarding the importance of biofilm formation in insects, in an artificial *Pseudomonas-Drosophila* infection model, biofilm formation was associated with virulence and host survival [21]. In a natural symbiotic association between bean bugs and *Burkholderia*, disruption of a purine biosynthesis gene in the bacterium reduce biofilm formation and colonization of the insect [20]. In mosquitoes, gut biofilm formation could also have implications for vector competence. *Chromobacterium*, which was isolated from *Aedes* mosquitoes, produced molecules that inhibited dengue virus only when grown *in vitro* as a biofilm but not when grown in a planktonic state [72], however it is unknown if biofilm formation occurred *in vivo* in the mosquito. Our data provide evidence that biofilms occur within the gut of mosquitoes and facilitate host colonization.

Although we have shown that the *ompA* gene of *C. neteri* is important for host colonization, we see no evidence that deletion of this gene alters mosquito development or growth rates. This is in contrast to the *Riptortus-Burkholderia* symbiosis whereby mutation of the *purT* gene in *Burkholderia* resulted in reduced growth rates and reduction in body weight of the host compared to insects that were infected with the WT bacterium [20]. The difference in our study to the findings in the *Riptortus-Burkholderia* symbiosis could be related to different requirements of the bean bug compared to the mosquito host as well as the different genes mutated in the symbionts. Our findings are consistent with a previous study in *Ae. aegypti* whereby an *ompA* mutant of *E. coli* did not influence growth when reared in a mono-association [73]. Using a similar gnotobiotic system that exploits the ability to sterilize mosquito eggs and rescue development by nutritional supplementation, several recent reports describe approaches to create bacteria-free mosquitoes [73, 74]. Here, we reared mosquitoes in a mono-association where they were only subjected to *C. neteri*. However, more than half the

adult mosquitoes inoculated with the $\Delta ompA$ mutant were not infected by bacteria, as evidenced by the inability to culture bacteria from these insects. Nevertheless, these mosquitoes had similar development and growth rates compared to mosquito possessing WT bacteria. The use of mutant bacteria that rescue development but have an impaired ability to colonize mosquitoes may provide a simple means to create axenic adult mosquitoes.

CRISPR/Cas9 gene editing has revolutionized genetic approaches in model and non-model bacteria [31–43]. However, there has been limited use of this technology in symbiotic microbes of arthropods. Here we demonstrate that editing approaches functional in *E. coli* can be easily applied with minimal adaptation to phylogenetically related symbiotic bacteria that are found within the guts of mosquitoes. The application of CRISPR/Cas9 genome editing to gut-associated bacteria of mosquitoes has significant applied potential. Paratransgenesis strategies are being evaluated in a range of medical and agricultural systems to mitigate pathogen transmission from insect vectors, however, most approaches engineer symbionts by plasmid transformation [49–55, 75] and where genome integration has been accomplished in symbionts [58–61], it has often been done with technologies that did not allow for site specific integration. Paratransgenic approaches suitable for use in the field will need to stably integrate genes into the bacterial genome in a manner that does not compromise bacterial fitness. Exploiting the flexibility and specificity of the CRISPR/Cas9 system to integrate genes in intergenic regions of the bacterial chromosome will undoubtedly be beneficial for these applied approaches.

In summary, we have demonstrated that the CRISPR/Cas9 gene editing system can be applied to symbiotic bacteria that associate with eukaryotic hosts to interrogate the role of bacterial genes in host-microbe associations. We created knockout and knockin mutants by deleting and disrupting the *ompA* gene of *C. neteri*. The knockout mutant displayed a reduced ability to form biofilms and colonize the gut of *Ae. aegypti* mosquitoes in a mono-association demonstrating bacterial genetic factors are important determinants that influence colonization of mosquito guts. *Aedes* mosquitoes are becoming powerful systems to investigate the genetics of host-microbe interactions given the scientific community has simple and efficient approaches to alter both the microbes (this study) and mosquito host genome [76, 77] at their disposal, as well as methods to create mono-associated mosquito lines [7, 64]. Finally, rapid, efficient, and site specific gene editing approaches for gut bacteria that associate with mosquitoes will facilitate the development of novel paratransgenic approaches to control arthropod-borne disease [57].

Material and methods

Bacterial and mosquito strains

E. coli BL21(DE3) (NEB) and *Cedecea neteri* strain Alb1, previous isolated from a lab-reared colony of *Ae. albopictus* (Galveston) mosquitoes [7], were used in this study. To further classify the gut-associated bacteria we completed multilocus sequence typing [78]. DNA from the single colony was used as a template in a PCR to amplify genes for MLST analysis (S3 Table). Amplicons were resolved on a 1% agarose gel, extracted and purified, and Sanger sequenced. The *atpD*, *infB*, *gyrB* and *rpoB* genes were aligned separately, using the species diversity as in [79] with several *Cedecea* sp. sequences and then concatenated using seaview [79]. The phylogenetic tree was constructed using iqtree [80] under the general time-reversible (GTR) model with 1000 fast bootstrap replicates, which are shown as percentage branch support values (S4 Fig). The sequences of our isolate are available under accessions (MN329096 (*atpD*), MN329097 (*gyrB*), MN329098 (*infB*), MN329099 (*rpoB*)). For gene editing and mosquito infections, cultures were grown in liquid LB media at 37°C with the appropriate antibiotic unless

stated otherwise. Mosquitoes were reared in the UTMB insectary under conventional conditions or in a mono-association (described below).

CRISPR gene editing

Designing protospacer sequence and cloning: The *E. coli* BL21 *ompA* gene sequence was retrieved from NCBI (accession number LR536431). The *C. neteri* Alb1 *ompA* gene was PCR amplified and Sanger sequenced using primers (OmpA-F and OmpA-R, [S3 Table](#)), which were designed based on the *Enterobacter cloacae ompA* (accession number CP017990). Editing the *ompA* gene of *E. coli* and *C. neteri* was complete as described in Reisch and Prather [35]. Protospacer sequences for the *ompA* gene were designed using CHOPCHOP [81, 82]. To clone the protospacer sequences into pKDsgRNA-ack ([S2 Table](#); Addgene plasmid #62654) we amplified the entire plasmid with primers that contained the protospacer sequence and this amplicon was self-ligated. This PCR was done using 0.5 μ M of each primer ([S1 Table](#)), 1x reaction buffer, 200 μ M dNTPs, 0.5U of Phire Host Start Taq polymerase (Thermo Scientific) and 200 ng of plasmid DNA as template. The cycling condition consisted of an initial denaturation step 98°C for 2 minutes, followed by 35 cycles of 98°C for 2 seconds, 58°C for 15 seconds, and 72°C for 2 minutes and 30 seconds, and then a final extension at 72°C for 10 minutes before holding at 16°C. The PCR products had a 15–17 bp overlapping sequence which was used to ligate the plasmid. The PCR product was digested with *DpnI* to remove any template plasmid. PCR products were then ligated by transformation into *E. coli* harbouring the Red/ET plasmid following the REPLACR mutagenesis protocol [83], thereby creating plasmids pKDsgRNA-Ec-*ompA*-1, pKDsgRNA-Ec-*ompA*-2, pKDsgRNA-Ent-*ompA*-1, and pKDsgRNA-Ent-*ompA*-2 ([S2 Table](#)). Colonies were screened for the protospacer insertion by PCR and confirmed by Sanger sequencing.

Knockout of *ompA*

The two protospacers were evaluated by transforming plasmids into either *E. coli* or *C. neteri* containing the pCas9-CR4 plasmid ([S2 Table](#); Addgene plasmid 62655), which expressed Cas9 nuclease. Transformants were selected at 30°C on LB agar plate containing spectinomycin (50 μ g/mL), chloramphenicol (34 μ g/mL), and either with or without anhydrotetracycline (aTC; 100ng/mL). The escape rate was quantified by comparing colonies in the plates with or without aTC. The protospacer with a lack of or few escape mutants was used for further experiments. Colonies from the –aTC plate were grown overnight in LB broth with the appropriate antibiotic at 30°C. A 1:100 diluted overnight culture was (grown until 0.4 OD₆₀₀) supplemented with 1.2% arabinose to induce the expression of λ -Red recombinase for 20 min. Cells were then transformed with 1–1.5 μ g of double stranded donor DNA for homologous recombination. Donor DNA was created by PCR amplifying the flanking left arm (LA) and right arm (RA) from *E. coli* and *C. neteri* genomic DNA. Each arm had flanking regions of 250 bp homologous to the target DNA. The resulting fragment was assembled using Gibson assembly (NEB). The assembled product was amplified to generate full length dsDNA for transformation. Colonies were screened for mutations by colony PCR with primers flanking the integration site and positive clones were Sanger sequenced ([S3 Table](#)). Positive colonies were grown in LB broth and genomic DNA was isolated. For further validation, the flanking regions of deletion or insertions were amplified, and the PCR product Sanger sequenced.

Insertion of mCherry and gentamicin gene into *C. neteri* genome

The plasmid pKDsgRNA-Ent-*ompA* was transformed into *C. neteri* and the gene editing procedure was repeated as described above. To generate the donor sequence for homologous

recombination the mCherry or gentamicin sequence (driven by the AmTr promoter) and each homology arm were amplified and ligated. The assembled product was amplified to generate a full length dsDNA fragment for transformation.

Stability of insertion

The stability of the knockout $\Delta ompA$ mutant and the knockin $ompA::gentamicin$ and $ompA::mCherry$ strains was assessed in LB media. The $ompA::mCherry$ and knockout $\Delta ompA$ mutant cultures were grown for 10 passages in LB broth. At each passage 40 μ l of culture was transferred into 4ml fresh LB media. The $ompA::gentamicin$ strain was grown with or without gentamicin (50 μ g/mL). Genomic DNA was isolated from the 0, 2, 4, 6, 8 and 10th subculture and PCR that amplified across the integration site was performed.

Complementation of *ompA* mutant

Functional rescue of the *ompA* mutation was achieved by complementing the mutant with the WT gene. The WT *ompA* gene was amplified from *C. neteri* genomic DNA and cloned into the pRAM-mCherry vector [7] in front of the *ompA* promoter, thereby creating pRAM-mCherry-*Ent-OmpA* plasmid. The Sanger sequence-verified plasmid was transformed into the $\Delta ompA$ mutant, thereby generating the $\Delta ompA/ompA$ complement strain. Colonies that acquired the plasmid were selected on LB plates containing kanamycin (50 μ g/mL).

In vitro characterization of *C. neteri* strains

To assess the impact of the gene deletion on bacterial growth the WT, $\Delta ompA$ mutant and $\Delta ompA/ompA$ complement were grown in LB broth and the density of bacteria (OD₆₀₀) was quantified by spectrophotometer. A 1:100 dilution of an overnight culture was inoculated into a 5 ml LB broth in a 50 ml tube and incubated at 37°C for 24 hrs. At 2, 4, 6, 8, 10, 12 and 24 hours growth was recorded at OD₆₀₀. The biofilm assay was performed as described previously [84, 85]. Briefly, biofilm formation by *C. neteri* strains was quantified on polystyrene microtiter plates after 72 h of incubation at 37°C by CV staining. Three independent experiments were performed, and the data were represented as CV OD₅₇₀ after normalizing by CFUs.

Mosquito infections

Mono-association in *Ae. aegypti* mosquitoes were done using gnotobiotic infection procedure [7, 64], with slight modifications. Briefly, mosquito eggs were sterilized for 5 min in 70% ethanol, 3 min in 3% bleach+0.01% Coverage Plus NPD (Steris Corp.), 5 min in 70% ethanol then rinsed three times in sterile water. Eggs were vacuumed hatched for 30–45 min and left overnight at room temperature to hatch any remaining eggs. Exactly twenty L1 larvae were transferred to T175 flask containing 60 ml of sterile water and fed on alternative days with 60 μ l of fish food (1 μ g/ μ l). Larvae were inoculated with 1x10⁷/ml of either the WT *C. neteri*, the $\Delta ompA$ mutant or the $\Delta ompA/ompA$ complement. The WT and $\Delta ompA$ strains were transformed with the pRAM-mCherry plasmid [7] that conferred resistance to kanamycin (but did not possess a functional *ompA* gene). We also performed gnotobiotic infections with WT *C. neteri*, knockin mutants all expressing mCherry from a plasmid. In order to confirm that eggs were successfully sterilized, a T175 flask containing twenty L1 larvae were reared in identical fashion to mono-associations, albeit without bacterial supplementation. These larvae did not develop beyond the L2 stage, indicating our rearing process was free from contamination. To quantify bacteria, L4 larvae were collected, washed three times with 1X PBS, and then homogenized in 500 μ l of 1X PBS and 50 μ l of homogenate was plated on LB agar containing 50 μ g/mL

kanamycin. Similarly, adult mosquitoes were collected 3–4 days post emergence and bacterial infection was quantified in the same manner as larvae. In order to assess the growth of the mosquitoes, time to pupation and growth rate were observed. Time to pupation was determined by quantifying the number of pupae each day post hatching, while survival to adulthood was calculated by quantifying the number of L1 larvae that reached adulthood. The experiment was repeated three times.

Reinfection of knockin mutants to mosquitoes

Knockin mutants were administered to 3–4 days adult *Ae. aegypti* in a sugar meal. These mosquitoes were reared under normal laboratory condition. Mosquitoes were fed with 1×10^7 of WT or the $\Delta ompA$::gentamicin strain for three days in 10% sucrose solution. After three days, mosquitoes were either administered sugar supplemented with gentamicin (50 $\mu\text{g}/\text{mL}$) or sugar without antibiotic. CFUs were determined at days 0, 2, 4, and 6 dpi by plating homogenized mosquitoes ($N = 10$) on LB agar. Similarly, the $\Delta ompA$::mCherry and WT *C. neteri* were fed to mosquitoes and midguts were dissected to assess colonization of bacteria in the tissue. For visualization of bacteria, midguts were fixed in 1% paraformaldehyde (PFA) in 1X PBS for 30 minutes and permeabilized with 0.01% Triton X-100 in 1X PBS for 20 min. The tissues were stained with 1:250 diluted Phalloidin (Sigma) for 20 minutes and samples were washed twice with 1X PBS for 10 minutes. Finally, midguts were then stained with 1:500 diluted DAPI (Invitrogen) for 10 min. Samples were transferred to slides and mounted with ProLong™ Gold Antifade (Invitrogen). The slides were observed using a Revolve FL microscope (ECHOLAB).

Supporting information

S1 Fig. Midgut infection of *C. neteri* and *E. coli* in mono-associations of *Aedes* mosquitoes.

Dissected gut tissue showing the conglomeration of bacterial cells when infected in mono-association in *Aedes* mosquitoes with *C. neteri* (A-C). However, *E. coli* (D-F) and $\Delta ompA$ (G-I) could not be seen in the midgut. Images were captured from the dissected midguts of different mosquitoes. Bacteria possessed the pRAM-mCherry plasmid which expressed the mCherry fluorescent protein.

(TIFF)

S2 Fig. Phylogenetic analysis of WT and mutants. Multilocus sequence analysis according to [78] indicates isolates to be members of the *C. neteri* species; the MLST genes were amplified in the wild type isolate and two mutants as shown in the tree (red).

(TIFF)

S3 Fig. Average CFUs of mosquito infection. Average CFU recovered in adult mosquitoes infected with *C. neteri* strains (WT, $\Delta ompA$ mutant and $\Delta ompA/ompA$ complement) reared in a mono-association using a gnotobiotic rearing approach. The uninfected mosquitoes were removed from the analysis. Box and whiskers show the 25th and 75th percentiles and the minimum and maximum values, respectively.

(TIFF)

S4 Fig. Mono-association infection and biofilm assessment. L1 axenic larvae were infected with WT *C. neteri* (A), $\Delta ompA$::mCherry (B) and $\Delta ompA$::gentamicin (C) and adults gut were analysed for presence of bacterial conglomerations (biofilm formation). For each treatment, 15–18 midguts were screened. Scale bar 130 μm .

(PDF)

S1 Table. gRNA sequence used in this study.

(XLSX)

S2 Table. Plasmids and bacterial strains used in this study.

(XLSX)

S3 Table. Primers used in this study.

(XLSX)

S4 Table. Infection and biofilm prevalence of mosquito midguts when reared in a mono-association.

(XLSX)

S1 Appendix. Multiple sequence alignment of WT and mutant *ompA* sequences of *C. neteri* and *E. coli*.

(DOCX)

Acknowledgments

We would like to thank the UTMB insectary core for providing mosquitoes. We also wish to thank Dr Ulrike Munderloh (University of Minnesota) for the kind gift of the pRAM plasmid.

Author Contributions

Conceptualization: Shivanand Hegde, Grant L. Hughes.

Formal analysis: Shivanand Hegde, Eva Heinz, Grant L. Hughes.

Funding acquisition: Grant L. Hughes.

Investigation: Shivanand Hegde, Pornjarim Nilyanimit, Elena Kozlova, Hema P. Narra, Grant L. Hughes.

Methodology: Shivanand Hegde, Pornjarim Nilyanimit, Elena Kozlova, Enyia R. Anderson, Hema P. Narra.

Project administration: Grant L. Hughes.

Supervision: Sanjeev K. Sahni, Grant L. Hughes.

Validation: Shivanand Hegde.

Writing – original draft: Shivanand Hegde.

Writing – review & editing: Eva Heinz, Grant L. Hughes.

References

1. Osei-Poku J, Mbogo CM, Palmer WJ, Jiggins FM. Deep sequencing reveals extensive variation in the gut microbiota of wild mosquitoes from Kenya. *Molecular Ecology*. 2012; 21(20):5138–50. <https://doi.org/10.1111/j.1365-294X.2012.05759.x> PMID: 22988916
2. Boissière A, Tchioffo MT, Bachar D, Abate L, Marie A, Nsango SE, et al. Midgut microbiota of the malaria mosquito vector *Anopheles gambiae* and interactions with *Plasmodium falciparum* infection. *PLoS Pathogens*. 2012; 8(5):e1002742. <https://doi.org/10.1371/journal.ppat.1002742> PMID: 22693451
3. David MR, Santos LMBD, Vicente ACP, Maciel-de-Freitas R. Effects of environment, dietary regime and ageing on the dengue vector microbiota: evidence of a core microbiota throughout *Aedes aegypti* lifespan. *Memórias do Instituto Oswaldo Cruz*. 2016; 111(9):577–87. <https://doi.org/10.1590/0074-02760160238> PMID: 27580348

4. Muturi EJ, Kim C-H, Bara J, Bach EM, Siddappaji MH. *Culex pipiens* and *Culex restuans* mosquitoes harbor distinct microbiota dominated by few bacterial taxa. *Parasites & Vectors*. 2016; 9(1):18. <https://doi.org/10.1186/s13071-016-1299-6> PMID: 26762514
5. Hughes GL, Dodson BL, Johnson RM, Murdock CC, Tsujimoto H, Suzuki Y, et al. Native microbiome impedes vertical transmission of *Wolbachia* in *Anopheles* mosquitoes. *Proceedings of the National Academy of Sciences of the United States of America*. 2014; 111(34):12498–503. <https://doi.org/10.1073/pnas.1408888111> PMID: 25114252
6. Coon KL, Brown MR, Strand MR. Mosquitoes host communities of bacteria that are essential for development but vary greatly between local habitats. *Molecular Ecology*. 2016; 25(22):5806–26. <https://doi.org/10.1111/mec.13877> PMID: 27718295
7. Hegde S, Khanipov K, Albayrak L, Golovko G, Pimenova M, Saldaña MA, et al. Microbiome interaction networks and community structure from laboratory-reared and field-collected *Aedes aegypti*, *Aedes albopictus*, and *Culex quinquefasciatus* mosquito vectors. *Frontiers in Microbiology*. 2018; 9:715.
8. Wang Y, Gilbreath TM, Kukutla P, Yan G, Xu J. Dynamic gut microbiome across life history of the malaria mosquito *Anopheles gambiae* in Kenya. *PLoS ONE*. 2011; 6(9):e24767. <https://doi.org/10.1371/journal.pone.0024767> PMID: 21957459
9. Weiss BL, Aksoy S. Microbiome influences on insect host vector competence. *Trends in Parasitology*. 2011; 27(11):514–22. <https://doi.org/10.1016/j.pt.2011.05.001> PMID: 21697014
10. Saldaña MA, Hegde S, Hughes GL. Microbial control of arthropod-borne disease. *Memórias do Instituto Oswaldo Cruz*. 2017; 112(2):81–93. <https://doi.org/10.1590/0074-02760160373> PMID: 28177042
11. Hegde S, Rasgon JL, Hughes GL. The microbiome modulates arbovirus transmission in mosquitoes. *Current Opinion in Virology*. 2015; 15:97–102. <https://doi.org/10.1016/j.coviro.2015.08.011> PMID: 26363996
12. Audsley MD, Ye YH, McGraw EA. The microbiome composition of *Aedes aegypti* is not critical for *Wolbachia*-mediated inhibition of dengue virus. *PLoS Neglected Tropical Diseases*. 2017; 11(3):e0005426. <https://doi.org/10.1371/journal.pntd.0005426> PMID: 28267749
13. Muturi EJ, Bara JJ, Rooney AP, Hansen AK. Midgut fungal and bacterial microbiota of *Aedes triseriatus* and *Aedes japonicus* shift in response to La-Crosse virus infection. *Molecular Ecology*. 2016; 25(16):4075–90. <https://doi.org/10.1111/mec.13741> PMID: 27357374
14. Kumar S, Molina-Cruz A, Gupta L, Rodrigues J, Barillas-Mury C. A peroxidase/dual oxidase system modulates midgut epithelial immunity in *Anopheles gambiae*. *Science*. 2010; 327(5973):1644–8. <https://doi.org/10.1126/science.1184008> PMID: 20223948
15. Pang X, Xiao X, Liu Y, Zhang R, Liu J, Liu Q, et al. Mosquito C-type lectins maintain gut microbiome homeostasis. *Nature Microbiology*. 2016; 1(5):16023. <https://doi.org/10.1038/nmicrobiol.2016.23> PMID: 27170846
16. Short SM, Mongodin EF, MacLeod HJ, Talyuli OAC, Dimopoulos G. Amino acid metabolic signaling influences *Aedes aegypti* midgut microbiome variability. *PLoS Neglected Tropical Diseases*. 2017; 11(7):e0005677–29. <https://doi.org/10.1371/journal.pntd.0005677> PMID: 28753661
17. Stathopoulos S, Neafsey DE, Lawniczak MKN, Muskavitch MAT, Christophides GK. Genetic dissection of *Anopheles gambiae* gut epithelial responses to *Serratia marcescens*. *PLOS Pathogens*. 2014; 10(3):e1003897. <https://doi.org/10.1371/journal.ppat.1003897> PMID: 24603764
18. Xiao X, Yang L, Pang X, Zhang R, Zhu Y, Wang P, et al. A Mesh-Duox pathway regulates homeostasis in the insect gut. *Nature Microbiology*. 2017; 2(5):17020. <https://doi.org/10.1038/nmicrobiol.2017.20> PMID: 28248301
19. Powell JE, Leonard SP, Kwong WK, Engel P, Moran NA. Genome-wide screen identifies host colonization determinants in a bacterial gut symbiont. *Proceedings of the National Academy of Sciences of the United States of America*. 2016; 201610856. <https://doi.org/10.1073/pnas.1610856113> PMID: 27849596
20. Kim JK, Kwon JY, Kim S-K, Han SH, Won YJ, Lee J-H, et al. Purine biosynthesis, biofilm formation, and persistence of an insect-microbe gut symbiosis. *Appl Environ Microbiol*. 2014; 80(14):4374–82. <https://doi.org/10.1128/AEM.00739-14> PMID: 24814787
21. Mulcahy H, Sibley CD, Surette MG, Lewenza S. *Drosophila melanogaster* as an animal model for the study of *Pseudomonas aeruginosa* biofilm infections in vivo. *PLOS Pathogens*. 2011; 7(10):e1002299–14. <https://doi.org/10.1371/journal.ppat.1002299> PMID: 21998591
22. Maltz M, Graf J. The type II secretion system is essential for erythrocyte lysis and gut colonization by the leech digestive tract symbiont *Aeromonas veronii*. *Applied and Environmental Microbiology*. 2011; 77(2):597–603. <https://doi.org/10.1128/AEM.01621-10> PMID: 21097598
23. Weiss BL, Wu Y, Schwank JJ, Tolwinski NS, Aksoy S. An insect symbiosis is influenced by bacterium-specific polymorphisms in outer-membrane protein A. *Proceedings of the National Academy of Sciences of the United States of America*. 2016; 113(12):3253–8. <https://doi.org/10.1073/pnas.1518888113> PMID: 27000000

- Sciences of the United States of America. 2008; 105(39):15088–93. <https://doi.org/10.1073/pnas.0805666105> PMID: 18815366
24. Maltz MA, Weiss BL, O apos Neill M, Wu Y, Aksoy S. OmpA-mediated biofilm formation is essential for the commensal bacterium *Sodalis glossinidius* to colonize the tsetse fly gut. *Applied and Environmental Microbiology*. 2012; 78(21):7760–8. <https://doi.org/10.1128/AEM.01858-12> PMID: 22941073
 25. Riehle MA, Jacobs-Lorena M. Using bacteria to express and display anti-parasite molecules in mosquitoes: current and future strategies. *Insect Biochemistry and Molecular Biology*. 2005; 35(7):699–707. <https://doi.org/10.1016/j.ibmb.2005.02.008> PMID: 15894187
 26. Dennison NJ, Saraiva RG, Cirimotich CM, Mlambo G, Mongodin EF, Dimopoulos G. Functional genomic analyses of *Enterobacter*, *Anopheles* and *Plasmodium* reciprocal interactions that impact vector competence. *Malaria Journal*. 2016; 15(1):425. Epub 3. <https://doi.org/10.1186/s12936-016-1468-2> PMID: 27549662
 27. Pei D, Jiang J, Yu W, Kukutla P, Uentillie A, Xu J. The *waaL* gene mutation compromised the inhabitation of *Enterobacter sp. Ag1* in the mosquito gut environment. *Parasites & Vectors*. 2015;1–10. <https://doi.org/10.1186/s13071-015-1049-1> PMID: 26306887
 28. Enomoto S, Chari A, Clayton AL, Dale C. Quorum sensing attenuates virulence in *Sodalis praecaptivus*. *Cell Host & Microbe*. 2017; 21(5):629–36.e5. <https://doi.org/10.1016/j.chom.2017.04.003> PMID: 28494244
 29. Dale C, Young SA, Haydon DT, Welburn SC. The insect endosymbiont *Sodalis glossinidius* utilizes a type III secretion system for cell invasion. *Proceedings of the National Academy of Sciences*. 2001; 98(4):1883–8. <https://doi.org/10.1073/pnas.021450998> PMID: 11172045
 30. Selle K, Barrangou R. Harnessing CRISPR–Cas systems for bacterial genome editing. *Trends in Microbiology*. 2015; 23(4):225–32. <https://doi.org/10.1016/j.tim.2015.01.008> PMID: 25698413
 31. Sander JD, Joung JK. CRISPR–Cas systems for editing, regulating and targeting genomes. *Nature Biotechnology*. 2014; 32(4):347–55. <https://doi.org/10.1038/nbt.2842> PMID: 24584096
 32. Barrangou R, van Pijkeren JP. Exploiting CRISPR–Cas immune systems for genome editing in bacteria. *Current Opinion in Biotechnology*. 2016; 37:61–8. <https://doi.org/10.1016/j.copbio.2015.10.003> PMID: 26629846
 33. Jiang Y, Chen B, Duan C, Sun B, Yang J, Yang S. Multigene editing in the *Escherichia coli* genome via the CRISPR–Cas9 system. *Applied and Environmental Microbiology*. 2015; 81(7):2506–14. <https://doi.org/10.1128/AEM.04023-14> PMID: 25636838
 34. Jiang W, Bikard D, Cox D, Zhang F, Marraffini LA. RNA-guided editing of bacterial genomes using CRISPR–Cas systems. *Nature Biotechnology*. 2013; 31(3):233–9. <https://doi.org/10.1038/nbt.2508> PMID: 23360965
 35. Reisch CR, Prather KLJ. The no-SCAR (Scarless Cas9 Assisted Recombineering) system for genome editing in *Escherichia coli*. *Scientific Reports*. 2015; 5:15096. <https://doi.org/10.1038/srep15096> PMID: 26463009
 36. Li Y, Lin Z, Huang C, Zhang Y, Wang Z, Tang Y-j, et al. Metabolic engineering of *Escherichia coli* using CRISPR–Cas9 mediated genome editing. *Metabolic Engineering*. 2015; 31:13–21. <https://doi.org/10.1016/j.ymben.2015.06.006> PMID: 26141150
 37. Ronda C, Pedersen LE, Sommer MOA, Nielsen AT. CRMAGE: CRISPR Optimized MAGE Recombineering. *Scientific Reports*. 2016; 6(1):1200. <https://doi.org/10.1038/srep19452> PMID: 26797514
 38. Tong Y, Robertsen HL, Blin K, Weber T, Lee SY. CRISPR–Cas9 Toolkit for *Actinomyces* Genome Editing. *Methods in molecular biology* 2018; 1671(1):163–84. https://doi.org/10.1007/978-1-4939-7295-1_11 PMID: 29170959
 39. Oh J-H, van Pijkeren JP. CRISPR–Cas9-assisted recombineering in *Lactobacillus reuteri*. *Nucleic Acids Research*. 2014; 42(17):e131–e. <https://doi.org/10.1093/nar/gku623> PMID: 25074379
 40. Mougialos I, Bosma EF, Weenink K, Vossen E, Goijvaerts K, van der Oost J, et al. Efficient genome editing of a facultative thermophile using mesophilic spCas9. *ACS Synthetic Biology*. 2017; 6(5):849–61. <https://doi.org/10.1021/acssynbio.6b00339> PMID: 28146359
 41. Li K, Cai D, Wang Z, He Z, Chen S. Development of an efficient genome editing tool in *Bacillus licheniformis* using CRISPR–Cas9 nickase. *Applied and Environmental Microbiology*. 2018:AEM.02608-17. <https://doi.org/10.1128/AEM.02608-17> PMID: 29330178
 42. Jiang Y, Qian F, Yang J, Liu Y, Dong F, Xu C, et al. CRISPR–Cpf1 assisted genome editing of *Corynebacterium glutamicum*. *Nat Commun*. 2017; 8:15179. Epub 2017/05/05. <https://doi.org/10.1038/ncomms15179> PMID: 28469274
 43. Cobb RE, Wang Y, Zhao H. High-efficiency multiplex genome editing of *Streptomyces* species using an engineered CRISPR/Cas system. *ACS Synth Biol*. 2015; 4(6):723–8. Epub 2014/12/03. <https://doi.org/10.1021/sb500351f> PMID: 25458909

44. Waller MC, Bober JR, Nair NU, Beisel CL. Toward a genetic tool development pipeline for host-associated bacteria. *Current Opinion in Microbiology*. 2017; 38:156–64. <https://doi.org/10.1016/j.mib.2017.05.006> PMID: 28624690
45. Wilke ABB, Marrelli MT. Paratransgenesis: a promising new strategy for mosquito vector control. *Parasites & Vectors*. 2015; 8(1):391–19. <https://doi.org/10.1186/s13071-015-0959-2> PMID: 26104575
46. Arora AK, Douglas AE. Hype or opportunity? Using microbial symbionts in novel strategies for insect pest control. *Journal of insect physiology*. 2017; 103:10–7. <https://doi.org/10.1016/j.jinsphys.2017.09.011> PMID: 28974456
47. Ricci I, Damiani C, Capone A, DeFreece C, Rossi P, Favia G. Mosquito/microbiota interactions: from complex relationships to biotechnological perspectives. *Current Opinion in Microbiology*. 2012; 15(3):278–84. <https://doi.org/10.1016/j.mib.2012.03.004> PMID: 22465193
48. Beard CB, Mason PW, Aksoy S, Tesh RB, Richards FF. Transformation of an insect symbiont and expression of a foreign gene in the Chagas' disease vector *Rhodnius prolixus*. *American Journal of Tropical Medicine and Hygiene*. 1992; 46(2):195–200. <https://doi.org/10.4269/ajtmh.1992.46.195> PMID: 1539755
49. Bisi DC, Lampe DJ. Secretion of anti-Plasmodium effector proteins from a natural *Pantoea agglomerans* isolate by using PeIB and HlyA secretion signals. *Applied and Environmental Microbiology*. 2011; 77(13):4669–75. <https://doi.org/10.1128/AEM.00514-11> PMID: 21602368
50. Hughes GL, Allsopp PG, Webb RI, Yamada R, Iturbe-Ormaetxe I, Brumbley SM, et al. Identification of yeast associated with the planthopper, *Perkinsiella saccharicida*: potential applications for Fiji leaf gall control. *Current Microbiology*. 2011; 63(4):392–401. <https://doi.org/10.1007/s00284-011-9990-5> PMID: 21850475
51. Medina F, Li H, Vinson SB, Coates CJ. Genetic transformation of midgut bacteria from the red imported fire ant (*Solenopsis invicta*). *Current Microbiology*. 2009; 58(5):478–82. <https://doi.org/10.1007/s00284-008-9350-2> PMID: 19159973
52. Bextine B, Lauzon C, Potter S, Lampe D, Miller TA. Delivery of a genetically marked *Alcaligenes sp.* to the glassy-winged sharpshooter for use in a paratransgenic control strategy. *Current Microbiology*. 2004; 48(5):327–31. <https://doi.org/10.1007/s00284-003-4178-2> PMID: 15060727
53. Hurwitz I, Hillesland H, Fieck A, Das P, Durvasula R. The paratransgenic sand fly: A platform for control of *Leishmania* transmission. *Parasites & Vectors*. 2011; 4(1):82. <https://doi.org/10.1186/1756-3305-4-82> PMID: 21595907
54. Durvasula RV, Gumbs A, Panackal A, Kruglov O, Taneja J, Kang AS, et al. Expression of a functional antibody fragment in the gut of *Rhodnius prolixus* via transgenic bacterial symbiont *Rhodococcus rhodnii*. *Medical and Veterinary Entomology*. 1999; 13(2):115–9. <https://doi.org/10.1046/j.1365-2915.1999.00175.x> PMID: 10484156
55. Wang S, Ghosh AK, Bongio N, Stebbings KA, Lampe DJ, Jacobs-Lorena M. Fighting malaria with engineered symbiotic bacteria from vector mosquitoes. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109(31):12734–9. <https://doi.org/10.1073/pnas.1204158109> PMID: 22802646
56. Wu SC-Y, Maragathavally KJ, Coates CJ, Kaminski JM. Steps toward targeted insertional mutagenesis with class II transposable elements. *Methods in molecular biology* 2008; 435:139–51. https://doi.org/10.1007/978-1-59745-232-8_10 PMID: 18370073
57. Tikhe CV, Martin TM, Howells A, Delatte J, Husseneder C. Assessment of genetically engineered *Tribolium odontotermis* as a 'Trojan Horse' for paratransgenesis in termites. *BMC Microbiology*. 2016; 16(1):355. <https://doi.org/10.1186/s12866-016-0822-4> PMID: 27595984
58. Pittman GW, Brumbley SM, Allsopp PG, O apos Neill SL. Assessment of gut bacteria for a paratransgenic approach to control *Dermolepida albobhirtum* larvae. *Applied and Environmental Microbiology*. 2008; 74(13):4036–43. <https://doi.org/10.1128/AEM.02609-07> PMID: 18456847
59. Pontes MH, Dale C. Lambda red-mediated genetic modification of the insect endosymbiont *Sodalis glossinidius*. *Applied and Environmental Microbiology*. 2011; 77(5):1918–20. <https://doi.org/10.1128/AEM.02166-10> PMID: 21216910
60. Wang S, Dos-Santos ALA, Huang W, Liu KC, Oshaghi MA, Wei G, et al. Driving mosquito refractoriness to *Plasmodium falciparum* with engineered symbiotic bacteria. *Science (New York, NY)*. 2017; 357(6358):1399–402. <https://doi.org/10.1126/science.aan5478> PMID: 28963255
61. Dotson EM, Plikaytis B, Shinnick TM, Durvasula RV, Beard CB. Transformation of *Rhodococcus rhodnii*, a symbiont of the Chagas disease vector *Rhodnius prolixus*, with integrative elements of the L1 mycobacteriophage. *Infection, genetics and evolution*. 2003; 3(2):103–9. [https://doi.org/10.1016/s1567-1348\(03\)00002-9](https://doi.org/10.1016/s1567-1348(03)00002-9) PMID: 12809804

62. Bextine B, Lampe D, Lauzon C, Jackson B, Miller TA. Establishment of a genetically marked insect-derived symbiont in multiple host plants. *Curr Microbiol.* 2005; 50(1):1–7. <https://doi.org/10.1007/s00284-004-4390-8> PMID: 15723145
63. Wu P, Sun P, Nie K, Zhu Y, Shi M, Xiao C, et al. A gut commensal bacterium promotes mosquito permissiveness to arboviruses. *Cell Host & Microbe.* 2019; 25(1):101–12.e5. <https://doi.org/10.1016/j.chom.2018.11.004> PMID: 30595552
64. Coon KL, Vogel KJ, Brown MR, Strand MR. Mosquitoes rely on their gut microbiota for development. *Molecular Ecology.* 2014; 23(11):2727–39. <https://doi.org/10.1111/mec.12771> PMID: 24766707
65. Zalewska-Piatek B, Wilkanowicz S, Bruzdziak P, Piatek R, Kur J. Biochemical characteristic of biofilm of uropathogenic *Escherichia coli* Dr(+) strains. *Microbiol Res.* 2013; 168(6):367–78. Epub 2013/02/05. <https://doi.org/10.1016/j.micres.2013.01.001> PMID: 23375236
66. Lindh JM, Borg-Karlson AK, Faye I. Transstadial and horizontal transfer of bacteria within a colony of *Anopheles gambiae* (Diptera: Culicidae) and oviposition response to bacteria-containing water. *Acta Tropica.* 2008; 107(3):242–50. <https://doi.org/10.1016/j.actatropica.2008.06.008> PMID: 18671931
67. Bäckman S, Näslund J, Forsman M, Thelaus J. Transmission of tularemia from a water source by transstadial maintenance in a mosquito vector. *Scientific Reports.* 2015; 5:7793. <https://doi.org/10.1038/srep07793> PMID: 25609657
68. Chavshin AR, Oshaghi MA, Vatandoost H, Yakhchali B, Zarenejad F, Terenius O. Malpighian tubules are important determinants of *Pseudomonas* transstadial transmission and longtime persistence in *Anopheles stephensi*. *Parasites & Vectors.* 2015; 8(1):36. <https://doi.org/10.1186/s13071-015-0635-6> PMID: 25604581
69. Chen S, Bagdasarian M, Walker ED. *Elizabethkingia anophelis*: Molecular Manipulation and Interactions with Mosquito Hosts. *Applied and Environmental Microbiology.* 2015; 81(6):2233. <https://doi.org/10.1128/AEM.03733-14> PMID: 25595771
70. Coon KL, Brown MR, Strand MR. Gut bacteria differentially affect egg production in the anautogenous mosquito *Aedes aegypti* and facultatively autogenous mosquito *Aedes atropalpus* (Diptera: Culicidae). *Parasites & Vectors.* 2016; 9(1):375. <https://doi.org/10.1186/s13071-016-1660-9> PMID: 27363842
71. Ribet D, Cossart P. How bacterial pathogens colonize their hosts and invade deeper tissues. *Microbes and infection.* 2015; 17(3):173–83. <https://doi.org/10.1016/j.micinf.2015.01.004> PMID: 25637951
72. Ramirez JL, Short SM, Bahia AC, Saraiva RG, Dong Y, Kang S, et al. *Chromobacterium Csp_P* reduces malaria and dengue infection in vector mosquitoes and has entomopathogenic and in vitro anti-pathogen activities. *PLoS Pathogens.* 2014; 10(10):e1004398. <https://doi.org/10.1371/journal.ppat.1004398> PMID: 25340821
73. Valzania L, Martinson VG, Harrison RE, Boyd BM, Coon KL, Brown MR, et al. Both living bacteria and eukaryotes in the mosquito gut promote growth of larvae. *PLoS Neglected Tropical Diseases.* 2018; 12(7):e0006638. <https://doi.org/10.1371/journal.pntd.0006638> PMID: 29979680
74. Correa MA, Matusovsky B, Brackney DE, Steven B. Generation of axenic *Aedes aegypti* demonstrate live bacteria are not required for mosquito development. *Nature Communications.* 2018; 9(1):R37. <https://doi.org/10.1038/s41467-018-07014-2> PMID: 30367055
75. Favia G, Ricci I, Damiani C, Raddadi N, Crotti E, Marzorati M, et al. Bacteria of the genus *Asaia* stably associate with *Anopheles stephensi*, an Asian malarial mosquito vector. *Proceedings of the National Academy of Sciences.* 2007; 104(21):9047–51. <https://doi.org/10.1073/pnas.0610451104> PMID: 17502606
76. Chaverra-Rodriguez D, Macias VM, Hughes GL, Pujhari S, Suzuki Y, Peterson DR, et al. Targeted delivery of CRISPR-Cas9 ribonucleoprotein into arthropod ovaries for heritable germline gene editing. *Nature Communications.* 2018; 9(1):245.
77. Li M, Bui M, Yang T, Bowman CS, White BJ, Akbari OS. Germline Cas9 expression yields highly efficient genome engineering in a major worldwide disease vector, *Aedes aegypti*. *Proceedings of the National Academy of Sciences of the United States of America.* 2017; 114(49):E10540–E9. <https://doi.org/10.1073/pnas.1711538114> PMID: 29138316
78. Brady C, Cleenwerck I, Venter S, Coutinho T, De Vos P. Taxonomic evaluation of the genus *Enterobacter* based on multilocus sequence analysis (MLSA): Proposal to reclassify *E. nimipressuralis* and *E. amnigenus* into *Lelliottia* gen. nov. as *Lelliottia nimipressuralis* comb. nov. and *Lelliottia amnigena* comb. nov., respectively, *E. gergoviae* and *E. pyrinus* into *Pluralibacter* gen. nov. as *Pluralibacter gergoviae* comb. nov. and *Pluralibacter pyrinus* comb. nov., respectively, *E. cowanii*, *E. radincitans*, *E. oryzae* and *E. arachidis* into *Kosakonia* gen. nov. as *Kosakonia cowanii* comb. nov., *Kosakonia radincitans* comb. nov., *Kosakonia oryzae* comb. nov. and *Kosakonia arachidis* comb. nov., respectively, and *E. turicensis*, *E. helveticus* and *E. pulveris* into *Cronobacter* as *Cronobacter zurichensis* nom. nov., *Cronobacter helveticus* comb. nov. and *Cronobacter pulveris* comb. nov., respectively, and emended description of the genera *Enterobacter* and *Cronobacter*. *Systematic and Applied Microbiology.* 2013; 36(5):309–19. <https://doi.org/10.1016/j.syapm.2013.03.005> PMID: 23632228

79. Gouy M, Guindon S, Gascuel O. SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol.* 2010; 27(2):221–4. Epub 2009/10/27. <https://doi.org/10.1093/molbev/msp259> PMID: 19854763
80. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol.* 2015; 32(1):268–74. Epub 2014/11/06. <https://doi.org/10.1093/molbev/msu300> PMID: 25371430
81. Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E. CHOPCHOP: a CRISPR/Cas9 and TALEN web tool for genome editing. *Nucleic Acids Research.* 2014; 42(W1):W401–W7. <https://doi.org/10.1093/nar/gku410> PMID: 24861617
82. Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. CHOPCHOP v2: a web tool for the next generation of CRISPR genome engineering. *Nucleic Acids Research.* 2016; 44(W1):W272–W6. <https://doi.org/10.1093/nar/gkw398> PMID: 27185894
83. Trehan A, Kielbus M, Czapinski J, Stepulak A, Huhtaniemi I, Rivero-Müller A. REPLACR-mutagenesis, a one- step method for site-directed mutagenesis by recombineering. *Scientific Reports.* 2015:1–9. <https://doi.org/10.1038/srep19121> PMID: 26750263
84. Kozlova EV, Khajanchi BK, Popov VL, Wen J, Chopra AK. Impact of QseBC system in c-di-GMP-dependent quorum sensing regulatory network in a clinical isolate SSU of *Aeromonas hydrophila*. *Microbial pathogenesis.* 2012; 53(3–4):115–24. <https://doi.org/10.1016/j.micpath.2012.05.008> PMID: 22664750
85. O' Toole GA, Kolter R. Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* bio-film development. *Molecular Microbiology.* 1998; 30(2):295–304. <https://doi.org/10.1046/j.1365-2958.1998.01062.x> PMID: 9791175