






## Quantifying hole-motion-induced frustration in doped antiferromagnets by Hamiltonian reconstruction

Henning Schlömer<sup>1,2,3</sup>, Timon A. Hilker<sup>2,4</sup>, Immanuel Bloch<sup>1,2,4</sup>, Ulrich Schollwöck<sup>1,2</sup>, Fabian Grusdt<sup>1,2</sup> & Annabelle Bohrdt<sup>2,3,5,6</sup>

Unveiling the microscopic origins of quantum phases dominated by the interplay of spin and motional degrees of freedom constitutes one of the central challenges in strongly correlated many-body physics. When holes move through an antiferromagnetic spin background, they displace the positions of spins, which induces effective frustration in the magnetic environment. However, a concrete characterization of this effect in a quantum many-body system is still an unsolved problem. Here we present a Hamiltonian reconstruction scheme that allows for a precise quantification of hole-motion-induced frustration. We access non-local correlation functions through projective measurements of the many-body state, from which effective spin-Hamiltonians can be recovered after detaching the magnetic background from dominant charge fluctuations. The scheme is applied to systems of mixed dimensionality, where holes are restricted to move in one dimension, but SU(2) superexchange is two-dimensional. We demonstrate that hole motion drives the spin background into a highly frustrated regime, which can quantitatively be described by an effective  $J_1$ - $J_2$ -type spin model. We exemplify the applicability of the reconstruction scheme to ultracold atom experiments by recovering effective spin-Hamiltonians of experimentally obtained 1D Fermi-Hubbard snapshots. Our method can be generalized to fully 2D systems, enabling promising microscopic perspectives on the doped Hubbard model.

<sup>1</sup>Department of Physics and Arnold Sommerfeld Center for Theoretical Physics (ASC), Ludwig-Maximilians-Universität München, D-80333 München, Germany.

<sup>2</sup>Munich Center for Quantum Science and Technology (MCQST), D-80799 München, Germany. <sup>3</sup>ITAMP, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA. <sup>4</sup>Max-Planck-Institut für Quantenoptik, D-85748 Garching, Germany. <sup>5</sup>Department of Physics, Harvard University, Cambridge, MA 02138, USA. <sup>6</sup>Institut für Theoretische Physik, Universität Regensburg, D-93035 Regensburg, Germany. ✉email: [H.Schloemer@physik.uni-muenchen.de](mailto:H.Schloemer@physik.uni-muenchen.de); [Annabelle.Bohrdt@physik.uni-regensburg.de](mailto:Annabelle.Bohrdt@physik.uni-regensburg.de)

Microscopically understanding the motion of mobile charge carriers doped into Mott insulators constitutes one of the key open problems in strongly correlated many-body physics. When hopping through an insulating spin environment, holes displace spins along their way, which effectively frustrates the magnetic background. The arising competition of kinetic energy gain via delocalization and associated magnetic energy cost leads to the formation of a plethora of strongly interacting many-body phases<sup>1,2</sup>, many of which yet seek to be explained on a microscopic footing. The Fermi–Hubbard (FH) model, believed to capture the essential physics of strongly correlated materials, has been subject to intense numerical studies that can resolve the intricate competition between various orders<sup>3–9</sup>. Nevertheless, despite ongoing theoretical and experimental efforts over the past decades, a precise microscopic understanding of the interplay between motional and spin degrees of freedom is still an unsolved task, whose long-sought understanding may help to reveal the origin of high-temperature superconductivity and possibly lead to the discovery of novel pairing mechanisms<sup>10,11</sup>.

Analog quantum simulation, e.g., via ultracold atoms, can shed new light on the microscopic mechanisms underlying strongly correlated quantum many-body states<sup>12–17</sup> and paradigmatic Hamiltonians like the FH model can now be experimentally explored<sup>11,18–24</sup>. In particular, these setups allow to perform genuine quantum projective measurements and sample snapshots of the many-body state in the Fock basis, which in turn allow for insights into the wave function beyond averages and local observables. This capability has already been used to unveil highly non-local order parameters and hidden correlations in many-body systems<sup>22,25</sup>.

As we demonstrate in our work, the huge amount of information stored in snapshots of many-body states can further be utilized to disentangle spin and charge sectors through non-local correlation functions, which allow us to recover emergent effective spin-Hamiltonians for parts of the system. The problem of reconstructing a Hamiltonian from measured correlations via machine learning schemes<sup>26–30</sup> has attracted considerable interest in recent years, including certifying quantum simulation devices<sup>31</sup>.

In this article, we present a snapshot-based Hamiltonian reconstruction scheme for the spin channel alone, which removes dominant charge fluctuations<sup>22,32</sup> from individual snapshots. This allows us to quantify the effective spin-Hamiltonian, which includes the back-action of mobile dopants on the spin environment. We exemplify the proposed method by considering a system in mixed-dimensions (mixD), where hole motion is restricted to one dimension (1D), but SU(2) spin-superexchange is two-dimensional (2D). We find that hole hopping drives and stabilizes the spins in a highly frustrated regime, which we show to be accurately described by a  $J_1$ – $J_2$ -type spin-Hamiltonian.

Our method is directly applicable to experimental data obtained from ultracold quantum gas microscopes. We showcase this by reconstructing effective Hamiltonians from 1D measurements of the FH model<sup>22</sup>, where spin-charge separation governs the physics of the chains. Furthermore, our insights could be used to effectively simulate the highly frustrated  $J_1$ – $J_2$  model in ultracold atom experiments by implementing the mixD setting and post-processing the measurements.

Our work sheds light on the long-standing question about the interplay of spin- and motional degrees of freedom in strongly correlated materials, and paves the way to gain deep microscopic insights into prototypical systems such as the 2D FH and  $t$ – $J$  model.

## Results

**The model.** We consider the  $t$ – $J$  model in mixD<sup>10,33–35</sup>, described by the Hamiltonian:

$$\hat{H} = -t \sum_{(i,j),x,\sigma} \hat{P}_{GW} (\hat{c}_{i,\sigma}^\dagger \hat{c}_{j,\sigma} + \text{h.c.}) \hat{P}_{GW} + J \sum_{(i,j)} (\hat{S}_i \cdot \hat{S}_j - \frac{\hat{n}_i \hat{n}_j}{4}). \quad (1)$$


Here,  $\hat{c}_i^{(\dagger)}$ ,  $\hat{n}_i$  and  $\hat{S}_i$  are fermionic annihilation (creation), charge density, and spin operators on site  $i$ , respectively;  $\langle i, j \rangle_{(x)}$  denotes a nearest-neighbor (NN) pair on a 2D square lattice (with subscript  $x$  indicating a NN pair only along the  $x$ -direction), and  $\hat{P}_{GW}$  is the Gutzwiller operator projecting out states with double occupancy. The mixD setting, Eq. (1), has successfully been implemented in ultracold atom setups using strong tilted potential gradients<sup>11</sup>, which effectively restrict hole motion perpendicular to the gradient direction while spin-spin interactions remain 2D<sup>36,37</sup>.

Recently, we demonstrated how hidden AFM correlations in the mixD  $t$ – $J$  model result in the formation of a remarkably resilient stripe phase (i.e., a coupled charge- and spin-density wave<sup>38,39</sup>), with critical temperatures on the order of the magnetic coupling  $J$ <sup>35</sup>. Above these critical temperatures of charge- and spin-density wave formation, holes were found to form a deconfined chargon gas, i.e., a phase without order<sup>34,35</sup>.

In the following, we focus on the latter regime, and study how hole motion distorts the spins in the background. The effect is qualitatively depicted in Fig. 1a. The upper panel shows an (idealized) real space snapshot of holes moving through an AFM Néel background. Bonds correspond to AFM interactions in the instantaneous charge configuration, illustrated by gray lines. In between holes on neighboring legs, spins are aligned, leading to a linearly increasing magnetic energy penalty via the formation of geometric strings<sup>21,40–42</sup> (depicted by green wiggly lines).

As a direct consequence of the restricted charge motion to 1D, spins can be relabeled by the new positions they have after moving all holes to the right in each chain—resulting in a distinct definition of squeezed space<sup>32,43</sup>. More formally, consider a Fock state  $\bigotimes_y |\sigma_{[1,y]}, \sigma_{[2,y]}, \dots, \sigma_{[L_x,y]}\rangle$ , where  $\{\uparrow, \downarrow\} \ni \sigma_{x,y}$  is the single particle basis of the  $t$ – $J$  model. These local spin charge configurations are relabeled upon squeezing, whereby each Fock state is now given by  $\bigotimes_y |\tilde{\sigma}_{[1,y]}, \sigma_{[2,y]}, \dots, \sigma_{[L_x,y]}\rangle \otimes \hat{h}_{[x_1,y]}^\dagger \dots \hat{h}_{[x_{N_y},y]}^\dagger |0\rangle$ <sup>34</sup>. Here,  $\tilde{\sigma}_{[\tilde{x},y]} = \uparrow, \downarrow$  (but note that  $\tilde{\sigma}_{[\tilde{x},y]} \neq 0$ ) denotes spins on the squeezed lattice  $\tilde{x} = 1, \dots, L_x - N_y$ , where  $N_y$  is the number of holes in rung  $y$ , and  $\hat{h}_{[x,y]}^\dagger$  creates a hard core fermionic chargon at site  $\mathbf{i} = [x, y]$ . By squeezing the spins out, spins on the squeezed and real space lattice relate as  $\tilde{\sigma}(\tilde{x}, y) = \sigma(\tilde{x} + \sum_{j < \tilde{x}} n_{[j,y]}^h, y)$ , where  $n_{[x,y]}^h$  refers to the number of chargons at real space lattice site  $[x, y]$ . The lower panel of Fig. 1a illustrates the squeezing process, where the initial Néel order is restored in the isolated spin background. However, interactions on diagonal bonds emerge (ocher lines), which cause geometric frustration of the spins in squeezed space. From now on, we refer to lattice sites in real and squeezed space by  $\mathbf{i}$  and  $\tilde{\mathbf{i}}$ , respectively.

**Characterizing the spin state in squeezed space.** In order to quantify the arising frustration on the squeezed lattice, we simulate the mixD  $t$ – $J$  model, Eq. (1), at finite temperature and fixed doping using imaginary time evolution schemes (purification) via matrix product states (MPS)<sup>44,45</sup>. For faster, more controllable numerics and to prevent post-selection of snapshots, we explicitly implement the system's enhanced U(1) symmetries in each ladder leg, i.e., we work in an ensemble where we allow for thermal spin fluctuations but keep the number of holes in each ladder leg constant<sup>35</sup>. In particular, we simulate the mixD  $t$ – $J$  model at intermediate temperature  $T/J = 5/3$  ( $\beta J = 0.6$ ), which lies inside the chargon gas phase (i.e., no charge- and spin-density waves form) and, furthermore, is in a temperature regime accessible for quantum gas microscopes.



**Fig. 1** Hole-motion-induced spin frustration. **a** Schematic of how hole hopping induces frustration in the spin background. Upper panel: Snapshot of holes moving through a Néel background. Spatial separation of holes on neighboring legs lead to the formation of geometric strings (green wiggly lines) costing magnetic energy. Lower panel: Upon transforming the snapshot to squeezed space, originally vertical bonds  $J_1^x$  in between two holes on neighboring legs become effective diagonal couplings  $J_2$ . The resulting energy penalty of aligned diagonal spins leads to frustration in the magnetic background. **b** Hamiltonian reconstruction results (blue) for hole dopings  $n^h = 0.05 \dots 0.2$  of a mixD  $t - J$  ladder with  $t/J = 3$ ,  $L_x \times L_y = 20 \times 2$  and  $T/J = 5/3 - 1.67$ . Reconstructions of squeezed space according to the input  $J_1^x - J_1^y - J_2$  Heisenberg Hamiltonian, Eq. (2), are presented in a  $J_1^y/J_1^x, J_2/J_1^x$  diagram. Light regions in the background signal the presence of either AFM or stripe AFM order in the purely magnetic  $J_1^x - J_1^y - J_2$  model in the ground state by plotting the sum of the spin structure factors  $S(\pi, \pi) + S(0, \pi)$ . Dark regions correspond to a highly frustrated regime without apparent order. Upon doping the system, the background spins are driven into a strongly frustrated state. Error bars correspond to the standard error to the mean when averaging over ten reconstruction runs. Red connected symbols show theoretical expectations assuming no spin-hole correlations in the mixD  $t - J$  model, i.e.,  $\hat{\rho} = \hat{\rho}_s \otimes \hat{\rho}_c$ , evaluated via Eqs. (4) and (5).

From the thermal MPS at inverse temperature  $\beta$ , we sample uncorrelated snapshots of the corresponding Gibbs state<sup>46,47</sup>. After post-processing the individual measurements by squeezing out the holes, spin-spin correlations  $\langle \hat{S}_i^z \hat{S}_j^z \rangle$  can directly be evaluated in squeezed space. In Fig. 2a, nearest-neighbor as well as diagonal spin-spin correlations are shown on the squeezed lattice. In the bulk of the squeezed ladder, both nearest neighbor as well as diagonal correlators are negative and comparable in magnitude, signaling strong frustration in the spin background induced by the motion of the holes. In contrast, at both edges of the ladder, low average hole concentrations lead to only marginal perturbations of the spin background—resulting in AFM type correlations that are negative (positive) along nearest (diagonal) neighbors.

Figure 2b shows nearest-neighbor rung (blue) and diagonal (red) correlators. Dashed-dotted lines correspond to rung and diagonal correlations for a Heisenberg ladder with solely nearest-neighbor couplings (where  $\beta J_1^x = \beta J_1^y = 0.44$ ), which describe the physics at the edges qualitatively well, but fail to reproduce the measured correlations in the bulk. Due to the frustrating effect of the hopping holes, diagonal couplings need to be taken into

account to accurately capture the physics of the squeezed background. To this end, we introduce an effective  $J_1^x - J_1^y - J_2$  Heisenberg model<sup>48</sup>, given by the Hamiltonian:

$$\hat{\mathcal{H}}_{\{J_H\}} = \sum_{\mu=x,y} J_1^\mu \sum_{\langle \hat{i}, \hat{j} \rangle_\mu} \hat{S}_i \cdot \hat{S}_j + J_2 \sum_{\langle \hat{i}, \hat{j} \rangle_{\text{diag}}} \hat{S}_i \cdot \hat{S}_j. \quad (2)$$

Here,  $\hat{S}_i$  is the spin-1/2 operator at site  $\hat{i}$ , and  $\{J_H\} = \{J_1^x, J_1^y, J_2\}$  are the coupling strengths of neighboring spins in  $x, y$  and diagonal direction on the squeezed lattice, respectively.

To quantitatively pin down the strength of the arising frustration, we perform a Hamiltonian reconstruction with input Hamiltonian  $\hat{\mathcal{H}}_{\{J_H\}}$ , Eq. (2), together with the measured spin-spin correlations in squeezed space. Couplings in  $\hat{\mathcal{H}}_{\{J_H\}}$  are chosen to be homogeneous throughout the bulk of squeezed space, justified by the approximately constant behavior of correlations in the bulk region of Fig. 2b—solid lines are averages of the correlations over the marked box. Due to the underlying SU(2) symmetry of the mixD  $t - J$  model, many-body snapshots along a single spin axis—here chosen along  $z$ —are sufficient to reconstruct the full effective Hamiltonian, Eq. (2). Results of the reconstruction correspond to the parameter configuration  $\{J_H\}$  that best describe the measured correlations.

On a more formal footing, we follow the procedure introduced in ref.<sup>30</sup> and minimize the objective function  $\mathcal{G}$  over all possible coupling parameters  $\{J_H\}$ :


$$\mathcal{G} = \ln Z(\beta, \{J_H\}) + 3\beta \left( \sum_{\mu=x,y} J_1^\mu \mathcal{M}_1^\mu + J_2 \mathcal{M}_2 \right), \quad (3)$$

with  $Z = \text{Tr}[e^{-\beta \hat{\mathcal{H}}_{\{J_H\}}}]$  the partition function and  $\mathcal{M}_1^\mu = \sum_{\langle \hat{i}, \hat{j} \rangle_\mu} \langle \hat{S}_i^z \hat{S}_j^z \rangle$ ,  $\mathcal{M}_2 = \sum_{\langle \hat{i}, \hat{j} \rangle_{\text{diag}}} \langle \hat{S}_i^z \hat{S}_j^z \rangle$  the summed correlations along nearest- and diagonal neighbors within the considered window in the bulk of squeezed space. Figure 2c shows how approximations for  $\mathcal{M}_1^{x,y}, \mathcal{M}_2$  quickly saturate with the number of used snapshots, suggesting a qualitatively satisfactory proxy for the spin-spin correlators after a few thousand projective measurements. For the rest of the analysis, we use sample sizes of 7000 snapshots for each approximation of the correlations.

The minimization process is done via standard gradient descent (GD) methods, where in each iteration the parameters are updated according to the gradient  $\nabla \mathcal{G}$  within the considered bulk window of squeezed space. The temperature  $\beta^{-1}$  of the  $J_1^x - J_1^y - J_2$  Heisenberg Hamiltonian is chosen identically to the underlying simulations of the mixD  $t - J$  system during the GD. Note that this choice might not reflect the actual effective temperature of the spin background. However, the relevant ratios  $J_1^y/J_1^x, J_2/J_1^x$  that quantify the frustration in the system are independent of the true temperature of the squeezed magnetic environment.

Intermediate temperature regimes  $T/J \gtrsim 1$ —as also chosen in our simulations—have been shown to work best for reconstructions of the underlying coupling parameters, as both in the low and high-temperature limit the energy landscape defined by  $\mathcal{G}$  is entirely flat<sup>30</sup>. Given a size  $L_x \times L_y$  of the mixD system, the dimensions of the reconstructed  $J_1^x - J_1^y - J_2$  Heisenberg ladder on the squeezed lattice is given by  $\tilde{L}_x \times \tilde{L}_y = (1 - n^h)L_x \times L_y$ .

Hamiltonian reconstruction results for a single run are presented in Fig. 2d. Evaluated correlations of the best fitting  $J_1^x - J_1^y - J_2$  model are seen to perfectly match the measured mean correlations in the bulk of squeezed space, hence strongly supporting that the physics of the magnetic background in the mixD  $t - J$  model is well captured by  $J_1^x - J_1^y - J_2$  Heisenberg interactions on a square lattice. We have explicitly checked that independent of the initially chosen parameter values for the GD,



**Fig. 2 Correlations in squeezed space and Hamiltonian reconstruction.** **a** Spin-spin correlations  $\langle \hat{S}_i^x \hat{S}_j^x \rangle$  on the squeezed lattice of an original  $20 \times 2$  mixD  $t - J$  ladder, with  $n^h = 0.2$ ,  $T/J = 5/3 - 1.67$ , and using 20,000 snapshots. In the bulk of squeezed space, hole motion distorts the spin background, leading to negative correlations across diagonals. In this region, effective  $J_1 - J_2$  physics is expected, as captured by the Hamiltonian Eq. (2). As holes are rarely located at the open boundaries of the system, correlations are left almost undisturbed and are of AFM type. Correlations along nearest neighbors in  $x$  go beyond the cutoff of the colorbar. **b** Rung  $C_1^x(\tilde{x}) = \langle \hat{S}_{[\tilde{x},0]}^x \hat{S}_{[\tilde{x},1]}^x \rangle$  and diagonal  $C_2^x(\tilde{x}) = \langle \hat{S}_{[\tilde{x},0]}^x \hat{S}_{[\tilde{x}+1,1]}^x \rangle + \langle \hat{S}_{[\tilde{x}+1,0]}^x \hat{S}_{[\tilde{x},1]}^x \rangle$  correlations. In the central bulk region of the ladder, correlations are approximately constant, the average being used as the input for  $J_1^x - J_1^y - J_2$  Hamiltonian reconstructions. In particular, we discard the two outer sites in squeezed space, as illustrated by the yellow box. Dashed-dotted lines correspond to rung and diagonal correlations for a nearest-neighbor Heisenberg model with  $\beta J_1^x = \beta J_1^y \sim 0.44$ , which captures the physics at the edges, but fails to describe the correlations in the bulk of squeezed space. The introduction of diagonal (frustrating) bonds is hence an essential step to describe the spin system on the squeezed lattice. **c** Summed correlations in the boxed bulk in (b) along rungs, legs and diagonals for varying snapshot set sizes. Light regions correspond to the standard error to the mean. After a few thousand measurements, convergence of the correlator proxies is reached. **d** Results show perfect agreement between the correlations emerging from a reconstructed effective  $J_1^x - J_1^y - J_2$  Heisenberg model (black crosses) and bulk averaged correlations of the doped mixD model in squeezed space (solid lines).

$\{J_H\}$  always converge to identical points in parameter space, underlining the robustness of the GD scheme—see Supplementary Note 1.

To characterize and classify the reconstructed spin states in squeezed space as a function of doping, we perform ground state calculations of the  $J_1^x - J_1^y - J_2$  Heisenberg model and evaluate the static spin structure factor (SSSF) given by  $S(q_x, q_y) = \frac{1}{L_x L_y} \sum_{i,j} e^{iq \cdot (i-j)} \langle \hat{S}_i \cdot \hat{S}_j \rangle$ . For  $J_1^x = J_1^y = J_1$ , it has been demonstrated that a highly frustrated magnetic regime exists for  $0.4 \lesssim J_2/J_1 \lesssim 0.6$  that is sandwiched by a Néel and stripe AFM phase<sup>49–54</sup>. Though the exact nature of the non-magnetic ground state in the frustrated regime is still controversial, it remains a promising candidate for the realization of a quantum spin liquid phase possibly described by Anderson’s resonating valence bond (RVB) paradigm<sup>55–62</sup>. We evaluate the hybrid order parameter  $S(\pi, \pi) + S(0, \pi)$  in the  $J_1^y/J_1^x - J_2/J_1^x$  parameter space, signaling whether AFM or stripe AFM order exists in the system. Dark regions in the background of Fig. 1b correspond to no apparent spin ordering, and hence signal the existence of a strongly frustrated spin state akin to the observations in the homogeneous  $J_1 - J_2$  model.

The reconstruction process, consisting of (1) approximating correlations in squeezed space using snapshots and (2) performing the GD, is repeated a total number of ten times. Averaging over the converged results of all runs leads to the main result of this paper, presented in Fig. 1b by blue connected symbols. We observe how the spin state in squeezed space rapidly approaches the highly frustrated regime upon increasing the doping level, until seemingly saturating within it to a certain configuration  $\{J_H^s\}$ . We note that at the considered system sizes, boundary effects become especially pronounced at low hole concentrations  $n^h = 0.05, 0.1$ . This leads to a slow saturation of the correlations in the bulk of squeezed space, which in turn shifts effective couplings averaged within the fixed window to smaller (larger) values of  $J_2/J_1^x$  ( $J_1^y/J_1^x$ ), see Supplementary Note 2. In the

thermodynamic limit, we in fact expect any finite hole doping in the chargin gas phase to drive the squeezed spin system into a highly frustrated state. A numerical study of longer ladders of size  $40 \times 2$  support this assumption, where already for  $n^h = 0.05$  the spin state is reconstructed to lie deep inside the frustrated regime.

The reconstruction scheme as introduced above takes into account spin-spin correlations directly measured in squeezed space, hence providing an unbiased platform for the analysis of the spin background by explicitly including the back-action of hole motion on the spins. Motivated from the separation of energy scales in the mixD  $t - J$  model with  $t/J \gg 1$ , we make the ansatz of a fully decoupled thermal density matrix given by separate spin (s) and charge (c) sectors,  $\hat{\rho} = \hat{\rho}_s \otimes \hat{\rho}_c$ , and aim to test the resulting predictions against the unbiased reconstruction output.

Within the separation ansatz, interaction strengths in squeezed space are obtained by conditioned probabilities in real space<sup>22</sup>. Two nearest neighbors along  $x$  in squeezed space interact only if the corresponding sites are nearest neighbors along  $x$  in real space, leading to an effective coupling strength (assuming homogeneous hole density  $\langle n_i^h \rangle = n^h$ )—see i.p. the Supplementary Materials of ref. 22:

$$J_n^x/J \propto \langle (1 - \hat{n}_i^h)(1 - \hat{n}_{i+e_x}^h) \rangle = 1 - 2n^h + g_x^{(2)}. \quad (4)$$

Here,  $g_\mu^{(2)} = \langle \hat{n}_i^h \hat{n}_{i+e_\mu}^h \rangle$  with  $e_\mu$  the unit vector in direction  $\mu = x, y$ . Vertical and diagonal bonds are obtained similarly by conditioning the correlators by the total number of holes to the left of site  $i$ ,  $\nu_{i=[x,y]}^h = \sum_{x' < x} N_{[x',y]}^h$ , with  $N_i^h$  the number of holes on site  $i$ . Diagonal coupling strengths  $J_n$  spanning a distance of  $\Delta x = n - 1$  (vertical bonds  $J_1^y$  correspond to  $J_1$  in this notation) are then given by:

$$J_n/J \propto \left\langle (1 - \hat{n}_i^h)(1 - \hat{n}_{i+e_y}^h) \right\rangle_{|\nu_i^h - \nu_{i+e_y}^h| = n-1}. \quad (5)$$




We evaluate the estimated effective couplings, Eqs. (4) and (5), using the mixD  $t-J$  snapshots. Results are shown by red connected symbols in Fig. 1b. We observe that the theoretically predicted expectations for  $J_1^x, J_1^y, J_2$  within the separation ansatz agree remarkably well with the full reconstruction. Deviations from the above description, in particular the consistent underestimation of relative diagonal coupling strengths  $J_2/J_1^x$ , are likely caused by non-trivial spin-hole correlations in the mixD  $t-J$  model, which are implicitly included in the reconstruction analysis but discarded in the separation ansatz.

We note that the conditioned correlators Eqs. (4) and (5) calculated from mixD  $t-J$  snapshots are numerically almost identical to calculations of free spinless fermions (free chargon gas), see Supplementary Note 3. This lets us conclude that holes, while behaving like free fermions in the chargon gas phase of the mixD  $t-J$  model, nevertheless correlate non-trivially with the spin background.

So far, our approach has been to restrict the effective Hamiltonian Eq. (2) to first order diagonal couplings  $J_2$ . To assess the systematic error related to this approximation, we estimate the magnitude of longer-range couplings  $J_n$  by evaluating the conditioned correlators Eq. (5) for  $n \geq 3$ . Relative strengths of couplings up to  $J_4$  are depicted in Fig. 3, where a rapid decrease with real space distance is observed. The relative strength of  $J_2$  in units of  $J_1^x$  is of order  $\sim 10\%$ , cf. Fig. 1b.  $J_3$ , on the other hand, reaches only a few percent in terms of  $J_1^x$ , suggesting that  $n \geq 3$  couplings are negligible for the effective description. We implement the Hamiltonian reconstruction scheme outlined above also for a  $J_1^x - J_1^y - J_2 - J_3$  Heisenberg model including interactions in squeezed space up to  $J_3$ . We find that this results in only very minor corrections to the reconstructed ratios  $J_1^y/J_1^x, J_2/J_1^x$ , supporting that the spin physics is well captured by nearest-neighbor and frustrating (first order) diagonal bonds—see Supplementary Note 4.

**Spin-charge separation in 1D.** In the 1D FH model, the ground state wave function is known to factorize into fully separated spin and charge channels in the strongly interacting limit, leading to the celebrated phenomenon of spin-charge separation (i.e., the exact absence of spin-hole correlations)<sup>43</sup>. In 1D, it has been demonstrated that hidden spin correlations—distorted in real space by the motion of holes—can be revealed by transformation to squeezed space, effectively described by a 1D Heisenberg Hamiltonian with nearest neighbor interaction  $J_1^x(n^h)$  on the squeezed lattice<sup>22,63</sup>, cf. Fig. 4a:

$$\hat{H}_{J_1^x} = J_1^x \sum_{\langle ij \rangle} \hat{S}_i \cdot \hat{S}_j. \quad (6)$$



**Fig. 3 Significance of longer ranged couplings.** By evaluating the conditioned correlators, Eqs. (4) and (5), we estimate the strengths of longer ranged couplings up to  $J_4$ . In units of the strongest interaction  $J_1^x$ , first order diagonal bonds  $J_2$  are of the order of  $\sim 10\%$ , whereas couplings  $J_3$  reach relative magnitudes of a few percent. Due to the finite system size,  $J_3$  ( $J_4$ ) and higher order couplings drop to zero for  $n^h = 0.05$  ( $n^h = 0.1$ ), as the corresponding conditioned probabilities Eq. (5) vanish for a single (two) hole(s) per leg.

We apply our squeezed space Hamiltonian reconstruction scheme and recover the effective spin-Hamiltonian of the doped 1D FH model using experimentally obtained snapshots in a degenerate two-component ultracold Fermi gas carried out by some of us<sup>22</sup>. Results are shown in Fig. 4b by blue data points, where a consistent decrease of effective coupling strength  $J_1^x(n^h)$  is observed upon increasing the hole doping—as expected from Eq. (4). For comparison, we further simulate the 1D  $t-J$  model with identical parameters as estimated in the experiment ( $t/J = 1.82$ ,  $T/J = 0.87$ ) and use sampled thermal snapshots in squeezed space for reconstructions, shown by red squares in Fig. 4b.

Effective interactions  $J_1^x/J$  reconstructed for an underlying 1D  $t-J$  model are seen to consistently lie below recovered coupling strengths of the 1D FH model. This discrepancy can be explained by higher order virtual processes in the FH model, which we illustrate by comparing the reconstructions to theoretical predictions within a separation ansatz. In the 1D  $t-J$  model, effective spin interactions in squeezed space can be calculated via Eq. (4), yielding:


$$J_1^x/J = 1 - n^h - \frac{1}{1 - n^h} [G(1)]^2. \quad (7)$$

Here,  $G(d) = \frac{1}{\pi} \int_0^\pi dk \cos(kd) n_F(n^h, T)$  with  $n_F(n^h, T)$  the Fermi-Dirac distribution of free chargons hopping on a 1D lattice at temperature  $T$ . When generalizing the  $t-J$  model to include next-nearest neighbor hole hopping processes mediated by doubly occupied virtual states as possible in the FH model, the effective coupling reads<sup>22</sup>:

$$J_1^x/J = 1 - n^h + G(2), \quad (8)$$

with  $J = 4t^2/U$  and  $U$  the Hubbard interaction.

Reconstructed values of  $J_1^x/J$  are observed to match the theoretical predictions for spin-charge separated systems well, depicted by red and blue dashed lines corresponding to Eqs. (7) and (8), respectively. This illustrates the (approximate) presence



**Fig. 4 Reconstructing 1D systems from experiments.** **a** Illustration of snapshots of the 1D FH model in real (top) and squeezed (bottom) space. **b** Evaluation of 1D FH snapshots of a cold atom experiment<sup>22</sup>. Reconstructions of the effective spin-Hamiltonian Eq. (6) in squeezed space for varying hole densities are shown by blue data points. Red data points correspond to reconstructions of the 1D  $t-J$  model, which we simulate using MPS for parameters as estimated in ref. <sup>22</sup>, i.e.,  $t/J = 1.82$  and  $T/J = 0.87$ . Results are compared to theoretical predictions (dashed lines) assuming spin-charge separation, Eqs. (7) and (8), showing a good match with the reconstructed data. In particular, higher order hopping processes lead to the FH measurement reconstructions of  $J_1^x/J$  to consistently lie above predictions for the  $t-J$  model. Error bars are too small to be visible for the  $t-J$  reconstructions on the scale of the plot. The  $T = \infty$  limit is shown by the gray dashed line, where a linear decrease  $J_1^x = 1 - n^h$  is expected for both the FH and  $t-J$  model.

of spin-charge separation in the 1D FH and  $t - J$  model away from the  $T = 0$  and strongly interacting limit, ultimately being mediated by their separation of energy scales.

## Discussion

Using Hamiltonian reconstruction schemes, we have proposed a method to quantify hole-motion-induced frustration in a doped antiferromagnet by exploiting the full information stored in many-body snapshots. An advantage of the reconstruction process as introduced above is that the effective Hamiltonian—defined on the squeezed lattice—describes a reduced number of degrees of freedom, i.e., its local Hilbert space dimension is smaller than the one of the original system. In particular, the effective spin-Hamiltonian in squeezed space is of local dimension  $d = 2$ , rendering reconstructions for a given set of snapshots feasible even for larger system sizes. Experimental data of 2D systems that are inaccessible with classical simulations but within reach of current experiments<sup>11</sup> could be used as input for a computational reconstruction of the spin background. We benchmarked this for wide Heisenberg ladders, where we reconstruct unknown coupling parameters to high accuracy from snapshots, see Supplementary Note 5.

By analyzing a setting in mixed-dimensions, we have firmly established a quantitative connection between the doped mixD  $t - J$  model and the paradigmatic frustrated  $J_1 - J_2$  model. In particular, we demonstrated how hole motion drives the spin background into a highly frustrated state, whereby effective diagonal, frustrating magnetic bonds are induced on the squeezed lattice formed by the spins alone. Our results match theoretical predictions based on spin-charge separation reasonably well, differences being likely caused by weak remaining spin-hole correlations that deserve further investigation in the future. We note that due to the formation of stripes below temperatures  $T \sim J/2$ <sup>35</sup> in the mixD  $t - J$  model, its ground state is not directly related to a quantum spin liquid phase. Nevertheless, the ordered stripe phase at lower temperature may be merely covering a disordered quantum phase that dominates the physics of the model once the stripe order is melted away above the stripe critical temperature. Though this perspective is admittedly speculative, a similar view is often evoked in the context of describing the pseudogap with its associated small Fermi surface as originating from a covered ground state fractionalized Fermi liquid, see e.g. ref. 64.

Utilizing snapshots of a cold atom experiment simulating the 1D FH model, we already demonstrated the direct applicability of the reconstruction method to existing experimental data. From a converse experimental point of view, the above insights could further be utilized to effectively simulate the highly frustrated  $J_1 - J_2$  model by implementing the mixD setting and post-processing the measurements.

The reconstruction scheme can be generalized and applied to a variety of many-body phases. In the stripe phase, for instance, fluctuating holes bound into stripes are expected to lead to spatial modulations of the couplings between spins which can be reconstructed using the scheme we described. Moreover, our method can be extended from mixD to fully 2D settings with homogeneous charge motion, where e.g., a weak easy-axis anisotropy of the Heisenberg interactions can enable string retracing<sup>40</sup> to remove dominant charge fluctuations and define a squeezed lattice. Applying our scheme to such snapshots will provide a microscopic perspective on the doped FH model and its relation to putative topological order in his enigmatic model. Making explicit use of all accessible correlation functions in squeezed space to further enhance the accuracy of the reconstructions is a promising

direction for future research, for instance by directly comparing the distributions of measured and reconstructed snapshots.

## Methods

**Finite-temperature DMRG.** We simulate the mixD  $t - J$  model at finite temperature using mixed state purification schemes while conserving the system's symmetries<sup>65</sup>. In particular, we expand the ladder system by introducing auxiliary sites, which act as a finite temperature bath via their entanglement to the physical system. In order to calculate thermal matrix product states, we first generate the infinite temperature, maximally entangled state  $|\Psi(\beta = 0)\rangle$  in a given symmetry sector<sup>35</sup>—see also Supplementary Note 6.

A pure state in the enlarged system at finite temperature is then calculated by evolving  $|\Psi(\beta = 0)\rangle$  in imaginary time under the physical Hamiltonian,  $|\Psi(\tau)\rangle = e^{-\tau H} |\Psi(\beta = 0)\rangle$ , where  $\tau = \beta/2$  with  $\beta$  the inverse temperature. The corresponding mixed state of the physical system is computed by tracing out all auxiliary degrees of freedom when computing expectation values in the physical subset.

During the imaginary time evolution, we conserve the particle number in each physical leg  $N_\ell$ ,  $\ell = 1 \dots L_y$ , the total particle number in the auxiliary system  $N_{\text{aux}}^{\text{tot}}$ , as well as the total spin  $S_{\text{phys.+aux.}}^z$  (the latter allowing for finite total magnetizations of the physical system at finite temperature). This results in a total of  $L_y + 2$  symmetries employed by the DMRG implementation.

Given a generic observable  $\hat{O}$  of the physical chain, the thermodynamic average can be calculated in the enlarged space by tracing out the ancilla degrees of freedom:

$$\langle \hat{O} \rangle_\beta = \frac{\langle \Psi(\beta) | \hat{O} | \Psi(\beta) \rangle}{\langle \Psi(\beta) | \Psi(\beta) \rangle}. \quad (9)$$

Here, the norm  $\langle \Psi(\beta) | \Psi(\beta) \rangle \propto Z(\beta)$  is proportional to the partition function at temperature  $\beta^{-1}$ .

The maximally entangled state needed as a starting point of the imaginary time evolution is generated using the concept of entangler Hamiltonians<sup>65,66</sup>, which we specifically tailor for our “leg-canonical” ensemble<sup>35</sup>. Since the maximally entangled state is usually of low bond dimension, we first employ global MPS imaginary time evolution schemes to evolve the system away from infinite temperature. Once bond dimensions are sufficiently high, we switch to local approximation methods. In particular, we use the Krylov scheme and the time-dependent variational principle (TDVP) for global and local evolutions, respectively<sup>45</sup>.

**Hamiltonian reconstruction.** Due to the non-Markovian nature of quantum states, it is a priori unclear whether measured correlations are sufficient to learn the quantum interactions of the underlying Hamiltonian<sup>67</sup>. However, it has been shown that the strongly convex property of the free energy with respect to the interaction parameters renders the Hamiltonian learning problem feasible<sup>30</sup>.

In each GD step, we compute the partition function and relevant correlations  $\mathcal{M}_1^{x/y}$ ,  $\mathcal{M}_2$  using the MPS schemes described above. Due to the numerical complexity, we do not consider advanced GD methods with varying step size e.g., given by the Amijo rule, but stick to a straightforward optimization using a fixed descent step. In particular, we choose the step  $\mathbf{a}$  to be 20% of the objective gradient, i.e.,  $\mathbf{a} = 0.2 \nabla \mathcal{G}$ , where  $\nabla'$  is the gradient in parameter space within the fixed window in the bulk of squeezed space as introduced in the main text. When the norm of the gradient reaches a certain threshold, here chosen as  $|\nabla \mathcal{G}| < 10^{-6}$ , we stop the descent and assume converged results.

In our simulations, we work at intermediate temperatures, i.p.  $T/J = 5/3$ . On the one hand, this ensures that the mixD  $t - J$  system is in the chargin gas phase, i.e., stripes do not form<sup>35</sup>. On the other hand, intermediate temperature regimes have been shown to yield best reconstruction results from projective measurements<sup>30</sup>. To illustrate the latter argument, consider for instance the ferromagnetic (FM) Ising model, featuring a FM ground state for any non-zero interaction strength. Therefore, at low temperatures close to the ground state, the energy landscape  $\mathcal{G}$  is nearly flat, resulting in bad reconstructions. Similarly, for  $T/J \gg 1$  measured correlations only weakly depend on the underlying coupling parameters (e.g., the infinite temperature state is identical for all interaction strengths), which hinder precise reconstructions. By reconstructing purely magnetic models for various temperatures, we demonstrate this explicitly in Supplementary Note 5.

## Data availability

The datasets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Code availability

The data analyzed in the current study have been obtained using the SyTen package<sup>68,69</sup>.

Received: 24 March 2023; Accepted: 10 July 2023;

Published online: 23 August 2023

## References

- Lee, P. A., Nagaosa, N. & Wen, X.-G. Doping a mott insulator: physics of high-temperature superconductivity. *Rev. Mod. Phys.* **78**, 17–85 (2006).
- Keimer, B., Kivelson, S. A., Norman, M. R., Uchida, S. & Zaanen, J. From quantum matter to high-temperature superconductivity in copper oxides. *Nature* **518**, 179–186 (2015).
- LeBlanc, J. P. F. et al. Solutions of the two-dimensional Hubbard model: benchmarks and results from a wide range of numerical algorithms. *Phys. Rev. X* **5**, 041041 (2015).
- Zheng, B.-X. et al. Stripe order in the underdoped region of the two-dimensional Hubbard model. *Science* **358**, 1155–1160 (2017).
- Jiang, H.-C. & Devereaux, T. P. Superconductivity in the doped hubbard model and its interplay with next-nearest hopping  $t'$ . *Science* **365**, 1424–1428 (2019).
- Qin, M. et al. Absence of superconductivity in the pure two-dimensional Hubbard model. *Phys. Rev. X* **10**, 031016 (2020).
- Schäfer, T. et al. Tracking the footprints of spin fluctuations: a multimethod, multimessenger study of the two-dimensional Hubbard model. *Phys. Rev. X* **11**, 011058 (2021).
- Jiang, H.-C. & Kivelson, S. A. Stripe order enhanced superconductivity in the Hubbard model. *Proc. Natl Acad. Sci.* **119**, e2109406119 (2022).
- Arovas, D. P., Berg, E., Kivelson, S. A. & Raghu, S. The hubbard model. *Annu. Rev. Condens. Matter Phys.* **13**, 239–274 (2022).
- Bohrdt, A., Homeier, L., Bloch, I., Demler, E. & Grusdt, F. Strong pairing in mixed-dimensional bilayer antiferromagnetic mott insulators. *Nat. Phys.* **18**, 651–656 (2022).
- Hirthe, S. et al. Magnetically mediated hole pairing in fermionic ladders of ultracold atoms. *Nature* **613**, 463–467 (2023).
- Bloch, I., Dalibard, J. & Zwirger, W. Many-body physics with ultracold gases. *Rev. Mod. Phys.* **80**, 885–964 (2008).
- Bloch, I., Dalibard, J. & Nascimbène, S. Quantum simulations with ultracold quantum gases. *Nat. Phys.* **8**, 267–276 (2012).
- Cheuk, L. W. et al. Quantum-gas microscope for fermionic atoms. *Phys. Rev. Lett.* **114**, 193001 (2015).
- Parsons, M. F. et al. Site-resolved imaging of fermionic  ${}^6\text{Li}$  in an optical lattice. *Phys. Rev. Lett.* **114**, 213002 (2015).
- Gross, C. & Bloch, I. Quantum simulations with ultracold atoms in optical lattices. *Science* **357**, 995–1001 (2017).
- Schäfer, F., Fukuhara, T., Sugawa, S., Takasu, Y. & Takahashi, Y. Tools for quantum simulation with ultracold atoms in optical lattices. *Nat. Rev. Phys.* **2**, 411–425 (2020).
- Esslinger, T. Fermi-hubbard physics with atoms in an optical lattice. *Annu. Rev. Condens. Matter Phys.* **1**, 129–152 (2010).
- Hart, R. A. et al. Observation of antiferromagnetic correlations in the Hubbard model with ultracold atoms. *Nature* **519**, 211–214 (2015).
- Cocchi, E. et al. Equation of state of the two-dimensional Hubbard model. *Phys. Rev. Lett.* **116**, 175301 (2016).
- Chiu, C. S. et al. String patterns in the doped Hubbard model. *Science* **365**, 251–256 (2019).
- Hilker, T. A. et al. Revealing hidden antiferromagnetic correlations in doped Hubbard chains via string correlators. *Science* **357**, 484–487 (2017).
- Koepsell, J. et al. Microscopic evolution of doped Mott insulators from polaronic metal to fermi liquid. *Science* **374**, 82–86 (2021).
- Bohrdt, A., Homeier, L., Reinmoser, C., Demler, E. & Grusdt, F. Exploration of doped quantum magnets with ultracold atoms. *Ann. Phys.* **435**, 168651 (2021).
- Endres, M. et al. Observation of correlated particle-hole pairs and string order in low-dimensional Mott insulators. *Science* **334**, 200–203 (2011).
- Di Franco, C., Paternostro, M. & Kim, M. S. Hamiltonian tomography in an access-limited setting without state initialization. *Phys. Rev. Lett.* **102**, 187203 (2009).
- Zhang, J. & Sarovar, M. Quantum hamiltonian identification from measurement time traces. *Phys. Rev. Lett.* **113**, 080401 (2014).
- Qi, X.-L. & Ranard, D. Determining a local Hamiltonian from a single eigenstate. *Quantum* **3**, 159 (2019).
- Cao, C., Hou, S.-Y., Cao, N. & Zeng, B. Supervised learning in Hamiltonian reconstruction from local measurements on eigenstates. *J. Phys. Condens. Matter* **33**, 064002 (2020).
- Anshu, A., Arunachalam, S., Kuwahara, T. & Soleimanifar, M. Sample-efficient learning of interacting quantum systems. *Nat. Phys.* **17**, 931–935 (2021).
- Bairey, E., Arad, I. & Lindner, N. H. Learning a local Hamiltonian from local measurements. *Phys. Rev. Lett.* **122**, 020504 (2019).
- Kruis, H. V., McCulloch, I. P., Nussinov, Z. & Zaanen, J. Geometry and the hidden order of luttinger liquids: the universality of squeezed space. *Phys. Rev. B* **70**, 075109 (2004).
- Grusdt, F., Zhu, Z., Shi, T. & Demler, E. Meson formation in mixed-dimensional t-J models. *SciPost Phys.* **5**, 57 (2018).
- Grusdt, F. & Pollet, L.  $z_2$  parton phases in the mixed-dimensional  $t - J_z$  model. *Phys. Rev. Lett.* **125**, 256401 (2020).
- Schlömer, H., Bohrdt, A., Pollet, L., Schollwöck, U. & Grusdt, F. Robust stripes in the mixed-dimensional  $t - j$  model. *Phys. Rev. Res.* **5**, L022027 (2023).
- Trotzky, S. et al. Time-resolved observation and control of superexchange interactions with ultracold atoms in optical lattices. *Science* **319**, 295–299 (2008).
- Dimitrova, I. et al. Enhanced superexchange in a tilted Mott insulator. *Phys. Rev. Lett.* **124**, 043204 (2020).
- White, S. R. & Scalapino, D. J. Density matrix renormalization group study of the striped phase in the 2d  $t - J$  model. *Phys. Rev. Lett.* **80**, 1272–1275 (1998).
- Tranquada, J. M. Stripes and superconductivity in cuprates. *Physica B: Condensed Matter* **407**, 1771–1774 (2012).
- Grusdt, F. et al. Parton theory of magnetic polarons: mesonic resonances and signatures in dynamics. *Phys. Rev. X* **8**, 011046 (2018).
- Grusdt, F., Bohrdt, A. & Demler, E. Microscopic spinon-charge theory of magnetic polarons in the  $t - j$  model. *Phys. Rev. B* **99**, 224422 (2019).
- Bohrdt, A. et al. Classifying snapshots of the doped Hubbard model with machine learning. *Nat. Phys.* **15**, 921–924 (2019).
- Ogata, M. & Shiba, H. Bethe-ansatz wave function, momentum distribution, and spin correlation in the one-dimensional strongly correlated Hubbard model. *Phys. Rev. B* **41**, 2326–2338 (1990).
- Schollwöck, U. The density-matrix renormalization group in the age of matrix product states. *Ann. Phys.* **326**, 96–192 (2011).
- Paeckel, S. et al. Time-evolution methods for matrix-product states. *Ann. Phys.* **411**, 167998 (2019).
- Ferris, A. J. & Vidal, G. Perfect sampling with unitary tensor networks. *Phys. Rev. B* **85**, 165146 (2012).
- Buser, M., Schollwöck, U. & Grusdt, F. Snapshot-based characterization of particle currents and the Hall response in synthetic flux lattices. *Phys. Rev. A* **105**, 033303 (2022).
- Starykh, O. A. & Balents, L. Dimerized phase and transitions in a spatially anisotropic square lattice antiferromagnet. *Phys. Rev. Lett.* **93**, 127202 (2004).
- Mezzacapo, F. Ground-state phase diagram of the quantum  $J_1 - J_2$  model on the square lattice. *Phys. Rev. B* **86**, 045115 (2012).
- Hu, W.-J., Becca, F., Parola, A. & Sorella, S. Direct evidence for a gapless  $Z_2$  spin liquid by frustrating Néel antiferromagnetism. *Phys. Rev. B* **88**, 060402 (2013).
- Wang, L., Poilblanc, D., Gu, Z.-C., Wen, X.-G. & Verstraete, F. Constructing a gapless spin-liquid state for the spin-1/2  $J_1 - J_2$  Heisenberg model on a square lattice. *Phys. Rev. Lett.* **111**, 037202 (2013).
- Gong, S.-S., Zhu, W., Sheng, D. N., Motrunich, O. I. & Fisher, M. P. A. Plaquette ordered phase and quantum phase diagram in the spin-1/2  $J_1 - J_2$  square Heisenberg model. *Phys. Rev. Lett.* **113**, 027201 (2014).
- Jiang, H.-C., Yao, H. & Balents, L. Spin liquid ground state of the spin-1/2 square  $J_1 - J_2$  Heisenberg model. *Phys. Rev. B* **86**, 024424 (2012).
- Haghshenas, R. & Sheng, D. N.  $u(1)$ -symmetric infinite projected entangled-pair states study of the spin-1/2 square  $J_1 - J_2$  Heisenberg model. *Phys. Rev. B* **97**, 174408 (2018).
- Anderson, P. Resonating valence bonds: a new kind of insulator? *Mater. Res. Bull.* **8**, 153–160 (1973).
- Kivelson, S. A., Rokhsar, D. S. & Sethna, J. P. Topology of the resonating valence-bond state: solitons and high- $T_c$  superconductivity. *Phys. Rev. B* **35**, 8865–8868 (1987).
- Anderson, P. W. The resonating valence bond state in  $\text{La}_2\text{CuO}_4$  and superconductivity. *Science* **235**, 1196–1198 (1987).
- Kalmeyer, V. & Laughlin, R. B. Equivalence of the resonating-valence-bond and fractional quantum Hall states. *Phys. Rev. Lett.* **59**, 2095–2098 (1987).
- Kotliar, G. Resonating valence bonds and d-wave superconductivity. *Phys. Rev. B* **37**, 3664–3666 (1988).
- Nagaosa, N. & Lee, P. A. Normal-state properties of the uniform resonating-valence-bond state. *Phys. Rev. Lett.* **64**, 2450–2453 (1990).
- Baskaran, G., Zou, Z. & Anderson, P. The resonating valence bond state and high- $t_c$  superconductivity—a mean field theory. *Solid State Commun.* **88**, 853–856 (1993).
- White, S. R., Noack, R. M. & Scalapino, D. J. Resonating valence bond theory of coupled Heisenberg chains. *Phys. Rev. Lett.* **73**, 886–889 (1994).
- Coll, C. F. Excitation spectrum of the one-dimensional Hubbard model. *Phys. Rev. B* **9**, 2150–2158 (1974).
- Zhang, Y.-H. & Sachdev, S. From the pseudogap metal to the fermi liquid using ancilla qubits. *Phys. Rev. Res.* **2**, 023172 (2020).

65. Nocera, A. & Alvarez, G. Symmetry-conserving purification of quantum states within the density matrix renormalization group. *Phys. Rev. B* **93**, 045137 (2016).
66. Feiguin, A. E. & Fiete, G. A. Spectral properties of a spin-incoherent Luttinger liquid. *Phys. Rev. B* **81**, 075108 (2010).
67. Leifer, M. & Poulin, D. Quantum graphical models and belief propagation. *Ann. Phys.* **323**, 1899–1946 (2008).
68. Hubig, C. et al. The SyTen toolkit. <https://syten.eu> (2015).
69. Hubig, C. Symmetry-protected tensor networks. <https://edoc.ub.uni-muenchen.de/21348/> (2017).

### Acknowledgements

We are thankful for valuable discussions with F. Palm, M. Kebrić, and L. Homeier. This research was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy—EXC-2111—390814868, by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement number 948141)—ERC Starting Grant SimUc-Quam, and by the NSF through a grant for the Institute for Theoretical Atomic, Molecular, and Optical Physics at Harvard University and the Smithsonian Astrophysical Observatory.

### Author contributions

A.B., F.G. and H.S. conceptualized the idea. H.S. carried out the calculations under the supervision of A.B., F.G. and U.S. T.A.H. and I.B. supervised the application to the experimental data. All authors discussed the results. H.S. wrote the manuscript with input from all authors.

### Funding

Open Access funding enabled and organized by Projekt DEAL.

### Competing interests

The authors declare no competing interests.

### Additional information

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s43246-023-00382-3>.

**Correspondence** and requests for materials should be addressed to Henning Schlömer or Annabelle Bohrdt.

**Peer review information** *Communications Materials* thanks Dominik Hangleiter and the other, anonymous, reviewer(s) for their contribution to the peer review of this work. Primary Handling Editor: Aldo Isidori.

**Reprints and permission information** is available at <http://www.nature.com/reprints>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023